

RBSC-NVSS SAMPLE. I. RADIO AND OPTICAL IDENTIFICATIONS OF A COMPLETE SAMPLE OF 1500 BRIGHT X-RAY SOURCES

Franz E. Bauer¹

*National Radio Astronomy Observatory²
Department of Astronomy, University of Virginia,
520 Edgemont Road, Charlottesville, VA 22903
E-mail: fbauer@nrao.edu*

J. J. Condon

*National Radio Astronomy Observatory²
520 Edgemont Road, Charlottesville, VA 22903
E-mail: jcondon@nrao.edu*

Trinh X. Thuan

*Department of Astronomy, University of Virginia,
P.O. Box 3818, Charlottesville, VA 22903-3818
E-mail: txt@virginia.edu*

and

J. J. Broderick

*Department of Physics, Virginia Polytechnic Institute and State University
Blacksburg, Virginia 24061
E-mail: jjb@vt.edu*

ABSTRACT

We cross-identified the *ROSAT* Bright Source Catalog (RBSC) and the NRAO VLA Sky Survey (NVSS) to construct the RBSC-NVSS sample of the brightest X-ray sources (≥ 0.1 counts s⁻¹ $\sim 10^{-12}$ ergs cm⁻² s⁻¹ in the 0.1–2.4 keV band) that are also radio sources ($S \geq 2.5$ mJy at 1.4 GHz) in the 7.8 sr of extragalactic sky with $|b| > 15^\circ$ and $\delta > -40^\circ$. The sky density of NVSS sources is low enough that they can be reliably identified with RBSC sources having rms positional uncertainties $\geq 10''$. We used

¹Visiting Astronomer, Kitt Peak National Optical Observatories, operated by the Association of Universities for Research in Astronomy, Inc., under contract with the National Science Foundation

²The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

the more accurate radio positions to make reliable X-ray/radio/optical identifications down to the POSS plate limits. We obtained optical spectra for many of the bright identifications lacking published redshifts. The resulting X-ray/radio sample is unique in its size ($N \sim 1500$ objects), composition (a mixture of nearly normal galaxies, Seyfert galaxies, quasars, and clusters), and low average redshift ($\langle z \rangle \sim 0.1$).

Subject headings: catalogs — radio continuum: general — surveys — X-rays: general

1. INTRODUCTION

The *ROSAT* All-Sky Survey Bright Source Catalogue (Voges et al. 1999, RBSC revision 1RXS) contains the first large all-sky sample of the brightest X-ray sources, analogous in many respects to the optical NGC catalog. It was derived from the soft (0.1–2.4 keV) X-ray survey performed during the first half year of the *ROSAT* mission in 1990/91. The catalog sky coverage is 92%, and there are 8,547 sources above its 0.1 counts s^{−1} ($\sim 10^{-12}$ ergs s^{−1} cm^{−2}) completeness limit. Bade et al. (1998) found that about one third of the RBSC sources can be reliably identified with galactic stars, while most of the rest are extragalactic. The extragalactic content of the RBSC comprises a diverse blend of (1) normal spiral galaxies whose X-ray emission is powered by stars and stellar remnants, (2) elliptical galaxies with hot gaseous halos, (3) AGN in Seyfert galaxies, elliptical galaxies, quasars, and BL Lac objects, and (4) clusters of galaxies. The large number of sources in this catalog easily permits statistical analyses of *each* type of X-ray object. However, the essential properties of these X-ray sources cannot be determined from the X-ray data alone—we need observations in optical and other wavebands to measure their distances, identify their energy sources, etc. Such observations are possible only for those RBSC sources whose optical counterparts have been identified. In this paper we present reliable radio and optical identifications for sources in the RBSC complete sample.

Most RBSC sources have rms positional uncertainties $\geq 10''$ and the sky density of faint optical objects is high, so only the nearest extragalactic X-ray sources can be optically identified by position coincidence alone (cf. Bade et al. 1998). Fortunately, most extragalactic RBSC sources are also radio sources in the 1.4 GHz NRAO/VLA Sky Survey (Condon et al. 1998a, NVSS), whose sky density is low enough for identification with RBSC sources. Since the radio positions are significantly more accurate, the radio sources may be optically identified, yielding optical identifications for the corresponding X-ray sources as well. The NVSS covers the 10.3 sr of sky north of $\delta = -40^\circ$ and contains over 1.8×10^6 sources stronger than its 2.5 mJy beam^{−1} completeness limit. Since the NVSS was made with relatively low resolution (45'' FWHM), it does not discriminate against moderately extended radio sources in nearby galaxies and clusters. Its rms positional uncertainties range from $< 1''$ for the $N \approx 4 \times 10^5$ sources stronger than 15 mJy to 7'' for the faintest ($S = 2.3$ mJy) detectable sources, allowing us to make optical identifications with objects as faint as $R \approx 21$.

We optically identified the RBSC-NVSS sources with objects in the United States Naval Observatory catalog A2.0 (Monet et al. 1998, USNO). The USNO catalog contains 526,280,881 objects detected by the Precision Measuring Machine on the Palomar Optical Sky Survey I (POSS-I) blue O and red E plates, the UK Science Research Council SRC-J survey plates, and the European Southern Observatory ESO-R survey plates. The catalog was compiled from the blue/red overlaps (within $2''$) of the detection lists generated from scans of POSS-I O and E plates centered on $\delta > -18^\circ$ and SRC-J and ESO-R plates centered on $\delta < -20^\circ$. The stated astrometric and photometric errors are about $0''.25$ and 0.5 mag rms, respectively. The USNO catalog covers the entire sky and probes as deep as $B=21$ (O plates), $R=20$ (E plates), $J=22$, and $F=21$ for objects with appropriate colors.

Section 2 explains our method for making the identifications and assessing their reliabilities. The results are presented in Section 3.

2. Cross-identifications

In this section we present a sequence of increasingly powerful methods for making cross-identifications and directly evaluating the their individual reliabilities: (1) The simplest case is identification by position-coincidence between two wavebands (Sec. 2.1), associating NVSS radio sources with RBSC X-ray sources, for example. For each candidate we derive the probability that it is the correct identification (the reliability of that identification). (2) If the positions are not sufficiently accurate to guarantee reliable identifications by themselves, they may be supplemented by additional data (Sec. 2.2). To continue our example, RBSC X-ray sources have a flatter radio flux-density distribution than NVSS background sources, so radio flux densities affect identification reliabilities. The identification reliabilities derived directly in this section are similar to those obtained via the “likelihood ratio” method by Sutherland & Saunders (1992). (3) Even with the aid of additional data, the RBSC position errors are too large for making reliable X-ray identifications with faint optical objects. We show how accurate positions of NVSS radio sources associated with RBSC sources can be used to select the correct optical counterparts. The reliabilities of such multiwavelength linked position-coincidence identifications are derived in (Sec. 2.3). (4) Finally, linked cross-identifications can themselves be strengthened by applying additional constraints (e.g., optical magnitudes), and their reliabilities are obtained in Section 2.4.

2.1. Direct Position-coincidence Identifications

All identification programs begin with a set of identification candidates in some search area surrounding each source to be identified. This area should encompass all plausible candidates, but its exact size and shape are not critical. A larger than necessary search area increases the number of candidates which must be evaluated but does not significantly degrade identification reliability.

We used search circles of $180''$ in radius (Figure 1) to identify RBSC sources having rms positional uncertainties $\sigma_x \approx 10'' - 20''$. The search radius is much larger than the 3σ X-ray error circle to permit identification with extended and asymmetric radio sources. Each search area contains some numbers $m \geq 0$ of NVSS radio sources and $l \geq 0$ of optical objects from the USNO catalog.

We consider radio identifications of X-ray sources using only the X-ray and radio positions. Let $P(R)$ ($R = 1, \dots, m$) represent the probability that the R^{th} candidate is the correct radio identification of an X-ray source and $P(0)$ the probability that none is. Since there is only a negligible chance that the radio identification is exterior to the search area, $P(0)$ is just the probability that the actual identification is too faint to be recognized as a candidate. For example, we cannot identify radio counterparts fainter than the $2.5 \text{ mJy beam}^{-1}$ NVSS catalog limit. Such an identification is often called an “empty field.” If we make the *astronomical* assumption that not more than one of the m candidates is the correct identification, then the sum of these mutually exclusive probabilities is unity:

$$\sum_{R=0}^m P(R) = 1 . \quad (1)$$

The R^{th} candidate is the correct identification if (1) there exists a detectable identification, (2) that identification lies in the infinitesimal area element dA containing the position of the R^{th} candidate, and (3) the $(m - 1)$ remaining candidates are unrelated sources which happen to lie in the areas dA surrounding their positions (see Figure 1). The probability that all three independent events occur is the product of their individual probabilities, which we now evaluate.

(1) The *a priori* probability that there is a detectable radio identification is equal to the initially unknown fraction f_r of X-ray sources in the sample that actually have detectable radio identifications. We estimated f_r by guessing an initial value, making trial identifications for the whole source sample, replacing the initial value by the observed value, and iterating. If this procedure fails to converge rapidly, the resulting identifications are probably too unreliable to be useful. To estimate the observed value of f_r we summed the computed reliabilities of all the RBSC-NVSS identifications and divided that sum by the total number of X-ray sources. Since only a small percentage of bright galactic X-ray stars are radio sources, we calculated separate f_r values for fields containing optically bright ($m < 12$) stars and for all other fields.

(2) The probability of finding the correct identification in the infinitesimal area dA offset by the angle $\phi_{xr}(R)$ between the position of the X-ray source and the R^{th} candidate is $p[\phi_{xr}(R)]dA$, where $p[\phi_{xr}(R)]$ is just the normalized error distribution of the measured X-ray/radio offsets. These errors include both the radio and X-ray measurement errors and may be augmented by a contribution allowing for possible astronomical offsets of extended sources. For example, the centroid of a head-tail radio source will not coincide with its parent galaxy.

(3) If the R^{th} candidate is the correct identification, then the remaining $(m - 1)$ candidates must be unrelated sources lying in areas dA containing their positions. The probability of finding each unrelated source is $\rho_r dA$, where ρ_r is the mean sky density of unrelated radio candidates. For

NVSS candidates,

$$\rho_r = \int_{S_{\min}}^{\infty} n(S) dS , \quad (2)$$

where $n(S)$ is the differential source count at 1.4 GHz and $S_{\min} = 2.5$ mJy is the minimum flux density of the candidates. Possible clustering of candidates around the true identification could be addressed by increasing ρ_r from its global to local value. If there is no detectable identification, all m candidates must be unrelated sources.

The resulting set of $(m + 1)$ proportionalities

$$P(R) \propto f_r p[\phi_{\text{xr}}(R)] dA (\rho_r dA)^{m-1} \quad \text{if } R > 0 \quad (3a)$$

$$P(0) \propto (1 - f_r) (\rho_r dA)^m \quad (3b)$$

normalized by Equation 1 specify the $(m + 1)$ identification reliabilities:

$$P(R) = c f_r \frac{p[\phi_{\text{xr}}(R)]}{\rho_r} \quad \text{if } R > 0 \quad (4a)$$

$$P(0) = c(1 - f_r) , \quad (4b)$$

where

$$c^{-1} = \frac{f_r}{\rho_r} \sum_{j=1}^m p[\phi_{\text{xr}}(j)] + (1 - f_r) . \quad (4c)$$

Equation 4 gives the probability $P(R)$ that the R^{th} of m candidates is the correct identification and the probability $P(0)$ that none is. These probabilities depend on the search area only through the number m of candidates contributing to the sum in Equation 4c. Making the search area “too big” by definition only adds candidates having negligible $p(\phi_{\text{xr}})$ and hence little effect in Equation 4c.

In many cases, the positional error distributions of both the source and its identification candidates are nearly circular Gaussians. [See Condon et al. (1995) for the more complicated case of elliptical Gaussian error distributions with arbitrary orientations.] If the X-ray source and radio candidate positions have rms uncertainties σ_x and σ_r in each coordinate, then

$$p(\phi_{\text{xr}}) = \frac{1}{2\pi\sigma^2} \exp\left(-\frac{\phi_{\text{xr}}^2}{2\sigma^2}\right) , \quad (5)$$

where $\sigma^2 \equiv \sigma_x^2 + \sigma_r^2$. The variance of ϕ_{xr} is $\langle \phi_{\text{xr}}^2 \rangle = 2\sigma^2$ so a typical identification has $p(\phi_{\text{xr}}) \sim (2\pi\sigma^2 e)^{-1}$ and is highly reliable if both

$$\sigma^2 \ll \frac{f_r}{2\pi e \rho_r (1 - f_r)} \quad (6a)$$

and

$$\sigma^2 \ll \frac{1}{2\pi e \rho_r} \quad (6b)$$

Equation 6a ensures that true empty fields are not misidentified with background sources, a danger if the actual identification rate is very low ($f_r \ll 1$). Equation 6b ensures that there is no difficulty choosing the correct identification from multiple candidates. For NVSS identifications of extragalactic RBSC sources, $f_r \approx 0.61$ and the sky density of sources stronger than the NVSS catalog limit $S_{\min} \approx 2.5$ mJy is $\rho_r \approx 1.76 \times 10^5 \text{ sr}^{-1}$, leading to the fairly weak requirement $\sigma \ll 10^2$ arcsec which is easily satisfied by the RBSC positional uncertainties. The sky density of optical objects in the USNO catalog is two orders of magnitude higher, and the RBSC positional uncertainties are too large to satisfy the requirement $\sigma \ll 10$ arcsec for making optical identifications by position coincidence alone, even though most RBSC sources do have optical counterparts in the USNO catalog.

2.2. Pairwise Identifications With Additional Constraints

Uncertain identifications based on positional coincidence alone may be strengthened or rejected by non-positional data. For example, the flux-density distribution of radio sources identified with extragalactic RBSC sources peaks well above the NVSS sensitivity limit. Figure 2 shows the logarithm of the ratio of the probability $p[S|x]$ that a RBSC-NVSS source has flux density S to the probability $p[S|\bar{x}]$ that an unrelated NVSS sources has flux density S . Most unrelated radio sources are fainter than the correct identifications (indicated in Figure 2 by the positive logarithm of the ratio at all but the faintest radio flux levels), so the stronger of two radio candidates with similar $p(\phi_{\text{xr}})$ is the more likely identification. The reliabilities of such identifications can be calculated through the use of “likelihood ratios” (see Sutherland & Saunders 1992); here we obtain the identification reliabilities directly by extending the derivation of Equation 4.

Let $p[\phi_{\text{xr}}(R), S(R)]dAdS$ be the probability that the R^{th} candidate is the identification lying in the area dA surrounding its observed position and in the flux-density range dS containing its observed flux density $S(R)$. The multiplicative law of probabilities states that this probability is the product of $p[\phi_{\text{xr}}(R)]dA$ and $p[S(R)|x]dS$, where $p(S|x)$ is the normalized ($\int_{S_{\min}}^{\infty} p(S|x)dS = 1$) flux-density distribution of radio sources identified with X-ray sources. Like f_r , $p(S|x)$ is best estimated from the actual identification data by iteration. The probability that the R^{th} candidate is an unrelated (X-ray quiet) radio source with flux density S lying in area dA is $p[S(R)|\bar{x}]\rho_r dAdS$, where $p(S|\bar{x})$ is the flux-density distribution of all (background) NVSS sources. Thus Equations 3 and 4 can be replaced by:

$$P(R) \propto f_r p[\phi_{\text{xr}}(R)] \frac{p[S(R)|x]}{p[S(R)|\bar{x}]} dAdS \left\{ \prod_{j=1}^m p[S(j)|\bar{x}] \right\} (\rho_r dAdS)^{m-1} \quad \text{if } R > 0 \quad (7a)$$

$$P(0) \propto (1 - f_r) \left\{ \prod_{j=1}^m p[S(j)|\bar{x}] \right\} (\rho_r dAdS)^m \quad (7b)$$

and

$$P(R) = c f_r \frac{p[\phi_{xr}(R)]}{\rho_r} \frac{p[S(R)|x]}{p[S(R)|\bar{x}]} \quad \text{if } R > 0 \quad (8a)$$

$$P(0) = c(1 - f_r) \quad (8b)$$

where

$$c^{-1} = \frac{f_r}{\rho_r} \sum_{j=1}^m p[\phi_{xr}(j)] \frac{p[S(j)|x]}{p[S(j)|\bar{x}]} + (1 - f_r) \quad (8c)$$

Equation 8 gives the identification probabilities based on candidate flux densities as well as positional coincidence. It could be further extended to include additional continuous (e.g., spectral index) or discrete (e.g., morphological type) parameters which might prove useful for distinguishing between correct identifications and unrelated candidates.

2.3. Linked Position-coincidence Cross-identifications

The positional uncertainties of most RBSC sources are too large to yield reliable optical identifications with faint galaxies and quasars directly. However, the NVSS sources reliably identified with RBSC sources have sufficiently accurate radio positions that nearly all can be optically identified by position coincidence alone. The reliabilities of such linked X-ray/radio/optical position-coincidence identifications are derived below.

Let $P(R, V)$ be the probability that the R^{th} radio source and the V^{th} optically visible object are the correct identifications of an X-ray source, where $R = 0, 1, \dots, m$ and $V = 0, 1, \dots, l$. The values $R = 0$ and $V = 0$ correspond to radio and optical “empty fields,” respectively. The probabilities of these $(m + 1)(l + 1)$ mutually exclusive possible outcomes must add up to one:

$$\sum_{R=0}^m \sum_{V=0}^l P(R, V) = 1 . \quad (9)$$

Denote the fractions of X-ray sources in the sample having detectable ($R > 0$) radio and optical ($V > 0$) identifications by f_r and f_v , respectively. Let $p[\phi_{xr}(R)]$ and $p[\phi_{xv}(V)]$ be the probability densities of the X-ray/radio and X-ray/optical positional offsets ϕ if the R^{th} radio source and the V^{th} optical object are the correct identifications. The sky densities of background radio and optical candidates are ρ_r and ρ_v , respectively. Using the multiplication law $P(R, V) = P(R)P(V|R)$ yields:

$$P(R, V) \propto f_r p[\phi_{xr}(R)] dA(\rho_r dA)^{m-1} f_v p[\phi_{xv}(V)] dA(\rho_v dA)^{l-1} \quad \text{if } R > 0, V > 0 \quad (10a)$$

$$P(R, 0) \propto f_r p[\phi_{xr}(R)] dA(\rho_r dA)^{m-1} (1 - f_v)(\rho_v dA)^l \quad \text{if } R > 0 \quad (10b)$$

$$P(0, V) \propto (1 - f_r)(\rho_r dA)^m f_v p[\phi_{xv}(V)] dA(\rho_v dA)^{l-1} \quad \text{if } V > 0 \quad (10c)$$

$$P(0, 0) \propto (1 - f_r)(\rho_r dA)^m (1 - f_v)(\rho_v dA)^l \quad (10d)$$

The normalization Equation 9 then implies:

$$P(R, V) = c f_r f_v \frac{p[\phi_{xr}(R)]}{\rho_r} \frac{p[\phi_{xv}(V)]}{\rho_v} \quad \text{if } R > 0, V > 0 \quad (11a)$$

$$P(R, 0) = c f_r (1 - f_v) \frac{p[\phi_{xr}(R)]}{\rho_r} \quad \text{if } R > 0 \quad (11b)$$

$$P(0, V) = c (1 - f_r) f_v \frac{p[\phi_{xv}(V)]}{\rho_v} \quad \text{if } V > 0 \quad (11c)$$

$$P(0, 0) = c (1 - f_r) (1 - f_v) , \quad (11d)$$

where

$$\begin{aligned} c^{-1} &= \frac{f_r f_v}{\rho_r \rho_v} \sum_{j=1}^m p[\phi_{xr}(j)] \sum_{i=1}^l p[\phi_{xv}(i)] \\ &+ \frac{f_r (1 - f_v)}{\rho_r} \sum_{j=1}^m p[\phi_{xr}(j)] + \frac{(1 - f_r) f_v}{\rho_v} \sum_{i=1}^l p[\phi_{xv}(i)] + (1 - f_r) (1 - f_v) . \end{aligned} \quad (11e)$$

Unfortunately, applying Equation 11 does not yield good optical identifications of RBSC X-ray sources because both the X-ray positional uncertainties and the mean density ρ_v of optical candidates are large. Reliable radio identifications of X-ray sources can be made because the density ρ_r of radio candidates is much smaller, and reliable optical identifications of NVSS radio sources are possible because the NVSS positions are more accurate. If the *astronomical* assumption is made that the optical identifications of these radio identifications are also the optical identifications of the corresponding X-ray sources, then

$$\begin{aligned} P(R, V) &\propto f_r p[\phi_{xr}(R)] dA(\rho_r dA)^{m-1} f_v p[\phi_{xv}(V)] dA(\rho_v dA)^{l-1} p[\phi_{rv}(R, V)] dA(\rho_v dA)^{l-1} \\ &\quad \text{if } R > 0, V > 0 \end{aligned} \quad (12a)$$

$$P(R, 0) \propto f_r p[\phi_{xr}(R)] dA (\rho_r dA)^{m-1} (1 - f_v) (\rho_v dA)^l (\rho_v dA)^l \quad \text{if } R > 0 \quad (12b)$$

$$P(0, V) \propto (1 - f_r) (\rho_r dA)^m f_v p[\phi_{xv}(V)] dA (\rho_v dA)^{l-1} (\rho_v dA)^l \quad \text{if } V > 0 \quad (12c)$$

$$P(0, 0) \propto (1 - f_r) (\rho_r dA)^m (1 - f_v) (\rho_v dA)^l (\rho_v dA)^l, \quad (12d)$$

where $p[\phi_{rv}(R, V)]$ is the probability distribution of offsets between the R^{th} radio source and the V^{th} optical object. The normalization Equation 9 implies

$$P(R, V) = c f_r f_v \frac{p[\phi_{xr}(R)]}{\rho_r} \frac{p[\phi_{xv}(V)]}{\rho_v} \frac{p[\phi_{rv}(R, V)]}{\rho_v} \quad \text{if } R > 0, V > 0 \quad (13a)$$

$$P(R, 0) = c f_r (1 - f_v) \frac{p[\phi_{xr}(R)]}{\rho_r} \quad \text{if } R > 0 \quad (13b)$$

$$P(0, V) = c (1 - f_r) f_v \frac{p[\phi_{xv}(V)]}{\rho_v} \quad \text{if } V > 0 \quad (13c)$$

$$P(0, 0) = c (1 - f_r) (1 - f_v), \quad (13d)$$

where

$$c^{-1} = \frac{f_r f_v}{\rho_r \rho_v^2} \sum_{j=1}^m p[\phi_{xr}(j)] \sum_{i=1}^l p[\phi_{xv}(i)] p[\phi_{rv}(j, i)] \\ + \frac{f_r (1 - f_v)}{\rho_r} \sum_{j=1}^m p[\phi_{xr}(j)] + \frac{(1 - f_r) f_v}{\rho_v} \sum_{i=1}^l p[\phi_{xv}(i)] + (1 - f_r) (1 - f_v). \quad (13e)$$

Equation 13 specifies the reliabilities of linked X-ray/radio/optical identifications made on the basis of position-coincidence alone.

2.4. Linked Cross-identifications With Additional Constraints

Finally, differences between the magnitude distributions of the optical counterparts to X-ray sources and background optical objects can be used to improve the reliability of the optical identifications. Figure 3 gives the logarithm of the ratio of the probability $p[\mu|x]$ that the USNO identification of a RBSC-NVSS source has magnitude μ to the probability $p[\mu|\bar{x}]$ that an unrelated USNO object has magnitude μ as a function of B magnitude, and shows, in contrast to the radio, that there is very little difference between the two populations for $B > 13$, but the ratio does become large for objects brighter than this limit. Let $\mu(V)$ denote the magnitude of the V^{th}

optical candidate. Let $p(\mu|x)$ and $p(\mu|\bar{x})$ be the normalized magnitude distributions of optical objects in the USNO catalog which are X-ray sources and unrelated optical objects, respectively. Let $p(S|x)$ be the normalized flux-density distribution of radio-detected X-ray sources and $p(S|\bar{x})$ be the the normalized flux-density distribution of unrelated radio sources stronger than the NVSS limit, 2.5 mJy. Then

$$\begin{aligned}
 P(R, V) &\propto f_r p[\phi_{xr}(R)] \frac{p[S(R)|x]}{p[S(R)|\bar{x}]} dAdS \left\{ \prod_{j=1}^m p[S(j)|\bar{x}] \right\} (\rho_r dAdS)^{m-1} \\
 &\times f_v p[\phi_{xv}(V)] \frac{p[\mu(V)|x]}{p[\mu(V)|\bar{x}]} dAd\mu \left\{ \prod_{i=1}^l p[\mu(i)|\bar{x}] \right\} (\rho_v dAd\mu)^{l-1} \\
 &\times p[\phi_{rv}(R, V)] dA (\rho_v dA)^{l-1} \quad \text{if } R > 0, V > 0
 \end{aligned} \tag{14a}$$

$$\begin{aligned}
 P(R, 0) &\propto f_r p[\phi_{xr}(R)] \frac{p[S(R)|x]}{p[S(R)|\bar{x}]} dAdS \left\{ \prod_{j=1}^m p[S(j)|\bar{x}] \right\} (\rho_r dAdS)^{m-1} \\
 &\times (1 - f_v) \left\{ \prod_{i=1}^l p[\mu(i)|\bar{x}] \right\} (\rho_v dAd\mu)^l (\rho_v dA)^l \quad \text{if } R > 0
 \end{aligned} \tag{14b}$$

$$\begin{aligned}
 P(0, V) &\propto (1 - f_r) \left\{ \prod_{j=1}^m p[S(j)|\bar{x}] \right\} (\rho_r dAdS)^m \\
 &\times f_v p[\phi_{xv}(V)] \frac{p[\mu(V)|x]}{p[\mu(V)|\bar{x}]} dAd\mu \left\{ \prod_{i=1}^l p[\mu(i)|\bar{x}] \right\} (\rho_v dAd\mu)^{l-1} (\rho_v dA)^l \quad \text{if } V > 0
 \end{aligned} \tag{14c}$$

$$\begin{aligned}
 P(0, 0) &\propto (1 - f_r) \left\{ \prod_{j=1}^m p[S(j)|\bar{x}] \right\} (\rho_r dAdS)^m \\
 &\times (1 - f_v) \left\{ \prod_{i=1}^l p[\mu(i)|\bar{x}] \right\} (\rho_v dAd\mu)^l (\rho_v dA)^l.
 \end{aligned} \tag{14d}$$

The normalized probabilities are:

$$P(R, V) = c f_r f_v \frac{p[\phi_{xr}(R)]}{\rho_r} \frac{p[S(R)|x]}{p[S(R)|\bar{x}]} \frac{p[\phi_{xv}(V)]}{\rho_v} \frac{p[\mu(V)|x]}{p[\mu(V)|\bar{x}]} \frac{p[\phi_{rv}(R, V)]}{\rho_v}$$

$$\text{if } R > 0, V > 0 \quad (15a)$$

$$P(R, 0) = cf_r(1 - f_v) \frac{p[\phi_{xr}(R)]}{\rho_r} \frac{p[S(R)|x]}{p[S(R)|\bar{x}]} \quad \text{if } R > 0 \quad (15b)$$

$$P(0, V) = c(1 - f_r)f_v \frac{p[\phi_{xv}(V)]}{\rho_v} \frac{p[\mu(V)|x]}{p[\mu(V)|\bar{x}]} \quad \text{if } V > 0 \quad (15c)$$

$$P(0, 0) = c(1 - f_r)(1 - f_v) , \quad (15d)$$

where

$$\begin{aligned} c^{-1} = & f_r f_v \sum_{j=1}^m \frac{p[\phi_{xr}(j)]}{\rho_r} \frac{p[S(j)|x]}{p[S(j)|\bar{x}]} \sum_{i=1}^l \frac{p[\phi_{xv}(i)]}{\rho_v} \frac{p[\mu(i)|x]}{p[\mu(i)|\bar{x}]} \frac{p[\phi_{rv}(j, i)]}{\rho_v} \\ & + f_r(1 - f_v) \sum_{j=1}^m \frac{p[\phi_{xr}(j)]}{\rho_r} \frac{p[S(j)|x]}{p[S(j)|\bar{x}]} + (1 - f_r)f_v \sum_{i=1}^l \frac{p[\phi_{xv}(i)]}{\rho_v} \frac{p[\mu(i)|x]}{p[\mu(i)|\bar{x}]} \\ & + (1 - f_r)(1 - f_v) . \end{aligned} \quad (15e)$$

It is with Equation 15 that we evaluated the reliabilities of the RBSC identification candidates found in Section 3.

The final identification process proceeded as follows. We started with a list of X-ray positions and errors from the RBSC. We then searched for corresponding NVSS radio and USNO optical sources within a $3'$ radius ($R = 0, \dots, m$ and $V = 0, \dots, l$ respectively). This particular search size was used because it was much larger than the 3σ X-ray error circle and large enough to contain all but the most extended radio and optical sources. The positions of these sources (X-ray, radio, and optical), their positional errors, and their fluxes and magnitudes were combined with the *a priori* probabilities of detecting radio and optical counterparts, f_r and f_v , the RBSC radio flux and optical magnitude distributions using Equation 15 to estimate the probabilities, $P(R, V)$. We estimated f_r and f_v initially, and then iteratively replaced these values with the actual identification rates until they converged to the values of 0.61 and 0.99. The RBSC flux and magnitude distributions were again found iteratively. The radio flux distribution of RBSC sources was compared to the distribution of background NVSS objects and the RBSC optical magnitude distribution was compared to the position dependent optical background distribution given by Bahcall & Soneira (1980).

2.5. Additional Parameters and Possible Concerns

Many astronomical objects have extended X-ray, radio, or optical structures which lead to offsets between wavebands beyond those accounted for by measurement errors.

To allow for an offset between the centroid and central component of a radio source smaller than the NVSS beam, we added one tenth of the deconvolved NVSS size or upper limit in quadrature to the rms uncertainty of the NVSS centroid position. Further, if this RBSC-NVSS matched field happened to lie within the area covered by the VLA FIRST Survey (White et al. 1997, 5'' resolution, 1 mJy flux limit), we then evaluated Equation 15, and replaced the NVSS sources with the FIRST sources if the reliability was better.

An attempt was also made to identify sources resolved into two or more NVSS components, such as double-lobed or head-tail radio galaxies. If two radio components in the search area were within 3' of one another and had a flux ratio < 3 , we initiated a test for a double source. This consisted of adding an artificial “source” with the combined flux of the two components and located at their radio centroid (this gave much better results than the *optimally* weighted centroid used in Windhorst et al. 1984) to the candidate list. We then re-evaluated Equation 15, keeping this new “source” if the reliability was greater than 0.8 or reverting back to the original reliability estimation otherwise. The rms major- and minor-axis position errors for these artificial sources were taken to be $\frac{1}{10}$ and $\frac{1}{30}$ the size of the component separation. We also found a few true double sources with flux ratios > 3 which were evaluated in the same manner.

We have not corrected for extended RBSC sources. This has the effect of selecting against large scale X-ray emission not coincident with optical and radio sources, such as clusters of galaxies.

Since the USNO catalog does not list optical angular sizes, we used the AIPS Gaussian fitting task IMFIT to measure the optical diameters of objects larger than 0'.5. An additional error of $\frac{1}{15}$ of the major axis was added in quadrature with the rms optical positional uncertainty (which we took to be 0'.5).

The USNO O plate magnitudes are converted into Johnson B magnitudes using $B = O - 0.119(O - E)$ (Evans 1988). We found that these USNO magnitudes are systematically too bright for galaxies with $B \lesssim 16$ by as much as three magnitudes in both R and B as shown by a comparison of USNO magnitudes with B magnitudes from the NASA/IPAC Extragalactic Database (NED 1999) (Figure 4). There is better agreement (the dispersion is ~ 2 mag) between USNO and NED magnitudes for galaxies with $B \gtrsim 16$. To avoid overestimating the identification reliabilities of objects brighter than $B \approx 16$, we corrected the USNO blue magnitudes using the polynomial fit

$$B_{\text{NED}} = -0.066 + 1.172B_{\text{USNO}} + 0.120(B_{\text{USNO}})^2 - 0.014(B_{\text{USNO}})^3 + 0.000376(B_{\text{USNO}})^4$$

indicated by the solid curve in Figure 4. The remaining scatter (± 2 mag) is not large enough to impact the reliability of our identifications since the ratio $p(\mu|x)/p(\mu|\overline{x})$ does not change by more than a factor of a few over the affected magnitude range (Fig. 3).

One problem associated with catalogued flux densities and magnitudes is Malmquist bias. Given some error in the flux and magnitude determination and the steep slope of the counts, the true brightness of an object will be overestimated. While this is a potential pitfall for faint flux and magnitude counts, it should not significantly affect the reliabilities calculated here for several reasons: 1) the flux and magnitude estimates are normalized by the same Malmquist biased background, 2) for the optical sources, the flat magnitude distribution (Fig. 3) carries little weight in the reliability determination anyway, and 3) the slope of the optical (and to a lesser extent the radio) RBSC source counts turns over well in advance of the survey limit, such that there are very few faint sources where the bias would be strongest.

A further complication is the high sky density of bright foreground stars at low Galactic latitudes. The likelihood that an optical star is an X-ray emitter is quite large (one third of RBSC sources are stellar in origin; (Bade et al. 1998)), while the likelihood that the same optical star is also a radio emitter is quite small ($\sim 0.1 - 2\%$, see Condon et al. (1997)). Thus a bright X-ray star near a background radio source may result in the identification of the X-ray star with that radio source. Because the density of background optical sources increases more rapidly toward fainter magnitudes than the density of RBSC-NVSS optical identifications does, a bright optical star will be assigned a greater identification reliability than a faint galaxy, even if the star is not quite as close to the radio position as the galaxy is. Weeding out X-ray objects which are likely to be stars will then decrease the chances of calling an interloper the correct identification. Thus we classified all objects brighter than $B < 13$ mag on the basis of appearance on the Digitized Sky Survey (Lasker et al. 1990, DSS) as stars (with diffraction spikes, saturated point source, no diffuse halo) or galaxies. X-ray fields containing one or more bright stars but no bright galaxies within 3σ of the X-ray position were considered stellar. In these cases, the value of f_r was forced to be 0.02 for stellar fields rather than 0.61, the extragalactic identification fraction. We may have missed a few radio stars with high proper motions because the USNO2 and radio positions were measured at different epochs.

Finally, we used the locally measured, rather than global, surface density ρ_v of background objects because many RBSC sources reside in optically overdense regions (clusters of galaxies).

3. Identification Results

For the parent sample, we selected the 5441 RBSC sources above the $0.1 \text{ counts s}^{-1}$ completeness limit with Galactic latitudes $|b| > 15^\circ$ (to reduce the number of objects which might confuse optical identification and minimize extinction at both optical and X-ray wavelengths) and J2000 $\delta > -40^\circ$, the NVSS declination limit. Of these, 1773 sources are readily identified as stars, in agreement with Bade et al. (1998). We attempted to make *linked* X-ray/radio/optical cross-identifications using X-ray/radio and radio/optical positional coincidence supplemented by radio flux densities and optical magnitudes, as described in Section 2.4.

We visually confirmed each match by overlaying the contours of the NVSS image and the RBSC error box on the DSS (see Figure 5). In addition, the IRAS Faint Source Catalogue (Moshir et al. 1992, FSC) has been matched against the RBSC (see Condon et al. 1998b), so we have plotted the FSC 3σ error ellipses to aid identification and differentiate between physical emission processes. The nearest candidate usually has the highest calculated reliability (Equation 15), although the radio emission for some sources is extended and more care had to be taken in locating an optical counterpart. We use the term *association* to describe extended X-ray/radio sources which appear related but may not be spatially coincident. For instance, a cluster of galaxies may contain one or more radio galaxies and hot intercluster X-ray gas. We retained 19 identifications with low calculated reliabilities on the basis of other information, such as detection in a *ROSAT* pointed observation or no obvious optical counterpart within the error circle. All deviations from the standard reliability estimate are noted in Section 3.1. The reliabilities of these sources are given as ‘–’.

Some RBSC-NVSS sources had no USNO sources listed near their NVSS positions, but objects were clearly visible on the DSS (i.e the red plate). Most were either very bright, extended galaxies or very faint objects. USNO sources were identified based on a positional coincidence of within $2''.0$ on the red and blue plates, so we might expect that some bright galaxies will have red and blue emission peaks separated by more than $2''.0$ and thus be rejected. For these we used positions and magnitudes from NED. Furthermore, the faint objects we find must only appear on the red plate (the lack of a blue plate counterpart presumably excluded them from the USNO catalog). We used the AIPS task IMFIT to measure their positions. Faint object magnitudes were estimated by comparison with nearby USNO sources.

These additional sources were added to the USNO list of optical candidates located within the search area and evaluated as part of the standard reliability calculation (see Equation 15). This additional step introduces a slight bias in the reliabilities of these new faint sources, because we have only added the faint sources in close proximity to the radio/X-ray positions, and not those from the rest of the $3'$ radius field. We can test the statistical error of this effect by simply doubling the number of local background objects; this is akin to probing roughly 2 magnitudes fainter than the USNO. While the change will depend on the exact field, we found the reliability with this background enhancement typically differed by only 1% (and at most 5%).

The Galactic absorption corrected X-ray fluxes come from the RBSC correlation catalogue (Voges et al. 1999), in which the RBSC count rate was converted to flux assuming a power law photon energy distribution of the form $E^{-\Gamma_x+1}$, common to AGN and clusters of galaxies. An average photon index $\langle\Gamma_x\rangle = 2.3$ was used, typical of extragalactic objects (Hasinger, Trümper & Schmidt 1991) over the *ROSAT* energy range. Corrections for Galactic absorption were based on hydrogen column densities N_H obtained from Dickey & Lockman (1990).

The error of the unabsorbed X-ray flux is about 30% taking into account errors in photon statistics and variable Galactic absorption. Flux estimates break down for cases when the actual

Γ_x deviates significantly from the mean value, there is additional intrinsic absorption, or when a single power law model inadequately describes the spectral shape (see Brinkmann et al. (1995) for a more detailed discussion of this problem).

A total of 1556 RBSC-NVSS sources were identified, on the criterion that the sum total of all radio identification reliabilities be greater than 0.50. The vast majority exceeded this cut-off limit a large margin. The distribution of RBSC-NVSS sources on the sky is fairly uniform (Figure 6). Roughly one half of the extragalactic RBSC sources did not have radio counterparts. Only 1 RBSC-NVSS source has no detectable optical counterpart (not surprising given the large RBSC error circles and the density of optical sources at the limiting magnitude of the POSS). Table 1 lists the 44 possible galactic stars so identified because they have $B < 13$ mag, diffraction spikes, faint irregular radio contours or a spectral type from SIMBAD. Many have been independently verified with high resolution VLA images to be radio stars by Condon et al. (1997). Table 1 is arranged in the following manner:

Column 1. *ROSAT* BSC name, format 1RXS JHHMMSS.S+DDMMSS.

Column 2. SIMBAD Common Name, if available.

Column 3. *ROSAT* RBSC X-ray Flux.

Column 4. NVSS position, unless otherwise noted.

Column 5. NVSS 1.4 GHz Flux density.

Column 6. Identification Probability $P(R,V)$ (Equation 15). Identifications with a ‘–’ have been accepted despite having a low probability.

Column 7. SIMBAD object type, if available. Abbreviations from SIMBAD as follows:

*	=	Star
Al	=	Eclipsing binary of Algol type
bL	=	Eclipsing binary of Beta Lyr type
Cl	=	Star or globular cluster
DQ	=	Cataclysmic variable of DQ Her type
Fl	=	Flare star
Mi	=	Variable star of Mira Cet type
Pu	=	Pulsating variable star
Ro	=	Rotationally variable star
RS	=	Star of RS CVn type
sr	=	Semi-regular pulsating star
TT	=	T Tau-type star
V	=	Variable star
PN	=	Planetary nebula

Column 8. SIMBAD B magnitude, if available; otherwise USNO magnitude enclosed in parentheses.

Column 9. SIMBAD spectral type, if available.

The remaining 1512 objects are extragalactic and listed in Table 2. Figure 7 shows the cumulative fraction of identifications versus reliability. The vast majority ($\sim 70\%$) of identifications have reliabilities greater than 99%. Table 2 is arranged in the following manner:

Column 1. *ROSAT* RBSC name, format 1RXS JHHMMSS.S+DDMMSS.

Column 2. NED Common Name, if available.

Column 3. *ROSAT* BSC X-ray Flux.

Column 4. NVSS position, unless noted otherwise.

Column 5. NVSS 1.4 GHz Flux density.

Column 6. Identification Probability $P(R,V)$ (Equation 15). Identifications with a ‘–’ have been accepted despite having a low probability.

Column 7. B magnitude from NED, if available; otherwise from USNO as indicated by ‘()’. Identifications with a ‘–’ denote empty fields.

Column 8. Redshift, if known. From NED unless noted otherwise. Identifications with a ‘–’ represent objects for which spectra was taken but a redshift could not be determined. We have obtained optical spectra with the KPNO 2.1 m telescope for many of the bright identifications which did not possess published redshifts. The detailed description of these spectra will be given in a subsequent paper (Bauer, Thuan & Condon 2000).

Column 9. NED optical morphology, if available.

Column 10. Spectral classification, if known. From NED unless noted otherwise in redshift column.

AGN	=	Active galactic nuclei, type unknown
Blazar	=	BL Lacertae type object with variable emission line spectra
BLLAC	=	BL Lacertae type object
BLRG	=	Broad-line radio galaxy
cD	=	Central dominant galaxy, Early-type stellar absorption continuum
Early	=	Early-type stellar absorption continuum
HII	=	Starburst galaxy with narrow emission-line spectra similar to HII regions
HPQ	=	High polarization QSO
LINER	=	Low ionization narrow emission-line region
LPQ	=	Low polarization QSO
QSO	=	Quasi-stellar object
Sy1-2	=	Seyfert galaxy, emission-line spectra classified from type 1 to 2

There are a small number of cases having low reliabilities, but which may in fact be true identifications. They generally fall into two categories; objects needing higher radio resolution and candidates that have optical/radio matches farther than 3σ from the X–ray source which did not have a convincing optical match. Many of the latter are likely to be associated with nearby clusters.

3.1. Notes on Individual Identifications

1RXS J000312.3–355541: Member of cluster Abell 2717.

1RXS J001031.3+105832: Member of triple galaxy III Zw 002.

1RXS J001118.5–285126: Member of cluster Abell 2734. Abell 2744. Diffuse radio emission; possible radio halo?

1RXS J001823.8+300357: Member of group WP 01.

1RXS J001906.0–202638: Cluster Abell S0026 is $50''$ from ID and overlaps within positional errors.

1RXS J002041.8–254307: Member of cluster Abell 0022.

1RXS J002136.8+280305: Group IV Zw 015 is $77''$ from ID and overlaps within positional errors.

1RXS J002430.5–292856: Extended radio emission.

1RXS J002534.9–330255: Greater than 3σ from X–ray position. Low X–ray flux and large error region imply unreliable

X-ray position. Optical and radio information point towards a correlation. Member of cluster ABELL S0041.

1RXS J002811.6+310342: Possible ID is not at 002812.61+310315.3 (as identified in RGB). Better radio resolution would strengthen ID.

1RXS J003037.6–241053: Member of cluster Abell 0047.

1RXS J003413.7–212619: Member of group SCG 05.

1RXS J004324.9–203728: Member of cluster Abell 2813.

1RXS J004924.0–293128: Member of cluster Abell S0084.

1RXS J010721.2+322254: Greater than 3σ from X-ray position. Low X-ray flux and large error region imply unreliable X-ray position. Optical and radio information point towards a correlation. Member of group IV Zw 038.

1RXS J010750.4–364342: Member of cluster Abell 2871.

1RXS J010850.5–152438: Member of cluster Abell 0151.

1RXS J011301.1+153041: Member of cluster Abell 0160. Given the low X-ray flux and large error region imply unreliable X-ray position. Optical and radio information point towards a correlation.

1RXS J011354.9–314538: Greater than 3σ from X-ray position. Given the low X-ray flux and large error region imply unreliable X-ray position. Optical and radio information point towards a correlation. Pointed *ASCA* observation by Arimoto et al. 1997 confirms ID. Member of cluster Abell S0141.

1RXS J011513.2+405743: More reliable ID is a blend of star and galaxy on DSS.

1RXS J012058.3–135100: Member of cluster CID 10.

1RXS J012337.2+331506: Member of pair ARP 229.

1RXS J012507.3+084124: Member of cluster Abell 0193.

1RXS J012910.8–214158: Galaxy composed of two compact optical cores.

1RXS J013152.8–133651: Member of cluster Abell 0209.

1RXS J013632.9+390556: KPNO spectrum obtained. No features revealed amid strong blue continuum.

1RXS J013715.1–091145: Member of cluster CID 12.

1RXS J015244.7+360855: Member of cluster Abell 0262.

1RXS J015705.3+412029: Member of cluster Abell 0276.

1RXS J020144.2–021158: Member of cluster Abell 0291.

1RXS J023727.6–263027: Member of cluster Abell 0368.

1RXS J024122.1–283919: Cluster Abell 3041 is $53''$ from ID and overlaps within positional errors.

1RXS J024620.0–301639: Supernova SN 1992bd is $1''$ from ID and is also a valid identification.

1RXS J024937.1–311114: Member of cluster Abell S0301.

1RXS J030150.6+355003: Member of group UGC 02489 and cluster Abell 0407. V Zw 311 NOTES02 appears to be a cD from DSS and is likely to be true ID. Better radio resolution needed to clarify.

1RXS J031611.4+090445: KPNO spectrum obtained. No features revealed amid the flat continuum.

1RXS J033828.8–352701: Member of Fornax cluster.

1RXS J041325.4+102756: Member of cluster Abell 0478.

1RXS J042551.3–083329: Member of cluster EXO 0422–086.

1RXS J042700.9–204321: Member of cluster Abell 0490.

1RXS J043337.7–131528: Member of cluster Abell 0496.

1RXS J043643.8+100323: Member of cluster MS 0433.9+0957.

1RXS J043902.0+052050: Member of cluster RX J04390+0520.

1RXS J044511.8–155118: Member of cluster CID 25.

1RXS J044803.3–202741: High resolution radio position from Owen, White & Ge (1993) confirms ID. Member of cluster Abell 0514.

1RXS J044814.8+095259: Member of cluster RX J0448.2+0952.

1RXS J045859.6–002904: Member of cluster CID 27.

1RXS J050049.4–384038: Member of cluster Abell 3301.

1RXS J050120.1–033223: High resolution radio position from Owen, White & Ge (1993) confirms ID. Member of cluster Abell 0531.

1RXS J050507.9–322007: Member of cluster Abell 3313.

1RXS J051611.5–000807: RBSC position incorrect. X-ray observations from Ceballos & Barcons (1996) confirm ID.

1RXS J054738.7–315228: Member of cluster Abell 3364.

1RXS J055040.7–321619: Cluster Abell S0549 ($z = 0.04030$) is $13''$ from ID and overlaps well within positional errors.

1RXS J055711.1–372813: Member of cluster Abell S0555.

1RXS J060552.7–351759: Member of cluster Abell 3378.

1RXS J062707.5–352917: Member of cluster Abell 3392.

1RXS J064326.5+421420: High resolution radio position from Laurent-Muehleisen et al. (1997) confirms ID.

1RXS J070427.0+631856: Member of cluster Abell 0566.

- 1RXS J070907.6+483657:** Member of cluster Abell 0569.
- 1RXS J071006.0+500243:** High resolution radio position from Laurent-Muehleisen et al. (1997).
- 1RXS J073221.5+313750:** Member of cluster Abell 0586.
- 1RXS J074144.8+741444:** Member of cluster ZwCl 0735.7+7421.
- 1RXS J074238.0+092213:** Member of cluster Abell 0592. Better radio resolution needed.
- 1RXS J081021.3+421657:** Member of cluster [VMF98] 047.
- 1RXS J081112.3+700300:** Member of cluster Abell 0621.
- 1RXS J081929.5+704221:** Very extended radio and optical emission associated with the dwarf galaxy Holmberg II.
- 1RXS J082209.5+470601:** Member of cluster Abell 0646.
- 1RXS J083811.0+245336:** Member of pair CGCG 120–011.
- 1RXS J083950.7–121424:** High resolution radio position from Li & Jin (1996) confirm ID as [HB89] 0837-120 and not [EYG89] 087.
- 1RXS J084255.9+292752:** Member of cluster ZwCl 0839.9+2937.
- 1RXS J091037.2+332920:** KPNO spectrum obtained. No features revealed among strong blue continuum.
- 1RXS J091949.1+334532:** X-ray position offset, but Einstein observations from Fabbiano, Kim & Trinchieri (1992) confirm ID. Member of cluster Abell 0779.
- 1RXS J092406.1+141002:** Member of cluster Abell 0795.
- 1RXS J094740.4–305655:** Member of triple galaxy AM 0945–304.
- 1RXS J095821.8–110344:** Member of cluster Abell 0907.
- 1RXS J100121.5+555351:** Member of cluster [YGK80] 0957+561CLUSTER. QSO [HB89] 0957+561 and galaxy [YGK81] 097 are 4'' and 9'' from ID and overlap within positional errors.
- 1RXS J100639.1+255450:** Member of cluster Abell 0923.
- 1RXS J101335.9–135108:** Member of cluster RX J10136–1350.
- 1RXS J101912.1+635802:** Member of pair CGCG 313–011.
- 1RXS J102228.9+500630:** Member of cluster Abell 0980.
- 1RXS J102339.3+041117:** Member of cluster ZwCl 1021.0+0426.
- 1RXS J102350.7–271522:** Member of cluster Abell 3444.
- 1RXS J102758.9–064804:** Member of cluster RX J1027.9-0648.
- 1RXS J103155.8–141659:** Multiple redshift systems overlap within positional errors.
- 1RXS J103459.5+304138:** Member of cluster Abell 1045.
- 1RXS J104043.7+395706:** Member of cluster Abell 1068.
- 1RXS J104431.7–070404:** Member of cluster Abell 1084.
- 1RXS J104651.9–253546:** Member of cluster RX J10468–2535.
- 1RXS J105344.2+492956:** Member of cluster MS 1050.7+4946.
- 1RXS J105825.9+564716:** Member of cluster Abell 1132.
- 1RXS J111137.2+405031:** Member of cluster Abell 1190.
- 1RXS J111422.6+582318:** High resolution radio position from Laurent-Muehleisen et al. (1997) confirms ID.
- 1RXS J111450.1–121351:** Cluster Abell 1211 is 89'' from ID and overlaps within positional errors.
- 1RXS J113121.4+333447:** Member of cluster Abell 1290.
- 1RXS J113153.7–195543:** Member of cluster Abell 1300.
- 1RXS J113222.4+555828:** Galaxy MCG +09–19–110 ($z = 0.05130$) is 17'' from ID and overlaps within positional errors. Member of cluster RX J1132.3+5558.
- 1RXS J113448.4+490438:** Member of cluster Abell 1314.
- 1RXS J114124.2–121632:** Member of cluster RX J1141.4–1216.
- 1RXS J114442.2+672435:** Member of cluster Abell 1366. Optical and radio information point towards a correlation. Possible X-ray cluster emission?
- 1RXS J114452.7+194706:** Greater than 3σ from RBSC position. Public *ROSAT* HRI observations place X–ray source much closer to reliable ID. Association with cluster Abell 1367.
- 1RXS J114947.0–121845:** Member of Abell 1391. Probable association.
- 1RXS J115518.9+232431:** Member of cluster Abell 1413. taken from the FIRST 1.4 GHz survey (White et al. 1997). NVSS flux density is 5.4 mJy. Member of cluster RX J11570+2415.
- 1RXS J115719.0+333645:** High resolution radio position from Owen, White & Ge (1993) confirms more reliable ID. Member of cluster Abell 1423.
- 1RXS J120511.7+392043:** Member of cluster RX J1205.1+3920
- 1RXS J121105.4+352005:** Member of cluster RX J1211.0+3520
- 1RXS J122752.7+632317:** Member of cluster RX J1227.8+6323.
- 1RXS J122945.9+075927:** Member of cluster Virgo S.
- 1RXS J123658.8+631111:** Member of cluster Abell 1576.

1RXS J124349.3–153320: Member of cluster Abell 1603.
1RXS J125233.7–311605: Member of cluster RX J12525–3116.
1RXS J125422.2–290034: Member of cluster Abell 3528.
1RXS J125710.0–172345: Pointed *ROSAT* observation from Peres et al. (1998) confirms ID. Member of cluster Abell 1644.
1RXS J125921.5–041131: Member of cluster Abell 1651.
1RXS J130250.3–023041: Member of cluster Abell 1663.
1RXS J130343.6–241506: Member of cluster Abell 1664.
1RXS J130346.7+191635: Member of cluster Abell 1668.
1RXS J130552.6+305405: Member of cluster Abell 1677.
1RXS J130916.1–013658: Member of cluster MS 1306.7–0121.
1RXS J131129.5–012017: Member of cluster Abell 1689.
1RXS J131506.8+514931: Member of cluster Abell 1703.
1RXS J132016.3+330828: Member of cluster RX J1320.1+3308.
1RXS J132542.1–022800: No optical counterpart on DSS.
1RXS J132549.2+591937: Member of cluster Abell 1744.
1RXS J132617.4+001329: Member of cluster RX J1326.3+0013.
1RXS J132758.9–313027: Greater than 3σ from RBSC position. Optical and radio information point towards a correlation. Member of cluster Abell 3558. Possible X-ray blend of star and galaxy?
1RXS J133226.0–330812: Better radio resolution confirms ID.
1RXS J134104.8+395942: Member of cluster Abell 1774.
1RXS J134152.6+262230: Cluster Abell 1775 ($z = 0.06960$) is $72''$ from ID and overlaps within positional errors.
1RXS J134730.5–114455: Member of cluster RX J13475–1145.
1RXS J134852.6+263541: Member of cluster Abell 1795.
1RXS J140102.1+025249: Member of cluster Abell 1835.
1RXS J140337.0–335840: Member of cluster Abell S0753.
1RXS J140728.4–270055: Member of cluster Abell 3581.
1RXS J141342.6+433938: Member of cluster Abell 1885.
1RXS J141357.3+711751: Association with cluster Abell 1895 ($z = 0.22492$).
1RXS J142139.7+371743: Member of cluster Abell 1902.
1RXS J143236.0+313855: Cluster Abell 1930 ($z = 0.13130$) is $119''$ from ID and overlaps within positional errors.
1RXS J143527.9+550747: Member of cluster Abell 1940.
1RXS J144428.4+311304: Cluster Abell 1961 ($z = 0.23200$) is $117''$ from ID and overlaps within positional errors.
1RXS J145307.8+215333: Member of cluster Abell 1986.
1RXS J145431.4+183834: Member of cluster Abell 1991.
1RXS J145434.1+080250: Greater than 3σ from RBSC position. Bade et al. (1998) claim optical—radio ID is the correct X-ray source. Optical and radio information point towards a correlation.
1RXS J145507.9+192025: High resolution radio position from Laurent-Muehleisen et al. (1997) confirms ID.
1RXS J145715.4+222026: Member of cluster MS 1455.0+2232.
1RXS J145904.1–084254: Better radio resolution needed to confirm ID.
1RXS J150020.7+212213: Member of cluster Abell 2009.
1RXS J151056.3+054431: Member of cluster Abell 2029.
1RXS J151127.2+062153: Greater than 3σ from RBSC position. Optical and radio information point towards a correlation. Member of cluster Abell 2033. Possible X-ray cluster emission?
1RXS J151642.4+070058: Member of cluster Abell 2052.
1RXS J151845.3+061340: High resolution radio position from Laurent-Muehleisen et al. (1997) confirms ID. Member of cluster Abell 2055.
1RXS J152151.0+074221: Member of cluster MKW 03s.
1RXS J152305.7+083550: ID greater than 3σ of RBSC position. There is a strong optical/radio candidate (CGCG 077-097) located within the cluster Abell 2063.
1RXS J153950.3+304305: Member of cluster Abell 2110.
1RXS J154009.4+141116: High resolution radio position from Laurent-Muehleisen et al. (1997) confirms ID (a star?).
1RXS J155611.0+662123: Association with cluster Abell 2146.
1RXS J160435.1+174333: High resolution X-ray position from Burstein et al. (1997) confirms ID.
1RXS J160456.8+235604: Member of cluster AWM 4.
1RXS J160740.7+254106: Optical ID is offset from strong radio source (FIRST ID), possibly indicative of a misidentification with a foreground star.
1RXS J161546.9–060841: Radio emission is very diffuse, from multiple components. Better radio resolution needed. Association.

1RXS J161711.4+063816: No object found within 3σ of X-ray source – X-ray errors too conservative? Identified as RGB J1617+066. Optical and radio information point towards a correlation. High resolution radio position from Laurent-Muehleisen et al. (1997).

1RXS J162032.0+295321: Member of cluster Abell 2175.

1RXS J162100.4+254547: Member of Abell 2177.

1RXS J162837.7+393249: Member of cluster Abell 2199.

1RXS J163246.8+053423: Member of cluster Abell 2204.

1RXS J164022.1+464215: Member of cluster Abell 2219.

1RXS J164238.9+272621: Member of cluster Abell 2223.

1RXS J170242.5+340336: Member of cluster Abell 2244.

1RXS J170941.2+342529: Member of cluster Abell 2249.

1RXS J171519.5+572430: Member of cluster CID 71.

1RXS J171706.8+293117: Member of cluster RBS 1634.

1RXS J171718.9+422652: Member of cluster RX J17173+4227.

1RXS J171746.9+194057: Association with cluster Abell 2254.

1RXS J171810.9+563955: Member of cluster ZwCl 1717.9+5636.

1RXS J172009.3+263727: Member of cluster RX J17201+2637.

1RXS J172226.7+320752: Member of cluster Abell 2261.

1RXS J173257.0+403635: Association with cluster Abell 2272.

1RXS J173301.7+434533: Member of cluster CID 72.

1RXS J174414.3+325925: Member of cluster ZwCl 1742.1+3306.

1RXS J175004.4+470037: High resolution radio position from Laurent-Muehleisen et al. (1997) confirms ID.

1RXS J175441.9+680334: Cluster ZwCl 1754.5+6807 ($z = 0.07700$) is $53''$ from ID and overlaps within positional errors.

1RXS J175706.9+535130: Member of cluster Abell 2292.

1RXS J182157.4+642051: High resolution radio position from Laurent-Muehleisen et al. (1997) confirms ID.

1RXS J203445.3–354921: Member of cluster Abell 3695.

1RXS J210027.7–170913: Better radio resolution needed to confirm ID.

1RXS J210707.7–252643: Cluster Abell 3744 is nearby and provides a considerable amount of diffuse radio emission as well. Better radio resolution needed.

1RXS J212706.7–120927: Greater than 3σ of RBSC position. Optical and radio information point towards a correlation.

1RXS J213414.4–311737: Better radio resolution needed to confirm ID.

1RXS J213504.7+085807: Optical ID is actually a composite of a bright star and another fainter source.

1RXS J214015.3–233946: Member of cluster MS 2137.3–2353.

1RXS J215336.1+174111: Member of cluster Abell 2390.

1RXS J215541.5+123144: Member of cluster Abell 2396.

1RXS J221020.8–121040: Member of cluster Abell 2420.

1RXS J221614.9–092033: Greater than 3σ of RBSC position. Optical and radio information point towards a correlation. Possible X-ray cluster emission?

1RXS J224919.5–372724: Member of cluster Abell S1065.

1RXS J225019.2+105406: Member of cluster Abell 2495.

1RXS J225334.1–334309: Member of cluster Abell 3934.

1RXS J230714.9–151322: Member of cluster Abell 2533.

1RXS J231607.5–202739: Member of cluster Abell 2566.

1RXS J231712.8+184237: Member of group HCG 094.

1RXS J231829.5+184246: Greater than 3σ of RBSC position. Optical and radio information point towards a correlation. Member of cluster Abell 2572. Possible X-ray cluster emission?

1RXS J232125.9–231230: Member of cluster Abell 2580.

1RXS J232519.4–120741: Member of cluster Abell 2597.

1RXS J233631.0+210848: Member of cluster Abell 2626.

1RXS J233641.8+235526: Member of cluster Abell 2627.

1RXS J233827.6+270045: Greater than 3σ of RBSC position. Optical and radio information point towards a correlation. Member of cluster Abell 2634. Pointed *ROSAT* HRI observations by Sakelliou & Merrifield 1998 show X-ray emission is a blend of cluster and individual galaxy emission.

J234741.2–280829: X-ray emission appears to be from bright knot in radio jet. Optical and radio information point towards a correlation.

1RXS J235034.7+292924: Cluster MS 2348.0+2913 ($z = 0.09500$) is $22''$ from ID and overlaps within positional errors. KPNO spectrum obtained. Strong absorption features at $z = -0.00020$, possibly a foreground star amid the cluster. Better optical and radio resolution needed to clarify.

1RXS J234741.2–280829: ID greater than 3σ of RBSC position. There is a strong optical/radio candidate nearby. Member of cluster Abell 4038.

3.2. Previous RBSC Misidentifications?

1RXS J000856.1+411034: Better radio resolution needed. Possible X-ray ID does not seem to be B3 0006+408 (as identified in RGB), but rather a bright, diffuse object at 00h08m56.0s +41d10m09.1s.

1RXS J001144.5+322441: Radio ID is 87GB 000913.4+320913 (as identified in RGB), and X-ray ID is cluster Abell 0007.

1RXS J220220.8+035306: Optical/X-ray ID greater than 3σ from radio position (NVSS and RGB). Background radio object misidentified with B=13.2 X-ray star?

3.3. A simple, well-defined sample of AGN

The RBSC-NVSS sample is defined by the following criteria:

- RBSC count rate ≥ 0.1 counts s^{-1}
- $f_r \geq 2.5mJy$
- $\delta \geq -40^\circ$
- $|b| \geq 15^\circ$

Columns 1 and 2 of Table 3 summarize the results of the selection process. We should note here, that while the method we employ here is quite robust, the large number of RBSC sources and the loose positional errors dictate that there will be some spurious spatial coincidences. To estimate the number of false matches we might have, we shifted the X-ray positions by $8'$ and performed the identification procedure again. The results are listed in column 3 of Table 3, where we find that the expected percentage of spurious matches is around 3% for reliabilities above 0.50. This fraction could be further reduced by comparing the flux density and magnitude distributions of the true and spurious objects; most spurious sources lie near the NVSS and USNO survey limits, while RBSC-NVSS sources do not.

Forthcoming papers in this series will present the radio, optical, and X-ray properties of the sample and optical identifications and classifications of a subset of objects. Using a number of criteria, we find that this sample of 1512 extragalactic objects is comprised almost entirely of AGN, making this the largest, complete sample of its kind. It represents a major step forward in the identification of RBSC objects and contains a large sample of both radio-loud *and* radio-quiet X-ray objects (previous surveys of this type have typically sampled only the radio-loud population).

FEB thanks W. Brinkmann and J. Siebert for their helpful comments and assistance with the RBSC. He also acknowledges support from a National Radio Astronomy Observatory Jansky Pre-doctoral Fellowship. JJB acknowledges support from the NSF through grant AST 9320547. This research has made use of data obtained through the High Energy Astrophysics Science Archive Research Center Online Service, provided by the NASA/Goddard Space Flight Center, the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, Caltech, under contract with the National Aeronautics and Space Administration and the SIMBAD database, operated at CDS, Strasbourg, France. The Digitized Sky Surveys were produced at the Space Telescope Science Institute under US government grant NAG W-2166.

Table 1. RBSC Galactic Identifications

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)
<i>ROSAT</i> BSC	SIMBAD	S _x	Position		S _{1.4GHz}	ID	SIMBAD		
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	(J2000) δ	mJy	P	Type	m _B	Sp. Class.
J001208.2–155031	SAO 147143	1.52e–11	00 12 07.78	–15 50 38.6	2.6	0.94	*	10.5	K0
J004703.1–115217	NGC 0246	1.01e–11	00 47 03.41	–11 52 19.1 ^O	126.7 ^E	1.00	PN	15.8	
J020135.5–160957	HD 12538	1.60e–12	02 01 35.23	–16 10 19.4	3.2	1.00	*	7.1	A3
J031634.9+321115	HD 20277	2.75e–11	03 16 34.00	+32 11 15.9	2.8	0.98	*sr	7.0	G8IV
J032635.1+284302	UX Ari	2.65e–10	03 26 35.40	+28 42 54.2	41.6	1.00	*RS	6.5	
J033647.2+003518	RGB J0336+005	7.10e–10	03 36 47.25	+00 35 15.5	123.2	1.00	*RS	5.7	G9
J040940.8–075327	V* EI Eri	1.34e–10	04 09 40.92	–07 53 31.7	3.9	1.00	*RS	7.7	G5IV
J041411.9+281230	HD 283447	5.86e–12	04 14 13.12	+28 12 14.8	6.0	1.00	*TT	11.8	K2
J052507.7+062103	SV* ZI 374	9.87e–12	05 25 08.18	+06 20 50.6	3.0	0.99	*V	1.4	B2III
J053516.6–052320	M42	1.30e–10	05 35 17.17	–05 22 32.7	105702.0	0.97	*Cl	3.0	III
J062150.0–341140	V* HY CMa	3.57e–12	06 21 49.55	–34 11 54.1	8.6	1.00	*V	10.3	K0
J072132.9+260939	V* V340 Gem	1.01e–11	07 21 32.89	+26 09 27.6	7.5	0.95	*sr	8.4	G2V
J072931.4+355607		6.47e–12	07 29 31.76	+35 55 51.6	7.3	0.99	*	(12.9)	
J074318.7+285306	HD 62044	2.15e–10	07 43 18.53	+28 53 02.5	6.5	0.98	*RS	4.3	K1III
J093346.5+624943	V* FF UMa	4.49e–11	09 33 46.94	+62 49 40.0	7.4	1.00	*V	8.9	G5
J103105.7+823327	NSV 4864	7.81e–12	10 31 05.02	+82 33 30.7 ^O	9.6	0.93	*V	5.6	F2V
J104520.5+453404	V* TX UMa	2.26e–12	10 45 20.50	+45 33 58.7 ^O	9.6	0.71	*Al	7.1	B8V
J113155.7–343632	CD–33 7795	1.93e–11	11 31 55.14	–34 36 29.4	5.6	0.97	*	(11.8)	M1
J115800.5+140222	V* FZ Leo	1.45e–11	11 58 02.16	+14 02 18.5	2.9	0.95	*V	8.9	G5
J133447.5+371100	V* BH CVn	2.78e–11	13 34 47.62	+37 10 53.8	8.8	0.91	*RS	5.3	F2IV
J142555.6+141148	StKM 1–1155	5.16e–12	14 25 56.00	+14 12 14.0 ^O	2.8	0.67	*	11.4	K4
			14 25 56.16	+14 12 02.1 ^O	2.8	0.26		(19.1)	
J143638.2+584309		1.41e–12	14 36 36.86	+58 42 55.8	4.5	1.00	*	(11.1)	
J150058.0–083103	V* del Lib	2.22e–11	15 00 58.38	–08 31 10.2	4.8	0.98	*Al	4.9	B9.5V
J151720.9–190103	HD 135743	1.39e–11	15 17 21.22	–19 00 58.9 ^O	3.3	0.93	*	10.0	G3/G5V
			15 17 21.26	–19 01 12.1 ^O	3.3	0.07		(17.2)	
J152153.0+205830	V* OT Ser	2.65e–11	15 21 52.98	+20 58 38.1	15.4	0.98	*V	11.4	M9
J154131.4–252043	CD–24 12231	1.39e–11	15 41 31.23	–25 20 36.2 ^O	3.0	0.83	*	10.7	K0
J161216.4–282500	SV* ZI 1217	8.58e–12	16 12 16.34	–28 25 00.9 ^O	5.9	0.76	*V	5.7	B9Vvar
J161441.0+335125	HD 146361 J	1.32e–10	16 14 41.01	+33 51 43.5	4.6	1.00	*RS	5.6	G0Ve
J165529.3–082008	V* V1054 Oph	1.85e–10	16 55 28.89	–08 20 10.2	10.7	1.00	*Fl	10.6	M3Ve
J172839.3+590158	V* GR Dra	1.92e–11	17 28 39.36	+59 02 07.0 ^O	3.3	0.87	*Pu	8.8	G0
J173241.3+741337	V* DR Dra	3.13e–11	17 32 40.45	+74 13 32.8	3.1	0.99	*RS	7.7	K0III
J173734.1+414618		2.11e–12	17 37 32.90	+41 46 12.4	10.4	0.99	*	(12.5)	
J204009.4–005216	V* AE Aqr	2.94e–11	20 40 09.02	–00 52 15.5 ^O	4.3	0.93	*DQ	10.4	K5IV–Vvar
			20 40 08.92	–00 52 23.7 ^O	4.3	0.05		(18.7)	
J204151.2–322604	V* AT Mic	1.15e–10	20 41 51.08	–32 26 03.2	3.3	0.99	*V	11.8	M4.5
J204512.3–350953	V* BX Mic	1.93e–11	20 45 11.79	–35 10 01.0	3.8	1.00	*bL	7.4	G0V
J210952.2+160920	V* NR Peg	4.20e–11	21 09 52.10	+16 09 26.7	2.7	0.78	*bL	8.7	G0
J212958.4+120959	NGC 7078	1.53e–10	21 29 59.27	+12 10 14.1	6.3	0.98	*Cl	–	
J221719.9–084801	GJ 852	6.98e–12	22 17 20.32	–08 48 02.2 ^O	7.0	–	*	(15.9)	M4.5Vmevar
J225302.0+165029	V* IM Peg	8.33e–11	22 53 02.39	+16 50 29.4 ^O	3.0	0.87	*RS	7.0	K1.5II–IIIe
J232301.2–063545	HD 220338	1.74e–11	23 23 01.09	–06 35 45.8	13.4	1.00	*	10.2	K0
J233152.6+195735	V* EQ Peg	2.62e–11	23 31 51.81	+19 56 15.2	7.8	–	*	11.5	M3.5
J234351.0–151655	V* R Aqr	1.73e–12	23 43 49.56	–15 17 03.7	18.4	0.99	*Mi	8.8	M7IIIpevar
J234940.8+362525	AG+36 2436	2.38e–11	23 49 41.00	+36 25 30.8	18.1	1.00	*Ro	6.7	G1IIIe

Table 1—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)
<i>ROSAT</i> BSC	SIMBAD	S _x	Position		S _{1.4GHz}	ID	SIMBAD		
1RXS name	Common name	ergs s ^{−1} cm ^{−2}	α	(J2000) δ	mJy	P	Type	m _B	Sp. Class.

^OOptical position given: because identification is more than 3σ from radio position or there are multiple optical candidates

^EExtended radio emission.

Table 3. RBSC-NVSS Results

Selection criteria imposed	No. of identifications found	No. of spurious identifications expected
RBSC sample	18811	–
Count rate $\geq 0.1 \text{ count s}^{-1}$	8547	–
$ b \geq 15^\circ, \delta \geq -40^\circ$	5441	–
RBSC-NVSS matches with Δ position $< 3\sigma$ and $f_r \geq 2.5 \text{ mJy}$	2300	919
Linked cross-identification matches	1556	51*

*The flux density and magnitude distributions of the spurious identifications are markedly different from the RBSC-NVSS sample,

REFERENCES

- Arimoto, N. , Matsushita, K. , Ishimaru, Y. , Ohashi, T. & Renzini, A. 1997, ApJ, 477, 128
- Bahcall, J. N. and Soneira, R. M. 1980, ApJS, 44, 73
- Bade, N., Engels, D., Voges, W., Beckmann, V., Boller, Th., Cordis, L., Dahlem, M., Engshauser, J., Molthagen, K., Nass, P., Studt, J. & Reimers, D. 1998, A&A, 127, 145 and <ftp://ftp.hs.uni-hamburg.de/pub/outgoing/rass-id/rass2idv2.0.cat>
- Bauer, F. B., Thuan, T. X., & Condon, J. J., in preparation
- Brinkmann, W., Siebert, J., Reich, W., Furst, E., Reich, P., Voges, W., Trumper, J., & Wielebinski, R. 1995, A&AS, 109, 147
- Brinkmann, W., Laurent-Muehleisen, S., Voges, W., Siebert, J., Becker, R., Brotherton, M., White, R. and Gregg, M., A&A, 2000, in press
- Burstein, D. , Jones, C., Forman, W., Marston, A. P. & Marzke, R. O. 1997, ApJS, 111, 163
- Ceballos, M. T. & Barcons, X. 1996, MNRAS, 282, 493
- Condon, J. J., Anderson, E., & Broderick, J. J. 1995, AJ, 109, 2318
- Condon, J. J., Cotton, W. D., Greisen, E. W., Yin, Q. F., Perley, R. A., Taylor, G. B., & Broderick, J. J. 1998, AJ, 115, 1693 and <http://www.cv.nrao.edu/~jcondon/nvss.html>
- Condon, J., Kaplan, D. & Yin, Q. 1997, BAAS, 191, 1402
- Condon, J. J., Yin, Q. F., Thuan, T. X. & Boller, Th. 1998, AJ, 116, 2682
- Dickey, J.M., & Lockman, F.J. 1990, ARA&A, 28, 215
- Evans, T. 1988, Ph.D. Thesis, Univ. Cambridge
- Fabbiano, G., Kim, D. -W. & Trinchieri, G. 1992, ApJS, 80, 531
- Hasinger, G., Truemper, J. & Schmidt, M. 1991, A&A, 246, L2
- Lasker, et al. 1990, AJ, 99, 2019
- Laurent-Muehleisen, S. A., Kollgaard, R. I., Ryan, P. J., Feigelson, E. D., Brinkmann, W., & Siebert, J. 1997, A&AS, 122, 235
- Li, J. & Jin, W. 1996, A&AS, 120, 201

- Monet, D., Bird, A., Canzian, B., Harris, H., Reid, N., Rhodes, A., Sell, S., Ables, H., Dahm, C., Guetter, H., Henden, A., Leggett, S., Levison, H., Luginbuhl, C., Martini, J., Monet, A., Pier, J., Rieke, B., Stone, R., Vrba, F., & Walker, R. 1996, USNO-A1.0, (U.S. Naval Observatory, Washington DC) and <http://www.usno.navy.mil/pmm/>
- Moshir, M. et al. 1992, Explanatory Supplement to the IRAS Faint Source Survey, Version 2, JPL D-10015 8/92 (Jet Propulsion Laboratory, Pasadena)
- NED 1999, <http://nedwww.ipac.caltech.edu/NED.html>
- Owen, F. N., White, R. A. & Ge, J. 1993, ApJS, 87, 135
- Peres, C. B., Fabian, A. C., Edge, A. C., Allen, S. W., Johnstone, R. M. & White, D. A. 1998, MNRAS, 298, 416
- Sakelliou, I. & Merrifield, M. R. 1998, MNRAS, 293, 489
- Sutherland, W. & Saunders, W. 1992, MNRAS, 259, 413
- Voges, W. et al. 1999, A&A, 349, 389
- White, R. L., Becker, R. H., Helfand, D. J., & Gregg, M. D. 1997, ApJ, 475, 479 and <http://sundog.stsci.edu/>
- Windhorst, R. A., Kron R. G., & Koo, D. C. 1984, A&AS, 58, 39

Fig. 1.— Diagram of a typical RBSC-NVSS field. Each search area contains some numbers $m \geq 0$ of NVSS radio sources and $l \geq 0$ of optical objects from the USNO catalog. In this case, there are three optical sources (O) and two radio sources (R) positionally offset from the X-ray source (X) by varying angles (ϕ).

Fig. 2.— Logarithm of the ratio of the probability $p[S|x]$ that a RBSC-NVSS source has flux density S to the probability $p[S|\bar{x}]$ that an unrelated NVSS sources has flux density S . This plot indicates that most background radio sources are much fainter than the true radio identifications of X-ray sources and can be exploited to enhance the probabilities of identifications of strong radio candidates.


Fig. 3.— Logarithm of the ratio of the probability $p[\mu|x]$ that a RBSC-NVSS optical identification has magnitude μ to the probability $p[\mu|\bar{x}]$ that an unrelated USNO object (Bahcall & Soneira 1980) has magnitude μ . This plot demonstrates that the magnitude distribution of background optical sources is similar to the optical identifications of RBSC sources, yielding little or no enhancement of the probabilities of optical counterparts except at bright ($B < 13$) magnitudes.

Fig. 4.— Comparison of B_{USNO} and B_{NED} . The straight line represents a one-to-one correspondence and the curve is a least-square fit to the data for $B_{\text{USNO}} < 16$ mag.

Fig. 5.— Sample finding chart of RBSC-NVSS sources. The NVSS radio contours are overlaid on the DSS images, and the crosses denote the *ROSAT* error box. In applicable fields, the 3σ error ellipses of IRAS sources have been plotted.


Fig. 6.— The sky distribution of the RBSC-NVSS sources. The overall distribution is fairly uniform in the area $|b| > 15^\circ$, $\delta > -40^\circ$.


Fig. 7.— The cumulative fraction f of RBSC-NVSS identifications as a function of reliability P (Equation 15).




$$m = 2$$


$$l = 3$$








J000409.2+452618








Table 2. RBSC Extragalactic Identifications

(1) <i>ROSAT</i> BSC 1RXS name	(2) NED Common name	(3) S_x ergs s ⁻¹ cm ⁻²	(4) Position α (J2000) δ		(5) $S_{1.4\text{GHz}}$ mJy	(6) ID P	(7) m_B	(8) Redshift	(9) NED Morphology	(10) Sp. Class.
J000000.0-392902	ESP 21616	1.61e-12	23 59 56.35	-39 28 46.3 ^O	3.3	0.25	19.0			
	ESP 21615		23 59 56.68	-39 28 37.2 ^O	3.3	0.34	18.9			
J000007.0+081653	UGC 12890	5.98e-12	00 00 06.98	+08 16 46.3	83.5	1.00	16.0	0.0387 ¹		Early
J000226.6+032105	NGC 7811	5.84e-12	00 02 25.94	+03 21 02.7	5.8	1.00	15.1	0.0255	cIm?	Sy1
J000312.3-355541 [†]	ESO 349- G 022	5.73e-12	00 03 12.88	-35 56 12.9	517.1	1.00	14.6	0.0498	S0?	
J000350.4+020340		4.32e-12	00 03 49.50	+02 03 52.7 ^O	11.2	0.29	(17.0)			
	NVSS J000349+020358		00 03 49.67	+02 03 59.3 ^O	11.2	0.71	(16.2)	0.0978 ¹		cD
J000409.2+452618		5.30e-12	00 04 09.48	+45 26 25.6 ^O	4.0	0.89	(16.9)	0.1209 ¹		Sy1
			00 04 09.97	+45 26 17.3 ^O	4.0	0.11	(19.1)			
J000416.8-290240		1.30e-12	00 04 17.40	-29 02 36.2	16.0	1.00	(17.5)			
J000459.1+114205	UGC 00032	4.48e-12	00 04 58.46	+11 42 03.3	31.4	1.00	17.0	0.0761	E	
J000559.1+160955	PG 0003+158	8.23e-12	00 05 59.41	+16 09 46.7	805.2 ^E	1.00	16.4	0.4509	Opt.var.	
J000618.9+201215	PG 0003+199	6.29e-11	00 06 19.45	+20 12 10.3	7.6	1.00	13.8	0.0256	S0/a	Sy1
J000620.3+105207	NVSS J000620+105150	4.20e-12	00 06 20.24	+10 51 50.4	12.4	1.00	17.3	0.1676 ¹		Early
J000729.3+024059	LBQS 0004+0224	2.58e-12	00 07 27.05	+02 41 11.4	7.4	1.00	17.3	0.3000		QSO
J000835.4-233917		4.09e-12	00 08 35.36	-23 39 27.8	36.1	1.00	(17.3)			
J001010.2-044225	MRK 0937	4.80e-12	00 10 09.84	-04 42 32.6	6.9	0.99	14.7	0.0295	SBa/b	Sy2
J001031.3+105832 [†]	MRK 1501	1.52e-11	00 10 30.91	+10 58 27.4	98.6	1.00	15.4	0.0893	Opt.var.	Sy1
J001118.5-285126 [†]	ESO 409- G 025	1.35e-11	00 11 21.65	-28 51 15.5 ^O	11.4	0.82	15.2	0.0609	E+4	
J001143.0-152312		2.23e-12	00 11 43.36	-15 23 19.3	32.0	1.00	(19.0)			
J001234.8-162809	PMN J0012-1628	1.62e-12	00 12 33.82	-16 28 06.5	94.7	1.00	17.7	0.1510		QSO
J001328.4+094942		5.56e-12	00 13 28.79	+09 49 31.0	31.0	1.00	(18.7)			
J001356.6-185408	NPM1G -19.0008	1.01e-11	00 13 56.11	-18 54 06.7	29.5	1.00	17.0	0.0944 ¹		Early
J001528.3+353641		3.45e-12	00 15 27.95	+35 36 40.5	11.1	1.00	(18.4)			
J001620.4-143008	PKS 0013-14	2.21e-12	00 16 19.94	-14 30 10.8 ^O	767.0 ^D	1.00	17.0	0.7664 ¹		QSO
J001823.8+300357 [†]	NGC 0071	3.72e-12	00 18 23.58	+30 03 48.0 ^{OR}	7.3 ^E	0.93	14.8	0.0224	SA0 ⁻ pec	
J001827.8+294730	RBS 0042	1.33e-11	00 18 27.75	+29 47 30.4	33.7	1.00	19.1	0.1000		BLLAC
J001906.0-202638 [†]		2.37e-12	00 19 07.98	-20 26 28.7 ^O	31.9	0.86	(16.7)			
			00 19 08.16	-20 26 29.9 ^O	31.9	0.14	(18.3)			
J001937.0+260316	[HB89] 0017+257	3.94e-12	00 19 39.81	+26 02 52.6 ^{OR}	684.3 ^E	1.00	15.4	0.2840		QSO
J001938.0+033646		2.67e-12	00 19 36.77	+03 37 14.0 ^O	9.0 ^A	0.45	(18.8)			
			00 19 36.81	+03 37 40.5 ^O	9.0 ^A	0.24	(19.1)			
			00 19 36.85	+03 37 43.8 ^O	9.0 ^A	0.21	(18.9)			
			00 19 37.69	+03 37 47.4 ^O	9.0 ^A	0.08	(19.0)			
J002041.8-254307 [†]	DUKST 473-042	5.07e-12	00 20 42.96	-25 42 39.5	23.8	0.99	(15.8)	0.0638		
J002108.1-190950	PGC 001348	1.70e-12	00 21 08.22	-19 10 10.3 ^R	1082.3 ^T	1.00	17.0	0.0952	E	
J002136.8+280305 [†]		4.18e-12	00 21 36.26	+28 03 11.0 ^O	3.5	0.69	(16.5)	0.0946 ¹		Early
	IV Zw 015 NOTES01		00 21 36.84	+28 03 02.2 ^O	3.5	0.19	(16.0)			
J002339.6-175352		1.88e-12	00 23 39.29	-17 53 54.8	41.6	1.00	(15.4)	0.0535 ¹		Sy1.5
J002430.5-292856 [†]	PKS 0021-29	4.22e-12	00 24 30.12	-29 28 48.9	2923.8 ^E	1.00	19.0	0.4060		Sy1
J002534.9-330255 [†]	ESO 350- G 015	3.58e-12	00 25 31.44	-33 02 46.4 ^O	1365.2 ^S	-	14.2	0.0498	E+3	
J002723.5-213325		1.71e-12	00 27 23.17	-21 33 05.8 ^O	3.0	0.33	(16.3)			

Table 2—Continued

(1) <i>ROSAT</i> BSC 1RXS name	(2) NED Common name	(3) S_x ergs s ⁻¹ cm ⁻²	(4) Position α (J2000) δ	(5) $S_{1.4\text{GHz}}$ mJy	(6) ID P	(7) m_B	(8) Redshift	(9) NED Morphology	(10) Sp. Class.
J002811.6+310342 [†]	MG3 J002814+3102	5.10e-12	00 27 23.85 -21 32 56.6 ^O	3.0	0.17	(20.0)			
J002836.2-233849	EDCC 447:002606.9-2355	2.62e-12	00 27 23.47 -21 33 02.8 ^O	3.0	0.44	(15.9)			
			00 28 10.73 +31 03 48.1 ^{OR}	207.2 ^S	0.99	18.0	0.4989 ¹		Sy1
			00 28 36.88 -23 38 37.4 ^O	3.9	0.17	17.8			
			00 28 37.33 -23 38 46.6 ^O	3.9	0.07	19.1	0.1108		
			00 28 37.47 -23 38 22.9 ^O	3.9	0.14	(16.7)			
			00 28 37.85 -23 38 31.0 ^O	3.9	0.17	(17.8)			
	APMBGC 473-051-063		00 28 36.89 -23 38 30.9 ^O	3.9	0.28	17.8	0.1120	E	
J002913.9+131605	PG 0026+129	1.26e-11	00 29 13.79 +13 16 03.6	7.4	1.00	15.4	0.1420		Sy1
J003037.6-241053 [†]	NVSS J003035-241115	1.67e-12	00 30 35.75 -24 11 15.3	8.0 ^E	0.93	(16.1)	0.1381		
J003322.8-203908	NPM1G -20.0015	4.63e-12	00 33 22.50 -20 39 08.5	8.6	1.00	17.1			
J003334.6-192130	NVSS J003334-192133	1.32e-11	00 33 34.33 -19 21 33.7	18.8	1.00	16.3	- ¹		BLLAC?
J003413.7-212619 [†]	ESO 540- G 001	4.49e-12	00 34 13.67 -21 26 16.9	43.5	1.00	14.0	0.0268	(R' ₁)SB(rl)bc	Sy1.8
J003514.9+151513	RBS 0082	6.11e-12	00 35 14.62 +15 15 02.8	18.7	1.00	16.3	0.2504 ¹		BLLAC
J003815.8-245858	[HB89] 0035-252	2.04e-12	00 38 14.74 -24 59 02.0	412.9	1.00	17.5	1.1960		QSO
J003908.2-222002	APM 003638.93-223630.6	2.14e-12	00 39 08.31 -22 20 01.6	117.0	1.00	14.5	0.0644	S0	
J004016.5-271913		4.15e-12	00 40 16.44 -27 19 12.2	160.5	1.00	(17.8)			
J004030.3+293328	UGC 00428	8.90e-12	00 40 28.21 +29 33 21.5	15.2	1.00	16.0	0.0713 ¹		Early
J004051.0+100332	3C 018	3.56e-12	00 40 50.35 +10 03 23.3	4603.3 ^E	1.00	18.5	0.1880	Radio galaxy	
J004149.7-091817	MCG -02-02-086	9.68e-11	00 41 50.38 -09 18 13.7	58.1	1.00	13.5	0.0552	cD;SB(r)0 ⁰ pec	
J004208.1+364115	NVSS J004208+364112	3.68e-12	00 42 08.22 +36 41 12.8	11.9	1.00	17.5	- ¹		BLLAC
J004226.6-133536		1.92e-12	00 42 27.04 -13 35 33.9	3.8	1.00	(17.1)	0.2903 ¹		Sy1/QSO
J004240.8+301742	RX J0042.6+3017	4.15e-12	00 42 39.95 +30 17 43.4	5.0	0.96	17.4	0.1407 ¹		Sy1.5
J004241.8+411535	MESSIER 031	3.26e-11	00 42 44.00 +41 16 06.0 ^O	120.4 ^E	1.00	4.4	0.0004	SA(s)b	LINER
J004324.9-203728 [†]		2.60e-12	00 43 24.71 -20 37 41.6	106.2	1.00	(18.2)			
J004333.7-044257		4.49e-12	00 43 34.13 -04 43 01.7	29.7	1.00	(17.3)	- ¹		
J004343.6+372521	IRAS F00409+3708	3.74e-12	00 43 42.55 +37 25 19.6	21.3	1.00	17.4	0.0801 ¹		Sy1
J004351.2+242422	87GB 004111.7+240742	1.28e-11	00 43 52.21 +24 24 21.5	49.9	1.00	17.2	0.0830	cD	
J004519.6+212735	87GB 004239.8+211104	3.62e-12	00 45 19.27 +21 27 42.8	50.1	1.00	17.8			BLLAC
J004733.3-251722	NGC 0253	3.59e-12	00 47 33.30 -25 17 20.0 ^O	2994.7 ^E	1.00	8.0	0.0008	SAB(s)c	HII
J004924.0-293128 [†]		3.83e-12	00 49 23.70 -29 31 43.6 ^O	3.4	0.55	(18.3)			
			00 49 24.04 -29 31 13.2 ^O	3.4	0.08	(15.7)			
	DUKST 411-062		00 49 22.94 -29 31 12.7 ^O	3.4	0.26	17.2	0.1049		
J004937.9-184751		1.86e-12	00 49 37.05 -18 47 52.6 ^O	3.0	0.82	(17.2)	0.2898 ¹		Sy1/QSO
			00 49 37.71 -18 47 46.0 ^O	3.0	0.17	(20.1)			
J005041.0-092855	[HB89] 0048-097	4.86e-12	00 50 41.27 -09 29 06.2	814.6	1.00	17.4		Opt.var.	BLLAC
J005118.2-283059	EDCC 485:004849.7-2847	3.01e-12	00 51 15.64 -28 31 32.9 ^{OR}	32.5 ^E	0.77	17.5	0.1111		
J005250.9+433617		4.63e-12	00 52 50.07 +43 36 34.3 ^O	4.0	0.05	(18.6)			
	NPM1G +43.0016		00 52 50.45 +43 36 14.3 ^O	4.0	0.91	17.7	0.0590 ¹		Sy1.9
J005335.3+124137	UGC 00545	2.38e-11	00 53 34.77 +12 41 33.8	8.8	0.98	14.4	0.0611	Compact	Sy1
J005447.2-245532		2.51e-12	00 54 46.76 -24 55 29.0	24.3	1.00	(17.5)			
J005620.1-093626	NPM1G -09.0033	8.85e-12	00 56 20.37 -09 36 23.5	200.7 ^E	1.00	16.7	0.1010		BLLAC

Table 2—Continued

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position	S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α (J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J005817.1+172300	RBS 0140	8.58e-12	00 58 16.77 +17 23 14.8	9.2	1.00	19.8			BLLAC
J005822.8-024126	NPM1G -02.0022	2.96e-12	00 58 22.36 -02 41 36.5 ^O	4.1	0.85	16.6	0.0728 ¹		Sy1.5
			00 58 23.21 -02 41 27.1 ^O	4.1	0.08	(18.0)			
J005932.3-351049		1.93e-12	00 59 31.43 -35 10 47.3	80.2	1.00	(18.5)			
J010013.4-151755	MCG -03-03-017	2.16e-12	01 00 15.91 -15 17 56.4	33.9	1.00	15.3	0.0542		
J010251.3-200154	PMN J0102-2001	2.31e-12	01 02 50.72 -20 01 55.9	64.7	1.00	(17.8)			
J010538.7-141610		2.19e-11	01 05 38.36 -14 16 10.9 ^R	5.8	1.00	(15.7)	0.0666 ¹		Sy1
J010721.2+322254 [†]	NGC 0383	3.62e-12	01 07 24.96 +32 24 45.1 ^{OR}	3785.4 ^T	0.08	13.4	0.0170	SA0 ⁻ :	
J010750.4-364342 [†]	GSN 093	1.70e-12	01 07 50.37 -36 43 22.1 ^O	9.5	0.91	16.4	0.1217		
			01 07 50.54 -36 43 33.7 ^O	9.5	0.07	(18.2)			
J010850.5-152438 [†]	MCG -03-04-008	5.28e-12	01 08 51.11 -15 24 24.5	6.5	1.00	15.5	0.0515	E	
J010908.3+181601	RBS 0157	9.57e-12	01 09 08.20 +18 16 06.6	90.8	1.00	16.2	- ¹		BLLAC?
J010910.5-323249		2.95e-12	01 09 10.27 -32 32 49.2 ^O	2.8	0.24	(20.1)			
	IRAS F01068-3248		01 09 11.20 -32 32 43.8 ^O	2.8	0.76	(16.5)			
J010950.3-125323		3.13e-12	01 09 51.53 -12 53 24.2	3.2	1.00	(21.5)			
J011034.9-164821	PMN J0110-1648	2.90e-12	01 10 35.13 -16 48 31.3	106.4 ^E	1.00	15.6	0.7787 ¹		Sy1
J011050.0-125455	RBS 0161	1.43e-11	01 10 49.96 -12 55 04.9	17.4	1.00	17.9	0.2315? ¹		QSO/Early?
J011058.7+330900	NGC 0410	3.54e-12	01 10 58.94 +33 09 05.9	6.3	1.00	12.5	0.0177	E+	LINER
J011114.6-161548		1.98e-12	01 11 13.78 -16 16 02.0 ^O	2.5	0.11	(18.4)			
	IRAS F01087-1631		01 11 14.26 -16 15 54.2 ^O	2.5	0.89	16.0	0.0518 ¹		Sy1.5
J011130.5+053612	NVSS J011130+053626	4.04e-12	01 11 30.30 +05 36 26.5	16.5	1.00	19.9	0.3460		BLLAC
J011205.2+224452	[HB89] 0109+224	4.27e-12	01 12 05.83 +22 44 38.0	385.8	1.00	15.7		Opt.var.	BLLAC
J011301.1+153041 [†]	VV 382 NED01	2.98e-12	01 12 59.68 +15 29 28.7 ^{OR}	468.1 ^D	-	14.1	0.0440	cD	
J011350.0-145041	MRK 1152	1.32e-11	01 13 49.78 -14 50 43.5	4.3	1.00	15.0	0.0527	S: pec	Sy1.5
J011354.9-314538 [†]	NGC 0439	2.00e-12	01 13 47.23 -31 44 50.3 ^{OR}	22.5	-	12.6	0.0194	SAB(rs)0 ⁻ ?	
J011501.3-340008		1.90e-12	01 15 01.62 -34 00 26.9 ^O	6.3	0.41	(19.2)			
			01 15 02.03 -34 00 27.8 ^O	6.3	0.58	(18.6)			
J011513.2+405743 [†]		5.09e-12	01 15 15.38 +40 57 30.7 ^O	6.4	0.14	(17.0)			
			01 15 14.89 +40 57 17.4 ^O	6.4	0.84	(15.5)	0.0978 ¹		Early
J011546.1+252006	87GB 011303.4+250427	5.45e-12	01 15 46.11 +25 19 54.5	38.2	0.99	19.0			BLLAC
J011555.6-274428		1.98e-12	01 15 55.54 -27 44 31.3	17.2	1.00	(19.0)			
J011637.7-281146		1.63e-12	01 16 37.06 -28 11 46.4	54.0	1.00	(18.5)			
J011746.6-244329		2.72e-12	01 17 46.90 -24 43 34.8	10.2	1.00	(17.9)			
J011811.6-265819		3.32e-12	01 18 11.15 -26 58 13.2	6.7	1.00	(16.4)			
J011834.7-184916	PKS 0116-19	2.55e-12	01 18 34.26 -18 49 14.9	1171.1	0.98	18.0	0.2800	N galaxy	
J011835.7-184334		1.33e-12	01 18 36.00 -18 43 32.2	5.0	1.00	(16.9)	0.4461 ¹		Sy1/QSO
J011905.4-145906		2.36e-12	01 19 04.70 -14 58 58.3	5.4	1.00	(17.7)			
J012058.3-135100 [†]	GIN 069	6.12e-12	01 20 58.77 -13 50 58.6	8.1	0.99	14.0	0.0508 ¹		Early
J012159.9-010223	[HB89] 0119-013	3.90e-12	01 22 00.06 -01 02 38.6	4.6	1.00	15.1	0.0543		Sy1
J012237.4-264646		4.23e-12	01 22 37.48 -26 46 45.0	3.8	1.00	(18.3)			
J012308.9+342048	87GB 012020.0+340518	5.86e-11	01 23 08.55 +34 20 47.5	45.6	1.00	15.2	0.2720		BLLAC
J012337.2+331508 [†]	NGC 0507	9.97e-12	01 23 39.77 +33 15 23.2 ^O	102.3 ^D	0.98	12.2	0.0164	SA(r)0 ⁰	

Table 2—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position		S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	δ (J2000)	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J012338.2–231100		8.47e–12	01 23 38.35	–23 10 59.4	27.5	1.00	(18.0)			
J012340.7+421019		5.72e–12	01 23 40.41	+42 10 18.3 ^O	21.2	0.06	(18.6)			
			01 23 40.48	+42 10 06.7 ^O	21.2	0.94	(19.3)	0.1856? ¹		BLLAC
J012354.4–350402	NGC 0526A	1.98e–12	01 23 54.57	–35 03 57.5	13.9	1.00	14.5	0.0191	S0pec?	Sy1.5
J012442.9–314327		2.08e–12	01 24 43.64	–31 43 41.4	13.4	1.00	(18.9)			
J012507.3+084124 [†]	IC 1695	2.16e–11	01 25 07.87	+08 41 59.2 ^O	59.2 ^S	0.88	14.9	0.0484	cD;S?	
J012529.4+014603	NGC 0533	5.33e–12	01 25 31.45	+01 45 34.2	29.1	1.00	12.4	0.0185	cD;E3:	
J012657.1+330730		4.47e–12	01 26 56.95	+33 07 27.4	6.9	1.00	(17.3)			
J012732.9+191043	UGC 01032	1.74e–11	01 27 32.20	+19 10 45.1	5.3	1.00	14.2	0.0174	SB0a	Sy1.5
J012910.8–214158 [†]	IRAS 01267–2157	1.04e–11	01 29 10.76	–21 41 57.3	2.4	1.00	18.5	0.0930	SB(s)a	Sy1
J013152.8–133651 [†]	ABELL 0209:[KSW78] A	4.63e–12	01 31 52.74	–13 36 59.6	19.1	0.99	17.3	0.2130		
J013240.5–080408	PKS 0130–083	4.17e–12	01 32 41.11	–08 04 05.7	307.5	1.00	(16.5)	0.1485 ¹		Early
J013312.0–351925		1.73e–12	01 33 12.26	–35 19 16.6	10.9 ^E	1.00	(17.3)			
J013350.9+303932	MESSIER 033	1.74e–11	01 33 51.26	+30 39 33.9	3.7 ^E	1.00	6.3	0.0005	SA(s)cd	
J013523.1–272807		1.55e–12	01 35 23.75	–27 28 13.0	8.3	1.00	(17.9)			
J013526.9–042636	IRAS F01329–0441	8.37e–12	01 35 27.18	–04 26 38.7	8.0	1.00	(16.5)	0.1547 ¹		Sy1
J013624.3+205712	[HB89] 0133+207	5.11e–12	01 36 24.45	+20 57 27.5 ^O	3901.9 ^D	1.00	18.1	0.4250	Opt.var.	QSO
J013632.9+390556 [†]	B3 0133+388	2.43e–11	01 36 32.43	+39 05 59.4	60.5	1.00	15.0	– ¹		BLLAC?
J013715.1–091145 [†]		8.20e–12	01 37 15.96	–09 11 55.2 ^O	179.6	0.10	(17.3)			
	MCG –02–05–020		01 37 15.38	–09 11 51.3 ^O	179.6	0.88	14.3	0.0411 ¹		Early
J013738.2–243053	[HB89] 0135–247	2.09e–12	01 37 38.09	–24 30 54.4	1181.9	1.00	17.3	0.8310		QSO
J013741.7+330931	[HB89] 0134+329	1.80e–11	01 37 41.27	+33 09 35.6	16018.2	1.00	16.1	0.3670	Opt.var.	Sy1?
J013801.3–215516		1.74e–12	01 38 03.22	–21 55 29.4 ^O	17.6	0.29	(19.6)			
			01 38 03.78	–21 55 31.6 ^O	17.6	0.62	(18.2)			
J013958.5+013151	[HB89] 0137+012	3.23e–12	01 39 57.48	+01 31 51.3	1311.6	0.99	17.1	0.2600	Opt.var.	QSO
J014005.1–153309	PMN J0140–1533	1.42e–12	01 40 04.37	–15 32 55.3	233.1	1.00	(16.3)	0.8188 ¹		Sy1/QSO
J014006.0+112918	RX J0140.1+1129	3.17e–12	01 40 04.51	+11 29 31.9	5.2	0.97	16.2	0.0650 ¹		Sy1
J014016.7–005005	[HB89] 0137–010	5.17e–12	01 40 17.82	–00 49 55.7	3.3 ^E	1.00	16.5	0.3340	Opt.var.	QSO
J014040.9–075850		9.06e–12	01 40 40.82	–07 58 48.6	27.2	1.00	(18.4)			
J014442.0–221339	ABELL 2938:001	1.80e–12	01 44 40.40	–22 13 48.8	10.3	1.00	17.4	0.2780	cD	
J014526.6–034945	MCG –01–05–031	3.50e–12	01 45 25.29	–03 49 43.5 ^R	9.2 ^E	1.00	14.0	0.0182	SB(rs)bc pec:	Sy2
J014645.6–004039	NPM1G –00.0070	2.88e–12	01 46 44.26	–00 40 49.2	8.7 ^E	0.99	15.9	0.0800		Sy1.5
J014828.3–040656	[ZGH93] 0145–0422	1.98e–12	01 48 27.61	–04 07 47.8	12.0	0.99	(16.3)	0.0862		
J014913.6+130352	NGC 0675	3.23e–12	01 49 08.65	+13 03 34.9 ^O	3.1 ^E	0.05	15.5	0.0177	S	
	NGC 0677		01 49 14.08	+13 03 19.1 ^O	20.6 ^E	0.89	13.2	0.0170	E	
J015227.1–231956	IRAS F01501–2334	1.16e–11	01 52 27.00	–23 19 52.5	5.0	1.00	15.6	0.1130		Sy
J015235.5–285247		2.60e–12	01 52 34.17	–28 53 30.4 ^O	8.7 ^A	0.21	(20.5)			
			01 52 34.39	–28 53 56.0 ^O	8.7 ^A	0.11	(19.6)			
			01 52 34.44	–28 53 37.9 ^O	8.7 ^A	0.25	(20.2)			
J015240.2+014718	NPM1G +01.0067	5.95e–12	01 52 39.58	+01 47 17.0	61.9	1.00	16.6	0.0790 ¹		Early
J015244.7+360855 [†]	NGC 0708	1.73e–11	01 52 46.45	+36 09 08.3	66.8 ^E	1.00	13.7	0.0161	cD;E	Sy2
J015332.8–011853	PMN J0153–0117	2.77e–12	01 53 34.34	–01 18 09.6	111.1 ^S	1.00	(19.0)	0.2458 ¹		Early

Table 2—Continued

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position	S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α (J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J015705.3+412029 [†]	HFLLZOA F215	8.19e-12	01 57 05.07 +41 20 31.3	46.4	0.99	13.8	0.0785 ¹		Early
J015935.1+104707	87GB 015652.7+103213	4.29e-12	01 59 34.41 +10 47 05.9	47.5	1.00	16.8			BLLAC
J015949.9-084949	PMN J0159-0849	2.62e-12	01 59 49.38 -08 49 58.5	38.7	1.00	(19.2)			
J015950.6+002337	PG 0157+001	3.88e-12	01 59 50.25 +00 23 38.9	26.7	1.00	15.7	0.1630		Sy1
J020014.5+312543	NGC 0777	4.41e-12	02 00 14.88 +31 25 45.5	7.4	1.00	12.5	0.0166	E1	
J020026.7+024012	MRK 0584	7.93e-12	02 00 26.37 +02 40 12.9	5.7	1.00	15.3	0.0780	Compact	Sy1
J020030.2+271238	RX J0200.5+2712	4.09e-12	02 00 29.49 +27 12 39.1	15.7	1.00	19.7			BLLAC
J020106.3+003400	MS 0158.5+0019	6.60e-12	02 01 06.16 +00 34 02.2	13.2	1.00	18.0	0.2990		BLLAC
J020144.2-021158 [†]	FIRST J020143.1-021148	5.18e-12	02 01 43.12 -02 11 48.0 ^{RF}	126.5 ^S	1.00	(16.7)	0.1950		cD
J020157.1-113238	[HB89] 0159-117	2.77e-12	02 01 57.19 -11 32 33.3	2753.3	1.00	16.4	0.6690	Opt.var.	Sy1
J020222.9+190421	UGC 01518	7.62e-12	02 02 18.78 +19 04 01.5	24.6	0.99	16.0	0.0650 ¹		Early
J020251.9-022323		1.96e-12	02 02 52.31 -02 23 21.2	68.0	1.00	(19.1)	0.2600 ¹		BLLAC?
J020301.2-240213		2.56e-12	02 03 00.89 -24 02 01.7	2.4	1.00	(16.3)			
J020413.6-333345		2.16e-12	02 04 13.22 -33 33 33.7	6.3 ^E	1.00	(18.3)			
J020615.9-001728	UGC 01597	6.72e-12	02 06 16.00 -00 17 29.3 ^F	4.3	1.00	14.3	0.0424	S0; merger?	Sy1.5
J020702.5+293035	RBS 0281	6.53e-12	02 07 02.22 +29 30 46.1 ^{OR}	2496.8 ^D	1.00	16.5	0.1094 ¹		Sy1.5
J020837.5+352314	MS 0205.7+3509	7.45e-12	02 08 38.03 +35 23 13.0	4.8	1.00	19.2	0.3180		BLLAC
J020917.6+444951		1.71e-11	02 09 16.93 +44 49 44.7	30.6 ^E	1.00	(18.7)			
J021206.0-255808		1.75e-12	02 12 05.95 -25 57 54.4	8.8	0.96	(18.4)			
J021231.5-350336		2.84e-12	02 12 30.50 -35 03 32.7	19.3	1.00	(18.4)			
J021359.6+004245		2.89e-12	02 13 59.31 +00 42 32.1	3.3	1.00	(18.0)	0.1820 ²		Sy1
J021412.0-353747		2.49e-12	02 14 11.72 -35 37 41.4	19.1	0.99	(19.2)			
J021433.7-004603	NGC 0863	5.30e-11	02 14 33.63 -00 46 00.5	16.8	1.00	13.9	0.0264	SA(s)a:	Sy1.2
J021632.3+231448	RBS 0298	1.44e-11	02 16 32.10 +23 14 50.2 ^{OR}	35.7 ^E	0.63	17.9			BLLAC
J021707.8+110403	[HB89] 0214+108	8.95e-12	02 17 07.67 +11 04 10.2 ^O	1376.2 ^D	0.99	17.0	0.4080	Opt.var.	QSO
J021754.1-195813		2.42e-12	02 17 53.67 -19 58 15.7 ^R	111.7 ^T	1.00	(16.5)			
J021952.9-300141		2.71e-12	02 19 52.92 -30 01 34.5	66.1	1.00	(18.6)			
J022441.4-190826	ESO 545- G 013	4.38e-12	02 24 40.45 -19 08 31.7	31.5	1.00	13.9	0.0337	SAB(s)c	Sy1.8
J022502.9-231254	PKS 0222-23	6.86e-12	02 25 02.61 -23 12 47.6	2096.4	1.00	18.5	0.2300		QSO
J022505.3+184654	MG1 J022505+1846	9.19e-12	02 25 04.64 +18 46 48.4	461.3	1.00	18.7			
J022716.6+020154	RGB J0227+020	3.34e-11	02 27 16.58 +02 01 58.9	36.8	1.00	18.8			BLLAC
J022738.2+441004		4.02e-12	02 27 39.63 +44 09 57.2	6.4	1.00	(17.1)	0.1754 ¹		Sy1
J022814.6+311838	NGC 0931	1.17e-11	02 28 14.40 +31 18 43.6	15.4	1.00	14.5	0.0167	Sbc	Sy1.5
J022904.0+380548	VV 107a	3.36e-12	02 29 01.32 +38 05 53.7	9.9	1.00	(15.3)	0.0375 ¹		Early
J023005.5-085951	MRK 1044	4.67e-11	02 30 05.37 -08 59 44.1	2.9	1.00	15.2	0.0165	SB0	Sy1
J023231.4+340435	IRAS 02295+3351	3.48e-12	02 32 33.14 +34 04 28.9	3.1	0.98	(16.3)	0.0796 ¹		Sy2
J023245.8+063706	NPM1G +06.0105	5.17e-12	02 32 46.23 +06 37 41.9	7.9	1.00	17.7	0.0708 ¹		Early
J023248.3+201722	87GB 022959.3+200408	1.67e-11	02 32 48.46 +20 17 16.2	83.4 ^E	1.00	18.0	0.1400		BLLAC
J023321.6-045455	PKS 0230-051	2.00e-12	02 33 22.51 -04 55 05.4	211.9 ^E	1.00	(17.0)	0.7786 ¹		Sy1
J023349.3+022933	[HB89] 0231+022	4.05e-12	02 33 48.99 +02 29 24.6	52.1	1.00	18.0	0.3210		QSO
J023438.0-084714	NGC 0985	2.60e-11	02 34 37.52 -08 47 15.8	17.3	1.00	13.8	0.0431	(R)SBb/c?	Sy1
J023507.3-040214	[HB89] 0232-042	2.36e-12	02 35 07.33 -04 02 05.1 ^R	1495.9	1.00	16.5	1.4500	Opt.var.	QSO

Table 2—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position		S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	(J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J023513.9–293616	ESO 416– G 002	5.18e–12	02 35 13.21	–29 36 17.2 ^R	240.2 ^E	0.99	15.1	0.0591	Sa:	Sy1.9
J023536.7–293845	PHL 1389	2.70e–12	02 35 36.73	–29 38 45.4	5.3	1.00	16.1			BLLAC
J023727.6–263027 [†]		2.17e–12	02 37 27.43	–26 30 25.9 ^O	5.4	0.32	(17.3)			
			02 37 27.73	–26 30 28.2 ^O	5.4	0.42	(16.5)			
			02 37 27.78	–26 30 20.8 ^O	5.4	0.18	(29.8)			
			02 37 28.21	–26 30 16.0 ^O	5.4	0.06	(19.6)			
J023733.9–360330		4.86e–12	02 37 33.98	–36 03 27.2	27.1	1.00	(16.5)			
J023827.5+015426	NGC 1019	6.82e–12	02 38 27.77	+01 54 34.7	7.6 ^E	1.00	14.3	0.0242	SB(rs)bc	Sy1
J023832.1+023336	PC 0235+0220	3.86e–12	02 38 32.71	+02 33 49.4 ^{O,R}	427.0 ^D	1.00	17.7	0.2090		Sy1
J023832.6–311658		9.75e–12	02 38 32.43	–31 16 58.8	71.4	1.00	(16.8)			
J023849.8–125854	TXS 0236–131	2.09e–12	02 38 49.34	–12 58 49.9	76.6	1.00	(17.6)	0.1951 ¹		Early
J024014.8–033129		2.34e–12	02 40 14.69	–03 31 36.3 ^O	118.8 ^D	0.98	(17.0)			
J024122.1–283919 [†]	TXS 0239–288	2.49e–12	02 41 21.40	–28 39 06.1	97.8	1.00	(17.5)		sym. dbl.	
J024215.2+053037	RBS 0345	1.37e–11	02 42 14.83	+05 30 42.1	3.1	1.00	(15.9)	0.0685 ¹		Sy1
J024235.9–213215	PKS 0240–217	4.54e–12	02 42 35.92	–21 32 25.8	1255.3	1.00	17.0	0.3140	N galaxy	
J024240.9–000046	MESSIER 077	4.21e–11	02 42 40.78	–00 00 47.7	4849.0 ^E	1.00	9.6	0.0038	(R)SA(rs)b	Sy2
J024241.2+005715	E 0240+007	4.25e–12	02 42 40.42	+00 57 29.4	8.1	1.00	16.6	0.5690		AGN
J024620.0–301639 [†]	NGC 1097	2.86e–12	02 46 18.93	–30 16 29.3	250.2 ^E	1.00	10.2	0.0043	(R' ₁ :)SB(r'l)b	Sy1
J024642.8–334340		2.16e–12	02 46 42.57	–33 43 36.0	2.9	0.94	(19.7)			
J024750.8–225003		1.78e–12	02 47 51.56	–22 50 05.7	3.1	1.00	(20.0)			
J024802.6–033206	[CGH98] J0248–0331	5.92e–12	02 48 03.40	–03 31 43.9	41.4	1.00	(16.5)	0.1883 ¹		Early
J024920.8+191813		9.45e–12	02 49 20.22	+19 18 21.3 ^O	4.1	0.11	(18.7)			
	IC 1854		02 49 20.71	+19 18 14.7 ^O	4.1	0.89	14.9	0.0310	S0/a	Sy1.5
J024937.1–311114 [†]	IC 1860	5.41e–12	02 49 33.85	–31 11 21.9	18.7	1.00	13.4	0.0229	E4	
J025018.2–212951		4.40e–12	02 50 18.87	–21 29 37.7	3.6	1.00	(19.2)			
J025038.8+171215	87GB[BWE91] 0247+1700	4.52e–12	02 50 37.79	+17 12 09.3	63.5	1.00	16.1	1.1003 ¹		QSO
J025055.4–361640		2.45e–12	02 50 55.31	–36 16 34.5	9.9	1.00	(16.7)			
J025250.3–011605	NGC 1132	4.58e–12	02 52 51.84	–01 16 29.3 ^F	7.9 ^E	1.00	13.2	0.0232	E	
J025740.7–163050	IRAS 02553–1642	3.59e–12	02 57 40.94	–16 30 47.1	10.6	1.00	16.0	0.0680		Sy1
J025815.7–210527		2.05e–12	02 58 18.98	–21 05 55.8	143.7 ^E	0.83	(16.5)			
J025930.7–242249	IRAS F02572–2434	5.84e–12	02 59 30.63	–24 22 54.9 ^O	3.9	0.92	15.7		S	
J030019.6+385508		4.99e–12	03 00 20.04	+38 54 57.7	8.7	1.00	(16.1)	0.1897 ¹		Sy1
J030136.9+015517	RBS 0381	7.52e–12	03 01 38.23	+01 55 15.2 ^O	393.6	0.67	18.5	0.1700	cD	
			03 01 38.43	+01 55 18.1 ^O	393.6	0.33	(19.3)			
J030150.6+355003 [†]	V Zw 311 NOTES05	6.83e–12	03 01 52.36	+35 50 28.4 ^O	763.0 ^T	0.71	19.0			
	V Zw 311 NOTES02		03 01 51.58	+35 50 27.1 ^O	763.0 ^T	0.09	17.7	0.0470 ¹		Early
	V Zw 311 NOTES04		03 01 51.88	+35 50 18.9 ^O	763.0 ^T	0.20	18.1			
J030326.3–240717	PKS 0301–243	9.42e–12	03 03 26.38	–24 07 10.2	701.0	1.00	16.5			BLLAC
J030330.0+055425	RBS 0384	1.13e–11	03 03 30.03	+05 54 29.8	29.6	1.00	17.9	0.1960		BLLAC
J030416.4–283215		5.06e–12	03 04 16.27	–28 32 20.7	8.2	1.00	(18.6)			
J030455.7–355651	PKS 0303–361	2.48e–12	03 04 56.11	–35 56 47.4	1149.5 ^E	1.00	(17.6)			
J030606.3–390212	NGC 1217	3.18e–12	03 06 05.98	–39 02 10.6	35.8	1.00	13.6	0.0210	SA:(s:)a pec	LINER

Table 2—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position		S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	(J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J030825.9+040637	NGC 1218	6.24e-12	03 08 26.26	+04 06 42.8	5748.1 ^E	0.98	13.8	0.0287	S0/a	Sy1
J031118.9-204621	IRAS F03090-2057	1.67e-11	03 11 19.35	-20 46 15.6	4.9	1.00	(16.0)			
J031227.4+391629	PGC 011941	9.21e-12	03 12 26.34	+39 16 30.5	1936.1 ^E	1.00	18.2	0.1610	Radio galaxy/n	
J031250.7+361527	V Zw 326	6.05e-12	03 12 50.21	+36 15 18.1	117.6	1.00	17.6	0.0712 ¹		Early
J031257.2+082236	PMN J0312+0822	8.24e-12	03 12 57.42	+08 22 08.0	42.7	0.99	(21.5)			
J031422.7+062001	RBS 0400	2.46e-11	03 14 23.90	+06 19 55.3	29.3	1.00	17.9			QSO
J031611.4+090445 [†]	RHS 21	6.87e-12	03 16 12.71	+09 04 44.0	55.4	1.00	17.4	— ¹		QSO
J031615.0-260748	RX J0316.2-2607	4.46e-12	03 16 15.00	-26 07 55.7	21.6	1.00	17.5			BLLAC
J031634.6-221612		2.34e-12	03 16 33.90	-22 16 11.5	4.2	1.00	(17.9)			
J031828.4-212216		4.06e-12	03 18 27.35	-21 22 04.8	3.8	0.98	(16.3)			
J031951.9+184539	[HB89] 0317+185	2.64e-11	03 19 51.70	+18 45 36.5	22.8	1.00	18.1	0.1900	Opt.var.	BLLAC
J032241.8-371239	NGC 1316	2.74e-12	03 22 41.63	-37 12 25.6	255.7 ^E	1.00	9.4	0.0059	(R')SAB(s)0 ⁰	LINER
J032350.5+071746		7.14e-12	03 23 50.79	+07 17 35.2 ^O	4.3	0.73	(21.0)			
			03 23 51.30	+07 17 39.3 ^O	4.3	0.27	(21.4)			
J032441.3+341056	GSC 2349 01904	1.09e-11	03 24 41.33	+34 10 45.0	614.3	1.00	15.7	0.0610		Sy1
J032525.2-060830	MRK 0609	1.14e-11	03 25 25.42	-06 08 37.0	30.4	1.00	15.1	0.0341	Im pec?	Sy1.8
J032540.8-164607		6.17e-11	03 25 41.15	-16 46 15.7	27.5	1.00	(16.7)			
J032613.6+022523	[HB89] 0323+022	3.54e-11	03 26 13.98	+02 25 15.4	68.2	1.00	16.5	0.1470	Opt.var.	BLLAC
J033051.9+053841	RX J0330.8+0538	1.04e-11	03 30 52.65	+05 38 34.8	4.1	1.00	16.7	0.0460		Sy1
J033107.6-210015	MRC 0328-211	5.11e-12	03 31 06.00	-21 00 33.0	168.8	1.00	(16.0)		sym. dbl.	
J033415.2-372538	PMN J0334-3725	1.29e-12	03 34 15.42	-37 25 43.7	222.0	1.00	(16.5)			
J033424.5-151325	IRAS 03321-1523	9.20e-12	03 34 24.74	-15 13 36.0	4.7	1.00	15.4	0.0348 ¹		Sy1.5-8
J033522.6+190739		1.28e-11	03 35 23.16	+19 07 23.0 ^R	4.6	0.99	(16.5)	0.1901 ¹		Sy1
J033623.3-034727		3.92e-12	03 36 24.00	-03 47 39.8	54.8	1.00	(17.4)			
J033630.3+321830	[HB89] 0333+321	1.02e-11	03 36 30.12	+32 18 29.2	2677.1	0.96	17.5	1.2580	Opt.var.	LPQ
J033701.9-095001		5.20e-12	03 37 01.34	-09 49 59.9 ^O	2.6	0.09	(18.0)			
			03 37 03.01	-09 50 01.1 ^O	2.6	0.91	(16.7)	0.2808 ¹		Sy1
J033810.1-244337	E 0336-248	1.87e-12	03 38 12.53	-24 43 50.4 ^O	13.9 ^E	0.80	(19.1)	0.2509		BLLAC
J033828.8-352701 [†]	NGC 1399	3.77e-12	03 38 29.02	-35 27 00.7	628.6 ^T	1.00	10.0	0.0048	E1 pec	
J033841.1+095824	PGC 013424	1.52e-10	03 38 40.62	+09 58 12.2	38.0 ^E	1.00	16.0	0.0346	cD	Sy2
J033851.5-353543	NGC 1404	3.96e-12	03 38 52.50	-35 35 40.0	4.3	1.00	10.9	0.0065	E1	
J033913.4-173553	APMBGC 548-090-125	4.34e-12	03 39 13.68	-17 36 00.9	170.6	1.00	16.2	0.0655 ¹		Early
J034118.9+152423	III Zw 054 NED01	2.36e-11	03 41 17.59	+15 23 44.3	20.8	1.00	17.0	0.0290		
J034203.8-211428	ESO 548- G 081	5.17e-12	03 42 03.19	-21 14 49.9	3.6	0.98	12.9	0.0145	SB(rs)a pec?	Sy1
J034205.4-370311	[HB89] 0340-372	1.69e-12	03 42 05.40	-37 03 20.9	2060.4	1.00	18.6	0.2840		QSO
J034255.1-370742	FCCB 1628	6.59e-12	03 42 54.08	-37 07 37.7	29.6	0.97	18.3		E	
J034308.0+185836	RX J0343.1+1858	8.62e-12	03 43 08.44	+18 58 20.2	3.1	1.00	16.7	0.1090 ¹		Sy1
J034838.5-274906	[HB89] 0346-279	1.36e-12	03 48 38.18	-27 49 14.3	840.0	1.00	19.4	0.9880		QSO
J034922.8-115923		2.99e-11	03 49 22.54	-11 59 14.4	24.7 ^D	1.00	(18.8)			
J035018.4-221708	HE 0348-2226	6.35e-12	03 50 19.17	-22 17 20.6	19.0	1.00	17.2	0.1110		Sy1
J035128.5-142903	[HB89] 0349-146	2.38e-12	03 51 28.54	-14 29 08.8 ^R	3045.1 ^D	1.00	16.2	0.6163		QSO
J035245.6-234254		4.10e-12	03 52 46.74	-23 42 57.1 ^O	4.1	0.92	(16.9)			

Table 2—Continued

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position	S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α (J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J035258.9+194117	RX J0352.9+1941:003	1.09e-11	03 52 58.97 +19 40 59.6	19.6	1.00	18.3	0.1080	cD	
J035308.3+825633		4.30e-12	03 53 09.49 +82 56 31.3	22.7	1.00	(16.0)	0.0694 ¹		Early
J035345.9+195825	IC 0355	5.91e-12	03 53 46.90 +19 58 21.8	7.8 ^E	1.00	15.4	0.0292 ¹		Sy1
J035410.1+024936	PGC 014064	1.49e-11	03 54 09.48 +02 49 30.8 ^O	7.4 ^D	0.62	16.3	0.0360		Sy1
J035432.5-134005		4.65e-12	03 54 32.69 -13 40 12.6	15.6	1.00	(16.6)	0.0766 ¹		Sy1
J035855.6-305437		3.00e-12	03 58 56.20 -30 54 46.0	9.6	1.00	(17.7)			
J040018.4+304127		4.80e-12	04 00 19.07 +30 41 19.5	13.2	1.00	(17.2)	0.1135 ¹		Sy2
J040129.1+815320		5.04e-12	04 01 29.78 +81 53 13.3	13.1	1.00	(19.6)	0.2149 ¹		Early
J040324.1-242931		4.25e-12	04 03 24.58 -24 29 46.8	7.2	0.96	(19.5)			
J040353.1-360448	[HB89] 0402-362	2.07e-12	04 03 53.79 -36 05 01.7	1151.1	1.00	17.2	1.4170		QSO
J040748.7-121133	[HB89] 0405-123	1.05e-11	04 07 48.44 -12 11 33.4	2940.9	1.00	14.8	0.5726	Opt.var.	Blazar
J041112.1-394131		4.28e-12	04 11 11.25 -39 41 42.6	5.1	0.99	(18.8)			
J041325.4+102756 [†]		6.06e-11	04 13 25.18 +10 27 54.1	37.6	1.00	(18.8)			
J041340.2+111218	3C 109	5.60e-12	04 13 40.33 +11 12 15.0 ^O	4070.4 ^D	1.00	18.7	0.3056	N galaxy;opt.var.	
J041357.5-380543	ESO 303- G 005	7.44e-12	04 13 58.80 -38 05 51.0 ^O	4.8 ^E	0.51	15.1	0.0498	cD?	
			04 13 58.44 -38 05 34.9 ^O	4.8 ^E	0.47	(16.3)			
J041652.6+010532	[HB89] 0414+009	7.13e-11	04 16 52.88 +01 05 26.5	120.6 ^E	1.00	16.4	0.2870		BLLAC
J041717.6-055343	[HB89] 0414-060	3.24e-12	04 17 16.95 -05 53 50.6 ^R	766.7 ^E	1.00	15.9	0.7750	Opt.var.	QSO
J041733.7-115412		5.79e-12	04 17 34.91 -11 54 34.2	30.8 ^E	1.00	(18.3)			
J041938.6+022439		5.64e-11	04 19 38.87 +02 24 43.4 ^O	20.0 ^E	0.07	(17.2)			
	NGC 1550		04 19 38.02 +02 24 34.9 ^O	20.0 ^E	0.93	13.1	0.0123	SA(s)0 ⁻ :	
J042132.7-062849		6.14e-12	04 21 32.84 -06 29 06.2	15.9	1.00	(19.1)			
J042156.8-181942	APMBGC 550-098-088	2.86e-12	04 21 57.77 -18 19 34.2	41.5	1.00	16.2	0.0954 ¹		Early
J042218.2+195047	MS 0419.3+1943	9.02e-12	04 22 18.33 +19 50 55.8	8.9	1.00	20.3	0.5120		BLLAC
J042307.3-395356	PKS 0421-400	2.32e-12	04 23 08.57 -39 54 01.7	219.6 ^E	1.00	(16.6)			
J042315.1-012019	[HB89] 0420-014	3.97e-12	04 23 15.80 -01 20 32.4	2726.3	1.00	17.8	0.9150		Blazer/HPQ
J042551.3-083329 [†]	VV 760 NED01	3.14e-11	04 25 51.30 -08 33 36.8	113.0	1.00	15.5			
J042700.9-204321 [†]	ABELL 0490:[GdP94]	2.12e-12	04 26 59.93 -20 43 12.2	2.7	0.95	(16.4)			
J042704.1+071631	KUV 04244+0710	5.37e-12	04 27 04.65 +07 16 26.7	3.4 ^E	1.00	16.5	0.0900		QSO
J042900.5-323633		6.25e-12	04 29 00.21 -32 36 38.1	9.2	1.00	(18.4)			
J042936.0-025301		3.51e-12	04 29 36.01 -02 53 06.4	139.3	1.00	(19.1)			
J043055.5-213107		2.35e-12	04 30 55.05 -21 31 19.4 ^{O R}	84.0 ^D	1.00	(17.2)			
J043311.2+052112	3C 120	7.25e-11	04 33 11.03 +05 21 15.4	3439.5 ^E	1.00	14.2	0.0330	S0:	Sy1
J043337.7-131528 [†]	MCG -02-12-039	6.66e-11	04 33 37.89 -13 15 42.3	121.2	1.00	13.3	0.0330	cD;E+?	
J043429.0+712757	IRAS 04288+7121	6.91e-12	04 34 28.62 +71 28 05.2	7.9	1.00	(16.5)	0.0248 ¹		Sy1
J043518.7-262120		2.90e-12	04 35 17.74 -26 21 25.2	16.8 ^E	1.00	(18.3)			
J043622.3-102226	MRK 0618	1.85e-11	04 36 22.34 -10 22 32.7	17.3	1.00	14.5	0.0356	SB(s)b pec	Sy1
J043643.8+100323 [†]	MRC 0433+099	4.88e-12	04 36 44.32 +10 03 10.0 ^R	12.1 ^E	1.00	18.5	0.1590	sym. dbl.	
J043737.6-295411	[HB89] 0435-300	2.22e-12	04 37 36.59 -29 54 05.3	1091.4	1.00	17.5	1.3280		QSO
J043902.0+052050 [†]	PMN J0439+0520	6.81e-12	04 39 02.19 +05 20 41.5	82.4 ^E	1.00	16.0	0.2080	cD	
J043931.4-320045		2.03e-12	04 39 32.22 -32 00 48.7	10.8	1.00	(18.8)			
J044017.8-245918		4.23e-12	04 40 18.30 -24 59 36.2 ^O	12.6	0.10	(18.6)			

Table 2—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position		S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	(J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J044122.7–270811	IRAS 04392–2713	1.32e–11	04 40 18.53	–24 59 34.1 ^O	12.6	0.90	(18.5)			
J044127.8+150455		4.00e–11	04 41 22.59	–27 08 20.3	34.2	1.00	16.5	0.0835		Sy1
J044229.8–001823		4.26e–12	04 41 27.43	+15 04 54.9	13.9	1.00	(19.7)			
J044237.8–282534	[HB89] 0440–285	2.72e–12	04 42 29.97	–00 18 35.4 ^R	5.4	1.00	(19.3)			
J044255.9–263452		2.42e–12	04 42 37.66	–28 25 30.6	357.1	1.00	19.2	1.9520		QSO
	APMBGC 485+062+086		04 42 56.58	–26 35 03.8 ^O	3.5	0.16	(18.1)			
J044312.2+064811	ABELL 0504:001	4.82e–12	04 42 55.99	–26 35 09.2 ^O	3.5	0.83	16.4		S0	
J044320.8–282039		2.68e–12	04 43 11.29	+06 47 55.0	14.8	0.98	17.4	0.1161 ¹	cD?	Early
J044428.6+122113		7.99e–12	04 43 20.82	–28 20 53.3	37.6	1.00	(16.4)			
J044510.0–383824		7.99e–12	04 44 28.75	+12 21 11.4	14.0	1.00	(16.5)			
J044511.8–155118 [†]	NGC 1650	2.44e–12	04 45 10.58	–38 38 32.1	661.0 ^E	1.00	(16.6)			
J044708.2–265731	IRAS 04451–2702	5.82e–12	04 45 11.53	–15 52 12.2 ^O	90.6	0.77	13.8	0.0363	E+	
J044803.3–202741 [†]	ABELL 0514:[OWG93] A	2.50e–12	04 47 09.56	–26 57 35.0	27.2	1.00	(16.6)	0.1144		Sy1.8
J044814.8+095259 [†]	RX J0448.2+0952:001	3.28e–12	04 48 02.99	–20 26 38.7 ^{O R}	126.5 ^S	0.86	14.5	0.0764		
J044837.7–163249	PMN J0448–1632	6.65e–12	04 48 12.64	+09 53 01.0	21.4	1.00	(18.7)	0.1525 ¹		Early?
J044942.1–391052	[HB89] 0448–392	7.12e–12	04 48 37.64	–16 32 42.8	59.0	1.00	(16.4)	– ¹		BLLAC
J045108.4–232541		1.63e–12	04 49 42.29	–39 11 09.2	542.7	1.00	16.5	1.2880	Opt.var.	QSO
J045142.3–034834	MCG –01–13–025	3.44e–12	04 51 07.51	–23 25 34.9	13.3	1.00	(18.8)			
J045230.4–295329	[HB89] 0450–299	7.27e–12	04 51 41.38	–03 48 34.0	8.0	1.00	15.0	0.0159	SAB(s)0 ⁺ pec	Sy1.2
J045608.8–215906	[HB89] 0454–220	3.20e–12	04 52 30.32	–29 53 36.7	9.7	1.00	16.0	0.2860		Sy1
J045859.6–002904 [†]	NGC 1713	4.05e–12	04 56 09.00	–21 59 11.5 ^{O R}	1991.4 ^D	0.89	16.1	0.5335		QSO
J045909.6+045831	UGC 03223	4.42e–12	04 58 54.53	–00 29 20.8 ^O	6.1	0.94	13.9	0.0151	cD;E+:	
J050049.4–384038 [†]	NGC 1759	1.58e–11	04 59 09.40	+04 58 29.7	20.0	1.00	13.8	0.0156	SBa	Sy1
J050120.1–033223 [†]	NPM1G –03.0221	4.37e–12	05 00 49.15	–38 40 18.8	6.1	0.99	14.1	0.0542	E+:	
J050209.1+033149	GHIGO	4.97e–12	05 01 20.01	–03 32 38.6 ^R	309.0 ^E	1.00	15.3	0.0877 ¹		Early
J050335.6–111504		8.31e–12	05 02 09.06	+03 31 48.0	19.2	1.00	14.8	0.0160	E	Sy1
J050417.8–294434		1.32e–11	05 03 35.54	–11 15 06.1	10.4	1.00	(19.2)			
J050419.3–095622		1.26e–12	05 04 19.54	–29 44 46.6	4.1	0.99	(16.6)			
J050507.9–322007 [†]		6.58e–12	05 04 19.46	–09 56 33.4	26.3	1.00	(18.4)			
		1.83e–12	05 05 08.58	–32 19 56.3 ^O	6.4	0.08	(18.2)			
			05 05 09.37	–32 19 55.2 ^O	6.4	0.91	(16.1)			
J050534.7+041600	PMN J0505+0416	9.83e–12	05 05 34.82	+04 15 54.8	168.0	1.00	16.7	0.0272? ¹		BLLAC
J050645.3–101121		4.02e–12	05 06 44.27	–10 11 41.7 ^O	6.7	0.14	(16.7)			
			05 06 44.97	–10 11 35.1 ^O	6.7	0.86	(16.7)	0.1246 ¹		Sy1.5
J050648.5–193651		2.43e–12	05 06 47.93	–19 36 51.0 ^O	4.2	0.93	(16.5)	0.0941 ¹		Sy1.5
			05 06 49.22	–19 36 46.3 ^O	4.2	0.06	(19.2)			
J050727.6–334628		2.87e–12	05 07 27.29	–33 46 32.6	36.8	1.00	(19.1)			
J050756.6+673721	87GB 050246.4+673341	3.86e–11	05 07 55.85	+67 37 24.0	26.2	1.00	18.5	0.0065 ¹		BLLAC?
J050857.4+062633		5.40e–12	05 08 57.97	+06 26 29.1	20.7	1.00	(18.6)			
J050938.3–040037	EXO 0507.1–0404	2.99e–11	05 09 38.17	–04 00 44.0	71.1	1.00	19.0	0.3040		BLLAC
J050940.0–251356		2.11e–12	05 09 39.83	–25 14 03.0 ^O	4.3	0.82	(19.3)			
			05 09 39.88	–25 13 39.9 ^O	4.3	0.06	(18.6)			

Table 2—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position		S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	(J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
			05 09 40.46	-25 13 42.6 ^O	4.3	0.12	(19.5)			
J051324.7-332012	IRAS F05115-3323	2.15e-12	05 13 24.71	-33 20 06.8	4.8	0.98	(16.6)			
J051611.5-000807 [†]	UGC 03271	9.19e-11	05 16 11.43	-00 09 00.2	12.4	—	14.1	0.0327	Sb/pec	Sy1
J051621.5-103341		1.21e-11	05 16 20.76	-10 33 51.2 ^O	3.7	0.11	(15.8)			
			05 16 22.12	-10 33 35.7 ^O	3.7	0.18	(17.4)			
	MCG -02-14-009		05 16 21.23	-10 33 41.3 ^O	3.7	0.67	15.5	0.0287 ¹		Sy1
J051912.1-311142		1.54e-12	05 19 11.53	-31 11 39.6 ^O	4.3	0.36	(20.4)			
			05 19 11.60	-31 12 02.3 ^O	4.3	0.29	(17.5)			
			05 19 13.36	-31 11 44.1 ^O	4.3	0.29	(18.7)			
J052042.4-132855	TXS 0518-135	4.36e-12	05 20 42.06	-13 28 47.2	93.4	1.00	(19.2)			
J052135.1+671808		6.12e-12	05 21 34.60	+67 18 07.9 ^O	3.9	0.69	(17.6)	0.0147 ¹		Sy1.5
			05 21 33.68	+67 18 10.0 ^O	3.9	0.27	(15.9)			
J052224.4-072515	PMN J0522-0725	6.84e-12	05 22 23.19	-07 25 13.4	55.7	0.99	(16.5)	0.1640 ¹		Sy1
J052258.0-362729	ESO 362- G 021	1.98e-11	05 22 57.93	-36 27 29.8	15620.0	1.00	14.5	0.0553	N galaxy	BLLAC/HPQ
J052439.5-261551		1.74e-12	05 24 39.81	-26 15 51.5	20.4	1.00	(17.5)			
J052712.4+041251		8.38e-12	05 27 10.92	+04 12 34.8 ^O	7.4 ^A	0.21	(17.3)	0.1537 ¹		Early
			05 27 11.71	+04 12 55.7 ^O	7.4 ^A	0.11	(19.4)			
			05 27 11.73	+04 12 26.8 ^O	7.4 ^A	0.43	(19.2)			
			05 27 12.28	+04 12 43.5 ^O	7.4 ^A	0.16	(18.4)			
J052815.9-294305		3.14e-12	05 28 15.92	-29 43 00.8 ^R	20.6	0.97	(16.0)			
J052855.9-392802		5.39e-12	05 28 53.01	-39 28 17.9 ^O	5.2	0.36	(17.8)			
			05 28 53.61	-39 28 19.6 ^O	5.2	0.17	(19.7)			
J053224.6-113206	ABELL 0545:001	1.12e-11	05 32 25.80	-11 32 33.8 ^O	46.4	0.52	17.0	0.1540		
J053256.0-370134		3.01e-12	05 32 54.82	-37 01 39.8 ^O	10.3 ^A	0.15	(20.3)			
			05 32 54.96	-37 01 43.2 ^O	10.3 ^A	0.26	(20.2)			
			05 32 54.17	-37 01 47.3 ^O	10.3 ^A	0.20	(21.0)			
			05 32 54.31	-37 01 53.7 ^O	10.3 ^A	0.19	(20.5)			
J053317.6-361931		3.85e-12	05 33 14.99	-36 18 59.1	12.2	0.93	(16.1)			
J053629.4-334302		8.67e-12	05 36 29.11	-33 43 02.4	94.7	1.00	(17.1)			
J054426.9-251018		3.76e-12	05 44 26.45	-25 10 29.9	2.9	0.99	(16.5)			
J054702.2-390425		2.93e-12	05 47 01.51	-39 04 26.2	31.7	0.95	(16.3)			
J054738.7-315228 [†]		7.64e-12	05 47 37.38	-31 52 24.8 ^O	16.0	0.12	(16.5)			
	ABELL 3364:001		05 47 37.67	-31 52 24.5 ^O	16.0	0.82	17.2	0.1658		
			05 47 37.74	-31 52 33.1 ^O	16.0	0.05	(18.1)			
J054915.4-242550	IRAS 05472-2426	4.84e-12	05 49 14.74	-24 25 54.5	2.6	0.94	17.8	0.0449		
J055040.7-321619 [†]	[HB89] 0548-322	4.32e-11	05 50 40.68	-32 16 20.7	344.6 ^E	1.00	15.5	0.0690	Opt.var.	BLLAC
J055229.5+592842	IRAS 05480+5927	1.80e-11	05 52 28.05	+59 28 32.1	4.2	1.00	(15.8)	0.0585 ¹		Sy1
J055410.7-275724		2.09e-12	05 54 11.13	-27 57 29.9 ^O	16.1	0.69	(18.5)			
			05 54 11.54	-27 57 20.9 ^O	16.1	0.31	(21.0)			
J055711.1-372813 [†]	ESO 364- G 018	5.26e-12	05 57 12.40	-37 28 38.0	309.5	1.00	15.0		E+2	
J055802.0-382001	IRAS F05563-3820	2.89e-12	05 58 02.20	-38 20 02.4	34.9	1.00	17.0	0.0344		Sy1
J055806.6-383829	EXO 0556.4-3838	1.64e-11	05 58 06.10	-38 38 29.5 ^R	105.7 ^E	1.00	17.1			Sy1/BLLAC

Table 2—Continued

(1) <i>ROSAT</i> BSC 1RXS name	(2) NED Common name	(3) S_x ergs s ⁻¹ cm ⁻²	(4) Position α (J2000) δ	(5) $S_{1.4\text{GHz}}$ mJy	(6) ID P	(7) m_B	(8) Redshift	(9) NED Morphology	(10) Sp. Class.
J055810.8–212344		3.00e–12	05 58 10.23 –21 23 39.5 ^O 05 58 10.93 –21 23 35.7 ^O	3.1 3.1	0.41 0.52	(16.6) (14.6)			
J055958.9+640952		5.36e–12	05 59 59.18 +64 09 59.2	32.3	1.00	(18.8)			
J060427.5–225542		3.64e–12	06 04 27.46 –22 55 40.0 ^R	5.3	0.99	(16.5)			
J060548.1–275439	IRAS F06038–2754	7.24e–12	06 05 49.21 –27 54 44.7	7.5	1.00	(16.4)			
J060552.7–351759 [†]	PKS 0604–352	1.06e–11	06 05 54.07 –35 18 07.7	1337.4	0.97	17.0		S0	
J060714.2–251855		4.14e–12	06 07 14.29 –25 18 59.5	12.0	1.00	(18.3)			
J061117.7–192255		5.38e–12	06 11 17.34 –19 23 00.0 ^R	117.2 ^S	1.00	(16.6)			
J061238.7–164839		7.14e–11	06 12 38.84 –16 48 33.5	8.4	0.99	(11.0)			
J061324.2–290029	IRAS F06114–2859	4.82e–12	06 13 24.36 –29 00 21.8	28.8	1.00	(14.3)			
J061949.8+573553		1.16e–11	06 19 49.23 +57 35 54.1	2.7	1.00	(19.5)			
J062150.0–341140		3.57e–12	06 21 49.55 –34 11 54.1	8.6 ^E	1.00	(19.1)			
J062221.4–260537	PMN J0622–2605	3.52e–12	06 22 22.07 –26 05 44.2	106.4	1.00	(17.2)			
J062233.8–231743	IRAS 06205–2316	6.11e–12	06 22 33.62 –23 17 44.2	4.3	0.97	10.4	0.0380		Sy1
J062334.6+644542		9.54e–12	06 23 35.95 +64 45 38.2	4.1	1.00	(16.5)	0.0864 ¹		Sy1.5
J062349.8–275333		3.46e–12	06 23 49.22 –27 53 27.9	7.7	1.00	(14.6)			
J062422.3–294449		7.95e–12	06 24 22.65 –29 44 50.5	15.5	1.00	(19.3)			
J062707.5–352917 [†]	PKS 0625–35	1.43e–11	06 27 07.12 –35 29 17.1 ^R	2372.5 ^T	1.00	16.5	0.0546	Elliptical	
J062736.9–260103		1.20e–11	06 27 37.15 –26 01 01.5	13.9	1.00	(16.7)			
J062910.4+460637		6.88e–12	06 29 15.32 +46 06 18.1 ^R	3.2	0.98	(21.5)			
J063248.0+634026	UGC 03478	1.31e–11	06 32 46.95 +63 40 25.6	12.9	1.00	13.6	0.0128	Sb	Sy1.2
J064326.5+421420 [†]	B3 0639+423	3.57e–11	06 43 26.76 +42 14 18.8 ^R	110.1 ^S	0.93	(17.0)	0.0893 ¹		Early
J065549.7+395957		4.38e–12	06 55 49.72 +39 59 43.6 ^O	5.1	0.05	(16.1)			
	UGC 03601		06 55 49.45 +40 00 01.4 ^O	5.1	0.95	14.8	0.0175 ¹		Sy1.2
J065611.5+423702	NPM1G +42.0131	4.02e–12	06 56 10.61 +42 37 01.6	933.6	0.99	16.9	0.0562 ¹		Early
J065938.5+541136	MRK 0374	2.55e–11	06 59 38.44 +54 11 45.0	14.5	1.00	15.0	0.0435	SBab	Sy1
J070427.0+631856 [†]		1.07e–11	07 04 29.59 +63 18 29.8 ^{OR}	36.6 ^E	0.06	(18.9)			
	KUG 0659+633		07 04 28.78 +63 18 39.1 ^{OR}	36.6 ^E	0.64	17.0	0.0942 ¹		Early
J070702.9+270650		7.04e–12	07 07 02.86 +27 06 49.7	46.4	0.97	(16.4)	0.0623 ¹		Early
J070713.5+643558	VII Zw 118	2.50e–11	07 07 13.34 +64 35 57.9	2.8	1.00	14.5	0.0797		Sy1
J070907.6+483657 [†]	NGC 2329	3.71e–12	07 09 07.83 +48 36 53.2	756.1 ^T	1.00	13.5	0.0193	S0–	
J071006.0+500243 [†]	87GB 070614.9+500820	4.39e–12	07 10 05.42 +50 03 08.1 ^R	113.9 ^E	1.00	16.1	0.0601 ¹		Early
J071030.0+590808	87GB 070609.2+591323	3.30e–11	07 10 30.65 +59 08 17.8	159.2	1.00	18.4	0.1250		BLLAC
J071148.0+321902	PGC 020369	1.17e–11	07 11 47.68 +32 18 35.7	65.7	1.00	15.8	0.0672 ¹		Early
J071218.9+571930	RX J0712.3+5719	2.96e–12	07 12 18.70 +57 19 22.0	7.9	1.00	19.9			BLLAC
J071339.7+382043	IRAS F07102+3825	4.11e–12	07 13 40.23 +38 20 38.1	11.3	1.00	(16.5)	0.1227 ¹		Sy1
J071435.7+740809	87GB 070816.7+741327	3.27e–12	07 14 35.93 +74 08 09.7	107.7 ^E	1.00	16.6	0.3710		QSO
J071726.9+440557	B3 0713+441	5.51e–12	07 17 26.71 +44 05 04.3	220.4	0.88	(16.6)	0.0652 ¹		Early
J071800.7+440527	IRAS F07144+4410	2.42e–11	07 18 00.69 +44 05 27.4	50.8	1.00	15.5	0.0614		Sy1
J072019.1+234904		6.64e–12	07 20 18.64 +23 49 04.0 ^F	6.6	1.00	(17.2)	0.1660 ¹		Early
J072153.2+712031	[HB89] 0716+714	4.87e–12	07 21 53.10 +71 20 36.7	727.2	1.00	15.5			BLLAC/HPQ
J072217.4+303052		4.77e–12	07 22 17.56 +30 30 52.4	6.5	1.00	(16.5)	0.0998 ¹		Sy1

Table 2—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position		S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	(J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J072927.4+243625	TXS 0726+247	6.01e-12	07 29 27.91	+24 36 22.8	68.7 ^E	1.00	(17.5)	0.1630 ²		HII
J073026.0+330727		3.28e-12	07 30 26.13	+33 07 21.9	9.2	1.00	(17.0)	0.1130 ¹		Early
J073152.4+280423	87GB 072846.0+281055	5.40e-12	07 31 52.75	+28 04 32.3	72.1	1.00	(17.6)	0.2495 ^{?1}		BLLAC?
J073221.5+313750 [†]		1.05e-11	07 32 20.44	+31 37 57.4	5.9	1.00	(17.6)			
J073329.3+351546	MG2 J073334+3515	4.10e-12	07 33 29.58	+35 15 42.4	102.3	1.00	15.4	0.1769 ¹		Early?
J073624.8+392608		1.12e-11	07 36 23.16	+39 26 17.4 ^F	3.8	1.00	(16.3)	0.1177 ¹		Sy1
J073657.0+584610		3.65e-12	07 36 56.46	+58 46 04.6 ^O	4.1	0.20	(19.2)			
	MRK 0009		07 36 57.05	+58 46 13.3 ^O	4.1	0.80	15.3	0.0399	S0 pec?	Sy1
J074058.5+552533	UGC 03957	1.40e-11	07 40 58.20	+55 25 38.0	53.2	1.00	(15.4)	0.0338 ¹		Early
J074144.8+741444 [†]	4C +74.13	6.23e-12	07 41 44.51	+74 14 39.0 ^O	22.7 ^E	0.73	(18.5)		cD	
			07 41 44.77	+74 14 32.9 ^O	22.7 ^E	0.27	(19.7)			
J074232.9+494830	UGC 03973	6.89e-11	07 42 32.84	+49 48 34.1	22.2	1.00	13.9	0.0222	SBb	Sy1.2
J074238.0+092213 [†]		4.58e-12	07 42 39.70	+09 22 26.4 ^O	5.1	0.64	(16.7)			
	LEDA 078089		07 42 40.62	+09 22 20.7 ^O	5.1	0.29	15.0	0.0635 ¹		Early
J074405.6+743356	MS 0737.9+7441	1.20e-11	07 44 05.44	+74 33 58.1	23.3	1.00	16.9	0.3150		BLLAC
J074541.2+314249	[HB89] 0742+318	3.22e-12	07 45 41.66	+31 42 56.6 ^F	1359.5 ^T	1.00	16.0	0.4611		Sy1
J074721.9+090607	RGB J0747+090	4.56e-12	07 47 22.22	+09 05 53.3	40.3	1.00	(17.1)	— ¹		BLLAC
J074729.4+605601	UGC 04013	1.68e-11	07 47 28.88	+60 56 00.4	6.3	1.00	13.3	0.0293	SBbc	Sy1
J074906.2+451039	B3 0745+453	3.18e-12	07 49 06.41	+45 10 33.8	147.8 ^E	1.00	16.3	0.1900		BLRG
J074929.4+745142	87GB[BWE91] 0743+7458	9.38e-12	07 49 29.67	+74 51 45.4	44.8	1.00	17.5	— ¹		BLLAC
J075122.1+551156		4.18e-12	07 51 22.35	+55 12 08.9	2.4	1.00	(16.3)	— ¹		BLLAC
J075124.3+173044		3.46e-12	07 51 25.08	+17 30 50.6	10.4	1.00	(18.2)	0.1863 ¹		Early
J075243.6+455653	NPM1G +46.0092	5.76e-12	07 52 43.60	+45 56 56.0	604.0 ^T	1.00	16.2	0.0517 ¹		Early
J075322.4+292154		2.96e-12	07 53 24.61	+29 21 31.9 ^F	4.1	0.96	(18.4)			
J075408.7+431611		3.45e-12	07 54 07.90	+43 16 09.5	29.6 ^T	1.00	(16.7)	0.3477 ¹		Sy1?
J075706.7+095647	[HB89] 0754+100	2.22e-12	07 57 06.60	+09 56 34.9	956.9	1.00	14.5	0.6600	Opt.var.	BLLAC
J075819.5+421938	IRAS F07548+4227	4.19e-12	07 58 19.66	+42 19 43.5	3.7	1.00	16.2	0.2100		Sy1
J075936.1+132123		8.64e-12	07 59 35.73	+13 21 20.5	5.1	1.00	(21.0)			
J080026.2+101313	IRAS 07577+1021	4.03e-12	08 00 26.84	+10 13 15.6	5.7	0.99	16.4	0.0478		Sy1.5
J080132.3+473618	87GB 075755.8+474440	8.58e-12	08 01 31.88	+47 36 16.9 ^R	123.1 ^T	1.00	15.5	0.1583		BLRG
J080323.1+481622		2.95e-12	08 03 23.02	+48 16 18.6	12.6	1.00	(20.1)			
J080525.8+753424	RX J0805.4+7534	7.57e-12	08 05 26.61	+75 34 25.2	52.6	1.00	18.1	0.1210		BLLAC
J080625.4+172556		1.95e-12	08 06 24.94	+17 25 04.1	40.9	1.00	(16.9)	0.1038 ¹		Early?
J080625.5+593105	87GB 080212.8+593933	4.95e-12	08 06 25.94	+59 31 06.8	60.9	1.00	16.2	— ¹		BLLAC
J080639.9+724831	87GB 080101.9+725723	3.18e-12	08 06 38.94	+72 48 20.1	50.1	1.00	16.1	0.0977 ¹		Sy2
J080938.5+345544	MG2 J080937+3455	9.21e-12	08 09 38.79	+34 55 35.0	286.4 ^S	1.00	17.0			
J080949.2+521855	87GB 080601.8+522753	1.89e-11	08 09 49.23	+52 18 58.1	182.8	1.00	15.3	0.1380		BLLAC
J081021.3+421657 [†]	CGCG 207-040	4.01e-12	08 10 23.25	+42 16 24.6	11.4	0.97	15.6	0.0632 ¹		Early
J081059.0+760245	PG 0804+761	2.93e-11	08 11 00.94	+76 02 45.6	3.6	1.00	15.1	0.1000		Sy1
J081112.3+700300 [†]	87GB 080602.4+701201	3.31e-12	08 11 12.85	+70 02 27.9	81.2	0.98	(18.6)	0.2230	cD	
J081201.8+023735	PMN J0811+0237	1.76e-11	08 12 01.51	+02 37 32.6	123.3 ^E	1.00	14.5			BLLAC
J081517.8+460429	KUG 0811+462	3.84e-12	08 15 16.68	+46 04 29.2	7.9	0.99	15.2	0.0410	Spiral	Sy1.5

Table 2—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position		S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	(J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J081523.2+015500	[HB89] 0812+020	4.10e-12	08 15 22.82	+01 55 05.2	1755.9	1.00	17.1	0.4020	Opt.var.	QSO
J081527.2-030819	3C 196.1	6.66e-12	08 15 27.81	-03 08 26.6	1963.2	1.00	18.5	0.1980	Radio galaxy	
J081624.6+573910	87GB 081222.1+574811	3.26e-12	08 16 22.68	+57 39 09.2	100.0	1.00	16.8	— ¹		BLLAC
J081815.0+012215		8.53e-12	08 18 14.66	+01 22 26.8 ^O	21.3	0.85	(16.5)	0.0890 ¹		Sy1.9
			08 18 14.99	+01 22 31.2 ^O	21.3	0.15	(18.2)			
J081917.6-075620		3.26e-12	08 19 17.51	-07 56 28.1	7.2	1.00	(18.9)			
J081917.9+642943	MCG +11-10-073	4.13e-12	08 19 16.46	+64 29 53.2 ^R	15.6	0.97	16.0	0.0390		Sy1
J081926.6+633741	KOS NP6 038	4.49e-12	08 19 25.81	+63 37 26.4	35.3	1.00	15.4	0.1191 ¹		Early
J081929.5+704221 [†]	Holmberg II	5.11e-12	08 19 28.86	+70 42 48.4 ^{OR}	9.9 ^E	0.85	11.1	0.0005	Im	
J082209.5+470601 [†]	RGB J0822+470	8.10e-12	08 22 09.54	+47 05 52.6	54.3	1.00	15.9	0.1273 ¹		HII
J082550.2+030930	PKS 0823+033	3.73e-12	08 25 50.37	+03 09 24.8	1400.6	1.00	18.5	0.5060		BLLAC/HPQ
J082705.9-070841	PMN J0827-0708	5.98e-12	08 27 06.10	-07 08 46.4	174.0	1.00	(17.6)	— ¹		BLLAC?
J082804.7+444602		4.22e-12	08 28 05.83	+44 46 00.7 ^F	34.9	1.00	(17.8)			
J082904.5+175415	MG2 J082906+1755	3.34e-12	08 29 04.60	+17 54 14.5	336.1 ^E	1.00	13.9	0.0894 ¹		Early
J083014.6-094455		4.10e-12	08 30 15.09	-09 44 56.6	12.8	1.00	(19.0)			
J083051.5+241053	[HB89] 0827+243	2.76e-12	08 30 52.11	+24 10 58.7	739.1	1.00	17.3	0.9390		Blazar/LPQ
J083201.1+461438		3.48e-12	08 32 01.78	+46 14 37.7 ^O	3.3 ^E	0.77	(18.3)			
	RX J0832.0+4614		08 32 02.15	+46 14 25.9 ^O	3.3 ^E	0.19	16.9	0.0459 ¹		Sy1
J083225.1+370737	FIRST J083225.3+370736	1.04e-11	08 32 25.22	+37 07 36.5	8.5	1.00	17.0	0.0906		Sy1.2
J083251.9+330011	RX J0832.8+3300	3.38e-12	08 32 51.49	+33 00 09.8	4.3	1.00	20.7	0.6710		QSO
J083454.3+553417	4C +55.16	2.58e-12	08 34 54.91	+55 34 21.0	8283.6	1.00	19.0	0.2420	Radio galaxy	
J083539.1-040508	NGC 2617	4.65e-12	08 35 38.84	-04 05 20.0	29.5	1.00	14.0	0.0143		Sy1.8
J083658.3+442612	[HB89] 0833+446	9.07e-12	08 36 58.58	+44 26 02.1	7.1	1.00	15.6	0.2550		QSO
J083811.0+245336 [†]	NGC 2622	5.03e-12	08 38 10.93	+24 53 43.0	65.6	1.00	15.0	0.0286	SBb	Sy1.8
J083950.7-121424 [†]	[HB89] 0837-120	1.20e-11	08 39 50.59	-12 14 34.2 ^O	1863.7 ^D	1.00	15.8	0.1976	Opt.var.	QSO
J084125.1+705342	[HB89] 0836+710	1.06e-11	08 41 24.46	+70 53 41.4	3823.7	1.00	16.5	2.1720		Blazar/LPQ
J084151.2+231845	RGB J0841+233B	3.32e-12	08 41 53.87	+23 19 54.9	192.0	1.00	(17.2)	0.4883 ¹		QSO
J084203.4+401830	87GB 083846.8+402925	5.13e-12	08 42 03.74	+40 18 30.8	22.7	1.00	15.6	0.1515 ¹		Sy1
J084206.6+075936	RX J0842.1+0759	9.10e-12	08 42 05.41	+07 59 26.7	17.2	1.00	17.7	0.1339 ¹		Sy1
J084255.9+292752 [†]	FIRST J084255.9+292727	3.62e-12	08 42 55.96	+29 27 27.6	25.2	1.00	(18.0)	0.1932		
J084406.3-140211	IRAS 08417-1351	9.70e-12	08 44 06.34	-14 02 10.9	7.8	1.00	(15.0)	0.0278		Sy1.5
J084553.0-024101		3.42e-12	08 45 53.41	-02 41 00.5	55.5	1.00	(16.6)	0.4690 ¹		QSO
J084600.6+070441	PMN J0846+0704	4.49e-12	08 45 59.95	+07 04 23.4	336.3 ^T	1.00	(16.6)	0.3418 ¹		QSO
J084628.5-121405	IRAS F08440-1202	1.26e-11	08 46 28.53	-12 14 10.4	15.7	1.00	(16.5)	0.1076 ¹		Sy1.8
J084713.3+113349	RGB J0847+115	2.31e-11	08 47 12.94	+11 33 50.5	33.2	1.00	16.6			BLLAC
J084755.9-070306	PMN J0847-0703	4.23e-12	08 47 56.78	-07 03 17.9	183.9	1.00	(16.4)	— ¹		BLLAC?
J084829.3+811145		2.26e-12	08 48 28.21	+81 11 49.8	12.9	1.00	(17.4)			
J084935.5-172116	TXS 0847-171	3.84e-12	08 49 34.99	-17 21 20.4	149.2	1.00	(21.5)			
J085011.2+360421	[CGH98] J0850+3604	2.44e-12	08 50 07.75	+36 04 21.7	21.7	0.98	(19.6)			
J085152.5+522828	RX J0851.8+5228:001	2.52e-12	08 51 51.39	+52 28 26.2 ^F	4.7 ^E	1.00	17.5	0.0640 ¹		Sy2
J085407.7+440840	B3 0850+443	3.16e-12	08 54 09.78	+44 08 30.7	79.8	1.00	19.0			BLLAC
J085439.9+174111	MRK 1220	4.15e-12	08 54 39.25	+17 41 23.6	5.8	1.00	16.5	0.0649	Compact	Sy1

Table 2—Continued

(1) <i>ROSAT</i> BSC 1RXS name	(2) NED Common name	(3) S_x ergs s ⁻¹ cm ⁻²	(4) Position α (J2000) δ	(5) $S_{1.4\text{GHz}}$ mJy	(6) ID P	(7) m_B	(8) Redshift	(9) NED Morphology	(10) Sp. Class.
J085451.5+200646	OJ +287	3.52e-12	08 54 48.87 +20 06 30.7	1512.3	1.00	14.0	0.3060	Opt.var.	BLLAC
J085610.0+541826	87GB 085229.1+543028	2.13e-12	08 56 08.20 +54 18 55.9	58.6	1.00	15.0	0.2540? ¹		Early
J085617.7-013809		2.53e-12	08 56 18.77 -01 38 16.7 ^O	2.9	0.07	(19.5)			
	CGCG 005-037		08 56 17.85 -01 38 07.6 ^O	2.9	0.93	15.5	0.0596 ¹		Sy2
J085916.5+834450		3.41e-12	08 59 10.12 +83 45 03.6	10.2 ^E	1.00	(17.6)			
J085920.6+004716	RGB J0859+007	5.56e-12	08 59 20.48 +00 47 13.2	39.0	1.00	19.0			QSO
J085930.1+745510	RX J0859.5+7455	2.44e-12	08 59 30.51 +74 54 59.8	6.1	1.00	16.8	0.2520 ¹		Sy1
J090315.3+405609	FIRST J090314.7+405559	4.20e-12	09 03 14.69 +40 56 01.0	38.2	1.00	18.7	0.1882 ¹		Early
J090533.4+184011	RX J0905.5+1840:099	3.05e-12	09 05 33.53 +18 40 02.9	18.7	0.98	(16.2)	0.1228 ¹		Early
J090604.9+194153	[HB89] 0903+198	2.57e-12	09 06 03.53 +19 41 39.7	16.8	1.00	16.4	1.2060	Opt.var.	QSO
J090632.0+164619	[HB89] 0903+169	3.82e-12	09 06 31.88 +16 46 13.0	1587.1 ^E	1.00	18.3	0.4121	Opt.var.	QSO
J090802.1-095928	NPM1G -09.0307	3.22e-12	09 08 02.38 -09 59 40.8	652.5 ^S	1.00	15.0	0.0533	E	
J090953.9+310558	MG2 J090953+3104	5.94e-12	09 09 53.75 +31 06 01.6	195.5	1.00	15.0	0.2740 ¹		BLLAC
J091036.6-103501	PMN J0910-1034	3.28e-12	09 10 36.06 -10 34 59.3	381.0 ^E	1.00	15.5	0.0921		
J091037.2+332920 [†]	TON 1015	2.40e-12	09 10 37.24 +33 29 25.0	105.6	1.00	16.0	— ¹		BLLAC
J091133.8+442305	B3 0908+445	1.52e-12	09 11 33.74 +44 22 49.6	446.9	1.00	17.2	0.2976 ¹		Sy1
J091230.8+155531	ABELL 0763:[OWB92] A	6.93e-12	09 12 30.72 +15 55 26.9 ^O	29.5	0.65	16.5	0.2121 ¹		Early
			09 12 31.04 +15 55 24.6 ^O	29.5	0.35	(15.6)			
J091300.4-210315		3.81e-11	09 13 00.38 -21 03 22.4	373.3 ^S	1.00	17.3			BLLAC
J091324.6+813318		3.09e-12	09 13 20.44 +81 33 06.2	4.9	1.00	(19.5)			
J091345.4+474208		2.05e-12	09 13 44.69 +47 42 10.5	17.5 ^E	1.00	(16.1)	0.0511 ¹		Early
J091346.0+405620	IRAS 09104+4109	1.14e-12	09 13 45.43 +40 56 30.7	16.7	1.00	18.0	0.4420		Sy2
J091401.0+050743	[HB89] 0911+053	2.87e-12	09 14 01.95 +05 07 47.7 ^R	456.0 ^T	1.00	17.4	0.3030		QSO
J091552.2+293335	[HB89] 0912+297	9.51e-12	09 15 52.40 +29 33 24.1	342.0	1.00	16.0		Opt.var.	BLLAC/HPQ
J091651.8+523829	87GB 091315.6+525108	5.30e-12	09 16 51.93 +52 38 28.5 ^{R^F}	157.6 ^S	1.00	19.5	0.1900		BLLAC
J091714.7-034302		2.56e-12	09 17 14.59 -03 43 15.3	32.1	1.00	(18.4)	— ¹		Early?
J091807.6-120543	Hydra A	6.77e-11	09 18 05.78 -12 05 41.3	40850.9 ^E	1.00	13.9	0.0538	(R')SA0 ⁻ :	Sy3
J091826.2+161825	MRK 0704	1.76e-11	09 18 26.05 +16 18 21.2	6.6	0.98	15.4	0.0299	SBa	Sy1.5
J091926.6+110659	RGB J0919+111	2.26e-12	09 19 25.69 +11 06 57.4	35.3	1.00	18.1			
J091949.1+334532 [†]	NGC 2832	2.71e-12	09 19 46.83 +33 44 58.8 ^O	5.5 ^E	1.00	12.9	0.0229	E+2::cD	
J092057.2-225716		7.82e-12	09 20 57.39 -22 57 21.2	14.3	1.00	(18.0)			
J092057.9+444200	[HB89] 0917+449	1.54e-12	09 20 58.48 +44 41 53.7	1017.6	1.00	19.0	2.1800		Blazar
J092108.3+453854	3C 219	1.66e-12	09 21 08.66 +45 38 56.9 ^O	5928.5 ^D	0.79	17.2	0.1744	Radio galaxy	
			09 21 09.12 +45 38 51.4 ^O	5928.5 ^D	0.13	(18.9)			
			09 21 10.59 +45 38 53.2 ^O	5928.5 ^D	0.08	(19.7)			
J092338.8-213541	NPM1G -21.0236	1.11e-11	09 23 39.09 -21 35 47.5	268.3	1.00	16.5	0.0530	Radio galaxy	
J092343.0+225437	CGCG 121-075	3.72e-11	09 23 43.03 +22 54 30.3	10.6	0.98	15.5	0.0330		Sy1
J092401.1+053350		4.75e-12	09 24 01.26 +05 33 42.8	7.5	1.00	(18.9)			
J092406.1+141002 [†]	ABELL 0795:[WKS75] G1	8.62e-12	09 24 05.31 +14 10 22.0	117.3	1.00	(16.9)	0.1364		
J092512.3+521716	PG 0921+525	2.30e-11	09 25 12.67 +52 17 08.6	10.1	1.00	16.0	0.0353	Pair?	Sy1
J092554.3+400414	FIRST J092554.7+400414	2.42e-12	09 25 54.79 +40 04 14.4 ^R	77.0 ^T	1.00	19.3	0.4700 ¹		Sy1
J092702.8+390221	[HB89] 0923+392	3.04e-12	09 27 03.04 +39 02 20.7	2885.1	1.00	17.9	0.6948	Opt.var.	LPQ

Table 2—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position		S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	(J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J092710.8+532741		1.80e-12	09 27 10.58	+53 27 31.6 ^{OF}	5.2	1.00	(17.8)	0.2082 ¹		Early
J092752.1-203437	[HB89] 0925-203	4.85e-12	09 27 52.05	-20 34 53.8	842.9 ^E	1.00	16.4	0.3480	Opt.var.	Sy1
J092804.2+744714	87GB 092308.0+745942	1.94e-12	09 28 02.86	+74 47 18.5 ^R	151.5 ^S	1.00	19.7	0.6380		BLLAC
J093037.1+495028	RGB J0930+498	2.69e-11	09 30 37.78	+49 50 26.4	21.4	1.00	17.2	0.1880		BLLAC
J093056.2+393332	FIRST J093056.8+393335	1.35e-12	09 30 56.94	+39 33 37.2	13.0 ^E	1.00	21.0	0.6380		BLLAC
J093318.2-171440		6.70e-12	09 33 18.11	-17 14 40.8	20.0	1.00	(16.7)	0.3132 ¹		QSO
J093430.2-172117		8.02e-12	09 34 30.26	-17 21 23.7	54.4 ^T	1.00	(18.1)	0.2500 ¹		Early
J093527.4+261714	RX J0935.4+2617	2.72e-12	09 35 27.12	+26 17 09.9 ^F	2.7	1.00	17.8	0.1220 ¹		Sy1
J093703.7+361555	IRAS F09339+3629	1.26e-12	09 37 03.65	+36 15 38.9	2.5	1.00	17.7	0.1795 ¹		Sy1
J093711.5+500848	87GB 093349.0+502204	1.31e-12	09 37 12.33	+50 08 52.5	167.1	1.00	18.0	0.2760		
J094020.7+614837	RX J0940.3+6148	2.49e-12	09 40 22.48	+61 48 25.1	12.8	1.00	18.0	0.2106 ¹		BLLAC
J094104.0+385400	[HB89] 0937+391	1.56e-12	09 41 04.18	+38 53 49.8	689.1 ^E	1.00	17.5	0.6180		QSO
J094202.2+085758	RX J0942.0+0857	2.79e-12	09 42 00.45	+08 57 43.9 ^O	7.2 ^A	0.10	(19.3)			
			09 42 00.81	+08 58 09.6 ^O	7.2 ^A	0.51	(19.6)			
	ZwCl 0939.3+0908:G1		09 42 01.14	+08 58 19.7 ^O	7.2 ^A	0.15	18.3	0.2070		
			09 42 01.43	+08 58 16.4 ^O	7.2 ^A	0.15	(19.4)			
J094204.0+234106	CGCG 122-055	2.25e-12	09 42 04.90	+23 41 05.5	5.9	1.00	15.3	0.0214		
J094211.6-224027		3.09e-12	09 42 14.83	-22 40 04.6 ^O	8.6 ^E	0.51	(16.2)			
J094355.3-070944		2.66e-12	09 43 55.39	-07 09 52.4	7.1	1.00	(18.8)			
J094436.5-263353		9.32e-12	09 44 36.45	-26 33 52.5 ^O	4.3	0.31	(18.7)			
			09 44 37.05	-26 33 56.1 ^O	4.3	0.65	(16.4)			
J094458.6-134707		4.07e-12	09 44 59.06	-13 47 42.2 ^{OR}	51.7 ^E	0.21	(18.9)			Early
			09 44 59.40	-13 47 51.8 ^{OR}	51.7 ^E	0.75	(17.4)	0.1782? ¹		Early
J094541.9-141927	NGC 2992	8.47e-12	09 45 42.00	-14 19 34.6	226.6	1.00	13.1	0.0077	Sa pec	Sy2
J094553.9+423854	IRAS F09427+4252	1.26e-12	09 45 54.49	+42 38 19.0 ^F	3.2	0.85	16.3	0.0740		Sy2
J094620.5+010459	NGC 2967 GROUP	4.17e-12	09 46 19.94	+01 04 53.4	14.9	1.00	(19.5)			
J094653.0+132000	MS 0944.1+1333	5.90e-12	09 46 52.73	+13 19 59.1	8.0	1.00	16.0	0.1310		
J094703.7+472143	IRAS F09438+4735	1.49e-12	09 47 05.02	+47 21 39.1	3.4	1.00	18.1	0.5410		QSO
J094709.2-254056		3.54e-12	09 47 09.44	-25 41 00.1	41.9	1.00	(17.0)			
J094713.2+762317	RBS 0797	4.62e-12	09 47 12.93	+76 23 12.8	22.4	1.00	19.2	0.3540		LINER
J094733.2+100522	RX J0947.5+1005	5.77e-12	09 47 33.77	+10 04 58.9	4.2 ^E	0.99	16.0	0.1392 ¹		Sy1
J094740.4-305655 [†]	ESO 434- G 040	9.18e-12	09 47 40.21	-30 56 57.4	14.6	1.00	14.1	0.0083	(RL)SA(l)0 ⁰	Sy2
J094953.8+170803		4.03e-12	09 49 52.74	+17 07 36.2	4.5 ^E	0.99	(21.5)			
J095154.9-064916	NGC 3035	6.95e-12	09 51 55.29	-06 49 29.3	6.2	1.00	13.5	0.0145	SB(rs)bc	Sy1
J095214.4+393627	FIRST J095214.7+393615	1.71e-12	09 52 13.98	+39 36 08.5	3.0	0.97	19.6			
J095225.8+750216	RX J0952.4+7502	4.20e-12	09 52 23.77	+75 02 13.2	12.2	1.00	19.9	0.1813 ¹		Early
J095303.4-084003	PMN J0953-0840	3.96e-12	09 53 02.70	-08 40 18.4	85.2	1.00	(17.2)	— ¹		BLLAC
J095331.5+253939	RX J0953.5+2539	1.92e-12	09 53 31.73	+25 39 45.6	3.7	0.97	17.1	0.0441 ¹		Sy1
J095340.4+014154		2.69e-12	09 53 41.27	+01 42 05.2	10.1	1.00	17.1	0.0983 ¹		Early
J095406.7+212250	4C +21.26	2.02e-12	09 54 07.02	+21 22 35.9 ^O	960.2 ^D	1.00	20.0	0.2954 ¹		QSO
J095410.8+491457	MS 0950.9+4929	2.03e-12	09 54 09.90	+49 15 01.1	2.7	1.00	19.3	0.2070		BLLAC
J095508.2+355054	FIRST J095507.8+355100	1.26e-12	09 55 08.04	+35 51 02.5	7.8	1.00	(19.8)			

Table 2—Continued

(1) ROSAT BSC 1RXS name	(2) NED Common name	(3) S _x ergs s ⁻¹ cm ⁻²	(4) Position α (J2000) δ	(5) S _{1.4GHz} mJy	(6) ID P	(7) m _B	(8) Redshift	(9) NED Morphology	(10) Sp. Class.
J095513.8–212301		3.44e–12	09 55 13.61 –21 23 03.2	60.2	1.00	(16.1)			
J095534.7+690338	MESSIER 081	2.65e–11	09 55 33.28 +69 03 55.0	88.9 ^E	1.00	7.9		SA(s)ab	Sy1.8/LINER
J095539.4+453218	[HB89] 0952+457	1.45e–12	09 55 39.94 +45 32 14.6	57.4	1.00	16.7	0.2590		QSO
J095550.4+694052	MESSIER 082	2.38e–11	09 55 51.88 +69 40 46.1	6204.9 ^E	1.00	9.3	0.0007	I0	HII
J095557.0–095856		4.52e–12	09 55 57.42 –09 59 02.4	21.0	0.98	(17.0)	0.1618 ¹		Early
J095627.2–095720		4.24e–12	09 56 28.14 –09 57 17.8	94.5	1.00	(17.6)	0.1585? ¹		Early
J095630.6–045259		2.36e–12	09 56 29.60 –04 53 33.4 ^O	3.0	0.07	(12.5)			
	IRAS F09539–0439		09 56 30.16 –04 53 16.7 ^O	3.0	0.80	(16.4)	0.1565 ¹		Sy1
			09 56 31.13 –04 53 22.4 ^O	3.0	0.08	(19.3)			
J095708.1+243319	RX J0957.1+2433	2.81e–12	09 57 07.72 +24 33 22.9	6.6	1.00	17.0	0.0819 ¹		Sy1.5
J095806.4–031729		2.61e–12	09 58 06.14 –03 17 38.2	7.5	1.00	(18.8)			
J095821.8–110344 [†]	TXS 0955–108	1.15e–11	09 58 21.84 –11 03 47.4	69.3 ^E	1.00	(16.9)	0.1610 ¹		Early
J095929.8+212340	87GB 095643.1+213755	8.05e–12	09 59 30.04 +21 23 19.2	40.8 ^E	1.00	17.4	— ¹		BLLAC?
J095942.1–311300	RX J0959.7–3113	2.12e–11	09 59 42.70 –31 13 03.4	8.3	0.96	(10.9)	0.0370		Sy1
J095949.3+222333	RGB J0959+224	2.58e–12	09 59 47.12 +22 24 12.3	48.0	0.95	15.6	0.2449 ¹		Early
J095955.4+130246	NGC 3080	4.90e–12	09 59 55.99 +13 02 47.7	3.2	1.00	14.3	0.0354	Sa	Sy1
J100028.9+440910	RX J1000.5+4409:002	2.27e–12	10 00 31.01 +44 08 43.0 ^F	4.4	1.00	(19.1)	0.1530		
J100121.5+555351 [†]	[YGK80] 0957+561G1	1.24e–12	10 01 21.07 +55 53 55.8	551.9	1.00	18.5	0.3900	cD	
J100200.3–080939	IRAS 09595–0755	6.51e–12	10 01 59.98 –08 09 41.0	12.8	1.00	15.2	0.0152	Ring galaxy	Sy1
J100235.9+324218	NGC 3099	2.69e–12	10 02 36.57 +32 42 26.2	10.5	0.99	15.4	0.0504 ¹		Early
J100508.4+341424	FIRST J100507.9+341424	1.46e–12	10 05 07.29 +34 14 31.3	3.0	0.98	(16.7)	0.1620 ²		Sy1
J100542.2+433244	IRAS 10026+4347	7.26e–12	10 05 41.03 +43 32 26.6	3.3	0.99	16.5	0.1772 ¹		Sy1
J100554.9–230318		4.01e–12	10 05 55.81 –23 03 27.9	7.0	1.00	(13.5)			
J100639.1+255450 [†]	PGC 029375	3.71e–12	10 06 38.82 +25 54 45.1	73.1	1.00	16.5	0.1165		
J100656.9+345446	FIRST J100656.4+345445	2.13e–12	10 06 56.33 +34 54 43.8	6.6	1.00	19.0			BLLAC
J100750.6–090437		3.24e–12	10 07 48.79 –09 04 49.8 ^O	10.7 ^E	0.12	(17.6)			
	IRAS F10053–0850		10 07 50.44 –09 04 44.3 ^O	10.7 ^E	0.86	(15.6)	0.0668 ¹		Sy1.5
J100811.5+470526	RX J1008.1+4705	1.08e–11	10 08 11.72 +47 05 26.2	4.7	1.00	19.0	0.3430		BLLAC
J100848.8–095442	NPM1G –09.0361	6.65e–12	10 08 49.34 –09 54 57.8	2.7	0.99	15.9	0.0575 ¹		Sy1
J101001.9+300334	[HB89] 1007+303	2.09e–12	10 10 00.79 +30 03 21.0 ^R	2.9	1.00	17.0	0.2600		QSO
J101015.9–311909		2.90e–11	10 10 15.87 –31 19 06.5	74.3	1.00	(16.4)			
J101027.9+413242	[HB89] 1007+417	2.97e–12	10 10 27.63 +41 32 36.6	499.0 ^E	1.00	16.5	0.6123		QSO
J101244.4+422959	B3 1009+427	7.30e–12	10 12 44.22 +42 29 56.7	79.5	1.00	16.5			BLLAC
J101330.4–283128	[HB89] 1011–282	3.64e–12	10 13 29.43 –28 31 09.0 ^O	833.3 ^D	0.99	16.9	0.2530		QSO
J101335.9–135108 [†]	PMN J1013–1351	3.78e–12	10 13 35.07 –13 51 30.6	257.6	1.00	(17.8)	0.1517 ¹		Early
J101504.3+492604	[HB89] 1011+496	1.81e–11	10 15 04.13 +49 26 01.1	378.1	1.00	16.1	0.2000		BLLAC
J101556.7–200219	CTS 0090	5.14e–12	10 15 56.47 –20 02 28.1	13.6	0.98	15.7			Sy1
J101616.4+410817	FIRST J101616.8+410812	5.50e–12	10 16 16.81 +41 08 09.4	14.8	1.00	(18.2)	0.2800 ¹		BLLAC
J101655.5+732359	NGC 3147	4.77e–12	10 16 53.86 +73 24 03.5	92.5 ^E	1.00	11.4	0.0094	SA(rs)bc	Sy2
J101718.0+291439	IRAS F10144+2929	7.41e–12	10 17 18.72 +29 14 27.5	2.5	1.00	15.8	0.0450 ¹		Sy2
J101900.8+375250	FIRST J101900.4+375240	9.35e–12	10 19 00.73 +37 52 48.0	4.0	1.00	17.1	0.1350 ²		Sy1
J101912.1+635802 [†]		5.51e–12	10 19 12.51 +63 58 13.9 ^O	4.1	0.06	(19.8)			

Table 2—Continued

(1) ROSAT BSC 1RXS name	(2) NED Common name	(3) S _x ergs s ⁻¹ cm ⁻²	(4) Position α (J2000) δ	(5) S _{1.4GHz} mJy	(6) ID P	(7) m _B	(8) Redshift	(9) NED Morphology	(10) Sp. Class.
J102212.5+512405	MRK 0141		10 19 12.60 +63 58 02.5 ^O	4.1	0.91	15.4	0.0417	E	Sy1.5
J102228.9+500630 [†]	MS 1019.0+5139	5.11e-12	10 22 12.82 +51 23 56.3	5.1	1.00	18.1	0.1410		BLLAC
J102232.2-103740	ABELL 0980:001	4.34e-12	10 22 28.50 +50 06 20.7	27.9	1.00	17.5	0.1582		
J102244.2-011257	[HB89] 1020-103	6.46e-12	10 22 32.72 -10 37 43.0	942.8	1.00	16.1	0.1970		BLLAC/HPQ
J102339.3+041117 [†]	RX J1022.7-0112	1.17e-11	10 22 43.77 -01 13 01.7	36.2	1.00	17.2	— ¹		BLLAC
J102350.7-271522 [†]	RX J1023.7+0411:003	1.02e-11	10 23 39.64 +04 11 15.4	7.4	1.00	(18.0)	0.2906		
	ABELL 3444:001	1.03e-11	10 23 50.23 -27 15 24.0 ^O	10.3	0.75	17.5	0.2542		
			10 23 50.50 -27 15 22.7 ^O	10.3	0.24	(18.9)			
J102658.5-174905		5.70e-12	10 26 58.52 -17 48 58.5	11.5	1.00	(16.6)	0.1142 ¹		BLLAC
J102730.9+352614	FIRST J102732.4+352621	1.94e-12	10 27 32.52 +35 26 20.2	11.9	1.00	18.7			
J102758.9-064804 [†]	LCRS B102528.0-063237	4.09e-12	10 27 58.77 -06 47 53.5	14.8	0.99	15.3	0.1165		
J102906.4+272918		3.11e-12	10 29 01.64 +27 28 51.6 ^O	4.6 ^E	0.50	(16.5)	0.0384 ¹		Sy1
J103024.5+551630	87GB 102712.9+553156	1.70e-12	10 30 25.74 +55 16 27.5	156.8	1.00	(19.3)			QSO
J103058.8+310305	[HB89] 1028+313	7.04e-12	10 30 59.12 +31 02 57.9	231.3 ^E	1.00	16.7	0.1782	Opt.var.	Sy1
J103118.6+505341	87GB 102814.4+510859	5.46e-11	10 31 18.51 +50 53 36.9	37.9	1.00	17.0	0.3610		BLLAC
J103137.7-260725		3.91e-12	10 31 37.98 -26 07 15.8	20.6	1.00	(18.3)			
J103155.8-141659 [†]	EC 10294-1401	4.93e-11	10 31 54.41 -14 16 50.9	14.6	1.00	13.9	0.0860	E1	Sy1
J103156.6-184628	IRAS 10295-1831	4.36e-12	10 31 57.21 -18 46 33.5	10.3	1.00	15.6	0.0403		Sy1
J103214.3+401607	FIRST J103213.9+401616	1.54e-12	10 32 13.97 +40 16 16.4 ^F	5.8	1.00	(15.7)	0.0776	cD	
J103239.1+212147	IRAS 10298+2137	1.94e-12	10 32 38.62 +21 21 48.3 ^F	4.2	1.00	(17.3)	0.1114 ¹		Sy2/HII
J103256.6-270727		5.09e-12	10 32 57.09 -27 07 21.0 ^O	7.6	0.13	(19.6)			
	IRAS F10306-2651		10 32 57.03 -27 07 30.7 ^O	7.6	0.85	16.2		SB	Sy1
J103423.1+734525	NGC 3252	4.13e-12	10 34 22.74 +73 45 53.1 ^O	4.4	0.87	14.1	0.0039	SBd? sp	
J103438.7+393834	KUG 1031+398	3.53e-11	10 34 38.57 +39 38 27.8	25.9	1.00	15.6	0.0424	Compact	Sy1
J103459.5+304138 [†]	LEDA 093943	3.58e-12	10 35 00.25 +30 41 37.5	9.1	1.00	17.1	0.1373		
J103744.9+571159	87GB 103431.3+572750	1.91e-12	10 37 44.28 +57 11 56.4	71.7	1.00	16.7			BLLAC
J103846.7+533002	NGC 3310	1.39e-12	10 38 45.81 +53 30 11.0	364.2 ^E	1.00	11.1	0.0033	SAB(r)bc pec	HII
J103944.0-084059	MCG -01-27-030	5.31e-12	10 39 43.26 -08 41 07.5	3.1	0.94	15.4	0.0649	cD;S0p	
J104043.7+395706 [†]		6.96e-12	10 40 43.77 +39 57 08.6	23.9	1.00	(17.6)			
J104148.9+390133	B3 1038+392	9.71E-13	10 41 49.03 +39 01 22.3	33.9	1.00	15.8	0.2082 ¹		Early
J104303.0+005423	[HB89] 1040+011 NED01	6.28e-12	10 43 03.87 +00 54 19.4	33.2	1.00	19.2	0.7300		QSO
J104427.6+271813		3.74e-12	10 44 28.41 +27 18 12.7	3.6 ^E	1.00	(16.2)	0.0760 ²		Sy1
J104431.7-070404 [†]		8.48e-12	10 44 32.78 -07 04 07.6	32.1	1.00	17.6	0.1324		
J104439.4+384541	CGCG 212-045	4.77e-12	10 44 39.13 +38 45 35.3 ^O	3.3	0.81	14.9	0.0357 ¹		Sy1.5
J104651.9-253546 [†]	RX J1046.8-2535:001	4.74e-12	10 46 51.44 -25 35 47.2	14.1	1.00	(18.4)	0.2426		
J104956.7+351644		1.68e-12	10 49 56.59 +35 16 41.0	4.4	1.00	(18.8)			
J105025.7-125045	NGC 3411	1.65e-11	10 50 26.07 -12 50 39.8	33.0 ^E	1.00	13.1	0.0153	E+	
J105034.8-023611	LCRS B104803.7-022022	3.94e-12	10 50 36.52 -02 36 18.4	14.5	1.00	16.8	0.1659 ¹		Early
J105037.1+801204	IRAS F10460+8027	4.03e-12	10 50 35.90 +80 11 55.3	5.7	1.00	(16.1)	0.1183 ¹		Sy1
J105125.1+394329		4.05e-12	10 51 25.40 +39 43 25.6	10.8	1.00	(18.7)			
J105130.2-091754		5.57e-12	10 51 28.25 -09 17 52.7 ^{OR}	964.2	0.08	(19.9)			
	PG 1048-090		10 51 29.93 -09 18 10.3 ^{OR}	2034.3 ^D	0.88	16.8	0.3440	Opt.var.	QSO

Table 2—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position		S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	(J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J105232.8+612523	[HB89] 1049+616	3.35e-12	10 52 32.72	+61 25 20.8	988.9	1.00	16.5	0.4220	Opt.var.	QSO
J105344.2+492956 [†]	87GB 105048.1+494610	1.27e-12	10 53 44.02	+49 29 55.2	64.6	0.99	17.4			
J105431.0+694933	RX J1054.5+6949	1.60e-12	10 54 31.63	+69 49 18.3	4.2	1.00	17.0	0.0920		
J105607.0+025215	RBS 0921	1.48e-11	10 56 06.26	+02 52 27.7	4.3 ^E	1.00	18.3	0.2363 ¹		Early
J105723.5+230317		5.28e-12	10 57 23.01	+23 03 15.2	7.9	1.00	(18.8)			
J105750.7-275354	RX J1057.8-2753	4.55e-12	10 57 50.80	-27 54 11.2	63.8	1.00	17.8	0.0920		BLLAC
J105755.9+191527		1.64e-12	10 57 55.77	+19 15 44.9 ^O	3.1	0.42	(17.5)			
	IRAS F10552+1931		10 57 55.73	+19 15 24.5 ^O	3.1	0.53	15.9	0.0772 ¹		Sy2
J105801.2+202937	MRK 0634	2.49e-12	10 58 01.15	+20 29 13.7 ^F	2.8	1.00	15.9	0.0662		Sy1
J105818.3+195149	[HB89] 1055+201	2.06e-12	10 58 17.90	+19 51 51.0 ^F	350.6 ^E	1.00	17.1	1.1100	Opt.var.	QSO
J105825.9+564716 [†]	ABELL 1132:[F ⁺ TC95] 008	4.51e-12	10 58 23.69	+56 47 36.6	40.4 ^E	0.99	15.6			
J105837.5+562816	87GB 105536.5+564424	3.83e-12	10 58 37.71	+56 28 11.8	228.5	1.00	15.8	0.1440		BLLAC
J110021.3+401933	FIRST J110021.0+401927	7.37e-12	11 00 21.02	+40 19 28.8	18.3	1.00	20.2	— ²		BLLAC
J110312.1+414200	CGCG 213-021	1.43e-12	11 03 11.38	+41 42 17.7	8.4	1.00	15.1	0.0309 ¹		Early
J110331.5-325115	[HB89] 1101-325	5.32e-12	11 03 31.50	-32 51 18.8	899.3 ^D	1.00	16.3	0.3554	Opt.var.	Sy1
J110337.7-232931	PMN J1103-2329	5.41e-11	11 03 37.51	-23 29 24.8	121.0	1.00	16.5	0.1860		BLLAC
J110412.4+765859	PG 1100+772	8.58e-12	11 04 12.22	+76 58 59.7	2340.6	1.00	15.7	0.3115	Opt.var.	QSO
J110427.1+381231	MRK 0421	3.29e-10	11 04 27.42	+38 12 32.9	899.6 ^T	1.00	13.3	0.0300	S?	Blazar
J110436.4+212431	87GB 110153.6+214051	1.48e-12	11 04 36.19	+21 24 18.5	66.0	1.00	16.4	0.1876 ¹		Sy1
J110643.5+723407	NGC 3516	3.31e-12	11 06 47.71	+72 34 10.3	32.4	1.00	12.5	0.0088	(R)SB(s)0 ⁰ :	Sy1.5
J110715.1+162810	[HB89] 1104+167	4.69e-12	11 07 15.02	+16 28 01.5	869.5 ^D	1.00	15.7	0.6320	Opt.var.	Sy1
J110748.2+150217	87GB 110510.0+151723	5.94e-12	11 07 48.14	+15 02 10.8	43.5	1.00	18.0	0.2594 ¹		BLLAC
J110858.0-014924		3.50e-12	11 08 58.22	-01 49 32.0	81.6	1.00	18.0	0.1056 ¹		Early
J110943.6+214519		2.17e-12	11 09 42.25	+21 44 31.6 ^O	5.6	0.28	(18.5)			
	[BBN91] 110702+220045		11 09 41.15	+21 44 25.3 ^O	5.6	0.40	(15.7)	0.0317		
J111047.1-283001	ESO 438- G 009	7.84e-12	11 10 48.03	-28 30 03.2	15.5	1.00	14.2	0.0245	(R' ₁)SB(rl)ab	Sy1.5
J111131.2+345212	FIRST J111130.8+345203	7.28e-12	11 11 30.97	+34 52 00.7	8.4	1.00	19.5	0.2120		QSO
J111137.2+405031 [†]	MCG +07-23-030	4.37e-12	11 11 39.03	+40 50 22.3 ^F	820.2 ^E	0.94	17.0	0.0737		
J111224.2+175131		2.37e-12	11 12 24.65	+17 51 21.9	14.3	1.00	(18.6)			
J111349.5+093518	IC 2637	4.77e-12	11 13 49.71	+09 35 10.5	44.0	1.00	14.0	0.0292	E+ pec	Sy1.5
J111357.7+144217	[HB89] 1111+149	1.48e-12	11 13 58.69	+14 42 27.5	642.3	1.00	18.0	0.8690	Opt.var.	QSO
J111422.6+582318 [†]	8C 1111+586	2.22e-12	11 14 21.65	+58 23 19.4 ^R	66.6 ^E	1.00	14.0	0.2060	cD	
J111450.1-121351 [†]		3.91e-12	11 14 49.80	-12 13 49.1 ^O	9.0	0.36	(18.8)			
	LCRS B111219.3-115728		11 14 50.28	-12 13 50.4 ^O	9.0	0.63	16.4	0.1905 ¹		Early
J111553.3+012947		4.37e-12	11 15 51.81	+01 29 55.5	17.2	1.00	20.1			
J111706.3+201410	87GB 111429.0+203022	5.62e-11	11 17 06.25	+20 14 07.9	103.1	1.00	15.2			BLLAC
J111717.3+000631	RX J1117.2+0006	3.53e-12	11 17 17.49	+00 06 33.5	14.2	1.00	(19.1)			QSO
J111803.6+450657	RX J1118.0+4506	3.70e-12	11 18 03.21	+45 06 44.8	16.0	1.00	16.5	0.1060		Sy1
J111908.1+211915	PG 1116+215	1.20e-11	11 19 08.57	+21 19 13.9	6.5	1.00	15.2	0.1765	E2	Sy1?
J111930.9+222647	MG2 J111929+2226	2.75e-12	11 19 30.28	+22 26 49.4	97.2	1.00	16.7	0.4200 ²		Sy1
J111941.0-304652		3.91e-12	11 19 39.60	-30 47 20.9 ^O	9.4	0.75	(19.3)			
J112016.7+125917	MESSIER 066	2.28e-12	11 20 15.98	+12 59 02.7 ^R	459.5 ^E	1.00	9.6	0.0024	SAB(s)b	Sy

Table 2—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position		S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	(J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J112047.5+421217	87GB 111803.1+422817	1.36e−11	11 20 48.06	+42 12 14.3	24.1	1.00	17.3	0.1240		BLLAC
J112109.9+535125	RX J1121.1+5351	2.74e−12	11 21 07.99	+53 51 22.9	2.6	1.00	16.1	0.1027 ¹		Sy1
J112147.3+114420	PG 1119+120	8.07e−12	11 21 47.21	+11 44 18.4	3.5	0.99	15.1	0.0502	Compact	Sy1
J112318.0−323219		4.48e−12	11 23 18.12	−32 32 16.6	20.3	1.00	(19.0)			
J112349.2+723002		6.60e−12	11 23 49.18	+72 30 00.3	12.5	1.00	(18.6)			
J112357.4+212914		3.57e−12	11 23 56.09	+21 28 58.5	24.0	1.00	(19.3)			
J112407.9+061256	CGCG 039−167	1.06e−11	11 24 08.74	+06 12 54.0	15.1	1.00	14.9	0.0372 ¹		Sy1
J112453.0+493408	RGB J1124+495	2.65e−12	11 24 53.76	+49 34 11.3	21.0	1.00	18.0			
J112551.6−074219		5.12e−12	11 25 51.96	−07 42 21.4	39.3	1.00	19.0	0.2786 ¹		BLLAC?
J112612.5−203724		4.95e−12	11 26 11.94	−20 37 24.0	22.7	1.00	(18.7)			
J112704.5−185731	PKS 1124−186	1.16e−11	11 27 04.41	−18 57 18.1	536.6	1.00	19.0	1.0500		QSO
J112716.6+190914		1.53e−12	11 27 16.24	+19 09 20.2	7.0	1.00	(16.1)	0.1055 ¹		Sy2
J112831.5+583343	NGC 3690	1.19e−12	11 28 32.61	+58 33 46.8	678.1	1.00	11.8	0.0105	SBm? pec	
J112914.7−383724		5.73e−12	11 29 14.76	−38 37 32.3	5.5	0.93	(16.2)			
J113007.5−144939	[HB89] 1127−145	3.04e−12	11 30 07.05	−14 49 27.2	5622.8	1.00	16.9	1.1870		Blazar/LPQ
J113046.7−313801		3.64e−12	11 30 46.14	−31 38 07.4 ^O	26.5	0.56	(17.0)			
			11 30 46.68	−31 38 05.6 ^O	26.5	0.44	(20.1)			
J113104.6−094353		2.25e−12	11 31 05.29	−09 44 10.7	21.9	1.00	(16.5)			
J113108.9+311409	[HB89] 1128+315	9.62e−12	11 31 09.66	+31 14 04.2	370.4	1.00	16.6	0.2890		QSO
J113109.0+263212	RX J1131.1+2632	1.81e−12	11 31 09.21	+26 32 08.4 ^F	3.0 ^E	1.00	17.0	0.2440		Sy1
J113121.4+333447 [†]	ABELL 1290:001	1.87e−12	11 31 20.93	+33 34 45.7	6.4	1.00	17.5	0.2203 ¹		Early
J113153.7−195543 [†]		4.64e−12	11 31 53.80	−19 55 33.0 ^{OR}	14.0 ^E	0.08	(19.8)			
	ABELL 1300:[POD97] 219		11 31 54.19	−19 55 40.2 ^{OR}	14.0 ^E	0.86	(18.4)	0.3077		
	ABELL 1300:[POD97] 121		11 31 54.80	−19 55 41.9 ^{OR}	14.0 ^E	0.06	17.0	0.3077		
J113155.4−123155		3.43e−12	11 31 51.51	−12 31 58.6 ^{OR}	16.3 ^E	0.11	(16.6)			
			11 31 53.36	−12 31 59.9 ^{OR}	4.3 ^E	0.86	(18.6)	0.6537 ¹		QSO
J113222.4+555828 [†]	RX J1132.3+5558:001	2.32e−12	11 32 22.86	+55 58 20.5	53.4 ^E	1.00	(16.8)	0.0493 ¹	cD?	Early
J113302.0+184655		1.67e−12	11 33 02.67	+18 47 07.3	3.7	1.00	(21.5)			
J113443.6−172853		3.45e−12	11 34 44.54	−17 29 03.9	4.8	1.00	(19.2)			
J113448.4+490438 [†]	IC 0712	3.70e−12	11 34 49.98	+49 04 43.5	32.8	1.00	14.7	0.0331 ¹		Early
J113626.6+700932	[HB89] 1133+704	5.77e−11	11 36 26.48	+70 09 25.8	328.4	1.00	15.5	0.0460	Compact	BLLAC
J113629.4+213552	NGC 3758	7.42e−12	11 36 29.17	+21 35 49.1	11.2	1.00	15.2	0.0299	Sab pec	Sy1
J113630.9+673708	87GB 113338.9+675347	2.45e−11	11 36 29.92	+67 37 06.0	45.8	1.00	17.3	0.1350		BLLAC
J113634.9−222928	IRAS F11340−2213	5.56e−12	11 36 35.86	−22 29 39.0	3.5	0.99	16.1			Sy1
J113737.4+103931		2.63e−12	11 37 37.93	+10 39 28.5	3.0	1.00	(16.6)	0.1745 ¹		Sy1
J113755.4−171034		4.41e−12	11 37 55.12	−17 10 31.0	5.1	1.00	(19.0)			
J113901.6−374420	NGC 3783	5.48e−11	11 39 01.66	−37 44 21.0	44.6	1.00	12.6	0.0097	(R')SB(r)a	Sy1
J113910.4−135027	[HB89] 1136−135	3.39e−12	11 39 10.69	−13 50 42.9	4225.3	1.00	16.1	0.5600		Sy1
J113957.9+654753	[HB89] 1137+660	3.20e−12	11 39 58.60	+65 47 47.1	3291.9 ^D	1.00	16.3	0.6460	Opt.var.	LPQ
J114023.5+152814		8.37e−12	11 40 23.55	+15 28 08.3	70.0	1.00	(17.4)	0.2447 ¹		Early?
J114106.1+024110	RX J1141.1+0241	2.12e−12	11 41 05.71	+02 41 17.1 ^O	2.5	0.90	16.9	0.0928 ¹		Sy1
J114116.2+215624	PG 1138+222	2.62e−11	11 41 15.87	+21 56 30.3	4.9 ^E	1.00	14.9	0.0632		Sy1

Table 2—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position		S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	(J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J114124.2−121632 [†]	LCRS B113851.7−115959	4.80e−12	11 41 24.26	−12 16 37.7	158.1	1.00	15.2	0.1195		
J114305.9+184348	87GB 114030.0+190057	2.26e−12	11 43 06.01	+18 43 43.2	57.0	1.00	16.4	0.3739 ¹		QSO
J114333.1+462343		6.66e−12	11 43 38.42	+46 24 03.2 ^F	13.5	1.00	(16.6)			
J114429.9+365314	KUG 1141+371	2.04e−11	11 44 30.10	+36 53 00.2	2.9	1.00	16.5	0.0400	Spiral	Sy1
J114442.2+672435 [†]	87GB 114155.4+674039	5.02e−12	11 44 36.82	+67 24 21.0 ^O	236.6 ^A	—	12.8	0.1159		
J114452.7+194706		7.63e−12	11 44 53.75	+19 46 32.1 ^O	2.5 ^A	0.14	(18.5)			
	CGCG 097−125		11 44 54.84	+19 46 35.2 ^O	2.5 ^A	0.50	15.7	0.0276	S0a	
J114509.3+304724	[HB89] 1142+310	6.97e−12	11 45 10.06	+30 47 13.4	3.3	1.00	16.0	0.0596		Sy1.5
J114517.3−342552	PKS 1142−341	9.64e−12	11 45 19.61	−34 25 59.0	369.3 ^S	0.97	16.0		dumbbell sys.	
J114518.3+152954	MCG +03−30−100	3.20e−12	11 45 22.03	+15 29 41.4	350.9	0.99	17.0	0.0679		
J114535.8−033948	LCRS B114301.7−032322	7.32e−12	11 45 35.09	−03 39 59.1	19.6	1.00	17.2	0.1675		
J114540.5−182716	IRAS F11431−1810	7.58e−11	11 45 40.36	−18 27 13.7	9.7	1.00	14.7	0.0330	Irr	Sy1
J114708.2−322954		4.43e−12	11 47 09.01	−32 30 02.9 ^O	3.7	0.51	(19.2)			
			11 47 09.51	−32 30 10.8 ^O	3.7	0.06	(20.5)			
			11 47 09.12	−32 30 06.4 ^O	3.7	0.40	(21.5)			
J114754.9+220548		3.08e−12	11 47 54.88	+22 05 34.3	4.1 ^E	1.00	(18.8)			
J114856.3−040408	[HB89] 1146−037	1.98e−12	11 48 55.82	−04 04 11.4 ^R	617.8 ^T	1.00	16.9	0.3410		Sy1/LPQ
J114930.4+243928	RGB J1149+246	9.62e−12	11 49 30.46	+24 39 22.6	28.5	1.00	18.0			BLLAC
J114947.0−121845 [†]		3.81e−12	11 49 48.89	−12 19 39.7 ^O	92.2 ^E	0.05	(19.6)			
	LCRS B114716.2−120301		11 49 49.20	−12 19 42.7 ^O	92.2 ^E	0.87	16.1			
J115035.2+415446	FIRST J115034.7+415440	3.69e−12	11 50 34.65	+41 54 42.0	26.3	1.00	17.0	1.0100 ²		QSO
J115125.1+585932	87GB 114846.6+591612	1.40e−12	11 51 24.66	+58 59 18.6	184.8	1.00	17.3			BLLAC
J115203.9−112223	PG 1149−110	1.41e−11	11 52 04.42	−11 22 18.5	10.9 ^E	0.99	15.5	0.0490		Sy1
J115227.1+320957	87GB 114949.8+322620	3.34e−12	11 52 27.54	+32 09 59.4	40.9 ^E	1.00	18.1	0.3740		Sy1
J115228.0−374837		3.54e−12	11 52 27.67	−37 48 34.9	2.5	0.98	(17.1)			
J115238.2−051229	MCG −01−30−041	1.93e−12	11 52 38.07	−05 12 29.5	12.4	1.00	14.7	0.0192	(R)SB(rs)ab	Sy1.8
J115323.4+583159	87GB 115044.1+584817	1.53e−12	11 53 23.95	+58 31 38.6 ^R	106.3 ^T	1.00	17.3	0.2026 ¹		Sy1.5
J115324.4+493108	[HB89] 1150+497	5.90e−12	11 53 24.51	+49 31 09.5	1572.7	1.00	17.1	0.3340	Opt.var.	HPQ
J115404.9−001008		4.04e−12	11 54 04.30	−00 10 08.7	10.6	1.00	(18.8)	— ¹		BLLAC?
J115518.9+232431 [†]	[HMS56] 1153+2341 01A	9.76e−12	11 55 17.86	+23 24 27.2 ^O	4.0	0.44	(19.3)	0.1428		
	MCG +04−28−097		11 55 17.99	+23 24 17.8 ^O	4.0	0.53	18.0	0.1429		
J115658.6+241523	RX J1157.0+2415:[AEF92]	2.63e−12	11 56 55.72	+24 15 35.4 ^F	5.4 ^E	0.99	(16.0)	0.1419		
J115719.0+333645 [†]	FIRST J115716.6+333653	2.10E−12	11 57 17.16	+33 36 54.7 ^O	34.3 ^E	0.19	(19.8)			
	7C 1154+3353		11 57 17.39	+33 36 39.9 ^O	34.3 ^E	0.69	17.0	0.2130	cD	
J115756.1+552717	NGC 3998	7.37e−12	11 57 56.14	+55 27 12.6	101.4	1.00	11.6	0.0035	SA(r)0 ⁰ ?;Sy1	LINER
J115941.4−195935	IRAS F11571−1942	6.93e−12	11 59 40.93	−19 59 22.3	10.0	1.00	16.2	0.4500		QSO
J120013.5−004632		2.68e−12	12 00 14.18	−00 46 38.6	40.2 ^E	1.00	(17.5)	0.1791 ¹		Sy1
J120105.6−000658		1.71e−12	12 01 06.24	−00 07 02.6	70.6	1.00	18.7	0.1651 ¹		Early
J120114.3−034040	MRK 1310	1.52e−11	12 01 14.41	−03 40 34.1	3.9	1.00	14.5	0.0194	E	Sy1
J120119.2−371547	IRAS F11587−3659	4.23e−12	12 01 18.29	−37 16 06.3	8.8 ^E	0.90	(14.8)			
J120226.9−012908	IRAS 11598−0112	2.44e−12	12 02 26.71	−01 29 17.0	14.9	1.00	17.9	0.1510 ¹		Sy1.5
J120234.1−052801	[HB89] 1200−051	5.56e−12	12 02 34.20	−05 28 02.5	530.5	1.00	18.0	0.3810		QSO

Table 2—Continued

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position	S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α (J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J120240.6+263143	MG2 J120242+2631	1.65e-12	12 02 40.30 +26 31 43.6	91.8 ^E	1.00	17.4	0.4800 ²		Sy1
J120246.0-034710	LCRS B120011.5-033039	1.90e-12	12 02 45.48 -03 47 21.7	3.1	1.00	16.4	0.0645		Sy1
J120308.9+443155	NGC 4051	4.92e-11	12 03 09.28 +44 31 54.0	98.1 ^E	1.00	10.8	0.0024	SAB(rs)bc	Sy1.5
J120329.8+480316	87GB 120056.9+481950	1.77e-12	12 03 29.92 +48 03 13.7	63.4	1.00	16.4	0.8133 ¹		QSO
J120413.0+114549		1.84e-12	12 04 12.17 +11 45 55.4	15.1	1.00	(16.9)	— ¹		
J120511.7+392043 [†]	RX J1205.2+3920:[CEF95]	7.47e-12	12 05 09.83 +39 20 31.8	3.2	0.99	(16.3)	0.0370	cD	
J120613.0-084743	MRC 1203-085	2.94e-12	12 06 12.11 -08 48 02.5	162.3 ^E	1.00	(21.5)			
J120711.1-174551	RBS 1068	6.07e-12	12 07 11.51 -17 46 05.5 ^{OR}	4.3	0.90	20.5			QSO
			12 07 12.03 -17 45 55.8 ^{OR}	4.3	0.10	16.7	0.0000 ¹		
J120946.0+321702	FIRST J120945.2+321700	2.84e-12	12 09 45.24 +32 17 00.9 ^{OF}	2.3	1.00	17.7	0.1450		Sy1
J120954.9+062806	IRAS F12073+0644	2.03e-12	12 09 54.37 +06 28 10.5	4.0	1.00	16.8	0.0796 ¹		Sy1
J121022.8+392926	[HB89] 1207+397	4.52e-12	12 10 26.65 +39 29 09.5	19.0	1.00	20.3	0.6150	Opt.var.	BLLAC
J121031.3+392450	NGC 4151	3.67e-12	12 10 32.52 +39 24 20.9	360.1	0.99	11.5	0.0033	(R')SAB(rs)ab:	Sy1.5
J121038.9+315704	[HB89] 1208+322	1.37e-12	12 10 37.56 +31 57 06.2 ^{OR}	370.4 ^D	1.00	16.0	0.3880		QSO
J121105.4+352005 [†]	87GB 120833.7+353632	1.46e-12	12 11 08.28 +35 19 59.1	74.9	1.00	(17.4)	0.1400 ¹		Early
J121158.1+224236		7.11e-12	12 11 58.68 +22 42 32.4	20.2	1.00	(18.7)			
J121301.8+512942		1.69e-12	12 13 00.87 +51 29 35.9	36.2	1.00	(18.4)			
J121321.6-261802	RBS 1080	1.06e-11	12 13 23.13 -26 18 10.1	7.3	1.00	19.0	0.2780		BLLAC
J121345.2+363754	NGC 4190	1.90e-12	12 13 44.75 +36 37 58.8	7.4 ^E	1.00	13.9	0.0008	Im pec	
J121500.4+050051		3.51e-12	12 15 00.71 +05 00 50.6	17.2	1.00	(16.4)			
J121506.7+331129	NGC 4203	6.05e-12	12 15 05.23 +33 11 52.7	6.6	1.00	11.8	0.0036	SAB0 ⁻ :	LINER
J121510.9+073205	PMN J1215+0732	4.79e-12	12 15 10.88 +07 32 05.2	137.7	1.00	16.0	0.1300		BLLAC
J121603.6-024302		2.20e-12	12 16 03.25 -02 43 04.5 ^F	11.0	1.00	(18.0)	0.1690? ¹		BLLAC
J121606.4+092915	MG1 J121604+0929	1.57e-12	12 16 06.34 +09 29 09.5	213.5 ^T	1.00	13.2	0.0935 ¹		Early
J121710.0+071127	NGC 4235	1.39e-12	12 17 09.62 +07 11 29.3	15.5 ^E	1.00	12.6	0.0080	SA(s)a	Sy1
J121721.0+305632	FBQS J1217+3056	1.42E-12	12 17 21.18 +30 56 30.6	7.8	0.99	17.0	0.3074		QSO
J121742.6+033943	PGC 039445	2.11E-11	12 17 39.76 +03 39 44.5 ^F	1117.4 ^E	0.84	14.9	0.0759	E	
J121752.1+300705	[HB89] 1215+303	4.01e-11	12 17 52.11 +30 07 00.0	572.7	1.00	15.5	0.1300	Opt.var.	BLLAC
J121827.0+294853	NGC 4253	6.75e-11	12 18 26.43 +29 48 47.3	39.7	1.00	13.7	0.0129	(R')SB(s)a:	Sy1.5
J121900.4+471747	MESSIER 106	2.26e-12	12 18 57.53 +47 18 14.1 ^{RF}	508.4 ^E	0.91	9.1	0.0015	SAB(s)bc	Sy1.9/LINER
J121946.0-031419		3.28e-12	12 19 45.70 -03 14 21.4	28.9	1.00	(17.6)	— ¹		BLLAC
J122011.9+020342	[HB89] 1217+023	4.77e-12	12 20 12.31 +02 03 41.7	673.7 ^E	1.00	16.5	0.2400	Opt.var.	QSO
J122044.5+690533		4.97e-12	12 20 44.21 +69 05 22.6	8.0	0.99	(19.4)			
J122106.9+474235	RX J1221.1+4742	1.45e-12	12 21 07.79 +47 42 29.7	42.4	1.00	19.0	0.2099 ¹		Early
J122121.7+301041	PG 1218+304	2.27e-11	12 21 21.92 +30 10 36.4	71.5	1.00	16.5	0.1820	Opt.var.	BLLAC
J122131.1+281402	W COMAE	2.23e-12	12 21 31.61 +28 13 58.3	732.1	1.00	16.5	0.1020	Opt.var.	BLLAC
J122131.4+082139	MG1 J122131+0821	1.75e-12	12 21 32.08 +08 21 43.8	176.9	1.00	17.1	0.1320 ¹		Early
J122144.4+751848	[HB89] 1219+755	1.74e-11	12 21 39.71 +75 18 34.1	7.7	0.95	14.5	0.0709	SB(r)ab pec	Sy1
J122208.7+030721		2.72e-12	12 22 08.58 +03 07 28.2	3.4	1.00	(21.5)			
J122222.8+041316	[HB89] 1219+044	1.45e-12	12 22 22.59 +04 13 17.3	800.8	1.00	18.0	0.9650		QSO
J122313.3+540906	4C +54.27	2.50e-12	12 23 13.23 +54 09 07.1	388.1	1.00	14.6	0.1570		BLRG
J122424.2+243618	87GB 122151.4+245224	2.54e-12	12 24 24.36 +24 36 23.7	25.9	1.00	17.6	0.2180		BLLAC

Table 2—Continued

(1) <i>ROSAT</i> BSC 1RXS name	(2) NED Common name	(3) S_x ergs s ⁻¹ cm ⁻²	(4) Position α (J2000) δ	(5) $S_{1.4\text{GHz}}$ mJy	(6) ID P	(7) m_B	(8) Redshift	(9) NED Morphology	(10) Sp. Class.
J122454.9+212252	PG 1222+216	3.29e-12	12 24 54.62 +21 22 47.2	2095.1	1.00	17.5	0.4350		Blazar
J122512.5+321354	87GB 122241.8+323117	2.26e-12	12 25 13.07 +32 14 01.2	49.8	1.00	13.0	0.0590 ²		Sy2
J122515.5+721450		1.37e-12	12 25 13.95 +72 14 50.2	5.2	0.98	(16.3)	0.1138 ¹		Early
J122539.9+245833	[HB89] 1223+252	2.11e-12	12 25 39.53 +24 58 35.9 ^O	365.4 ^D	1.00	16.0	0.2680	Opt.var.	QSO
J122623.9+324431	FIRST J122624.2+324429	4.76e-12	12 26 23.99 +32 44 32.8	2.8	1.00	17.2	0.2420 ²		Sy1
J122645.2+063906		1.48e-12	12 26 44.19 +06 38 54.1	17.2 ^E	1.00	(17.5)	— ¹		BLLAC
J122745.1+084147	RX J1227.7+0841	4.84e-12	12 27 44.64 +08 41 51.6	3.8	1.00	16.8	0.0852 ¹		Sy1
J122752.7+632317 [†]	87GB 122536.3+633947	2.12e-12	12 27 51.89 +63 23 02.2	233.6 ^E	1.00	15.5	0.1454		
J122906.5+020311	3C 273	1.05e-10	12 29 06.41 +02 03 05.1	54992.1	1.00	12.9	0.1583	Sy1	Blazar/LPQ
J122945.9+075927 [†]	MESSIER 049	1.39e-11	12 29 46.60 +08 00 02.9	221.3 ^E	1.00	9.4	0.0029	E2/S0(2)	
J123014.2+251805	MG2 J123013+2517	3.35e-12	12 30 14.09 +25 18 07.6	244.0	1.00	15.6	0.1350		BLLAC
J123132.5+641420	[HB89] 1229+645	4.55e-12	12 31 31.34 +64 14 19.4	58.8	1.00	16.9	0.1700		BLLAC
J123144.7+284754	MG2 J123143+2848	2.10e-12	12 31 43.85 +28 47 51.2	141.5	1.00	15.5	1.0300 ²		QSO
J123203.6+200930	PG 1229+204	1.03e-11	12 32 02.95 +20 09 26.4	3.1	1.00	15.3	0.0640	Compact	Sy1
J123416.9-385637		6.45e-12	12 34 17.15 -38 56 35.1 ^O	6.9	0.86	(18.3)			
			12 34 17.60 -38 56 44.6 ^O	6.9	0.14	(16.3)			
J123439.0+235014	NGP9 F378-0239966	2.22e-12	12 34 38.64 +23 50 13.0 ^F	15.3 ^E	1.00	19.3			
J123511.1-140332		2.52e-12	12 35 10.83 -14 03 23.7	4.0	1.00	(19.3)			
J123651.1+453907	CGCG 244-033	6.70e-12	12 36 51.09 +45 39 05.1 ^F	4.2	1.00	15.2	0.0302 ¹		Sy1.8/HII
J123651.2+250752	MG2 J123651+2507	1.45e-12	12 36 51.57 +25 07 50.7 ^F	270.4 ^D	1.00	17.6	0.5460		Sy1
J123658.8+631111 [†]	ABELL 1576:1234+6327	1.48e-12	12 36 57.46 +63 11 15.8	19.8	1.00	(21.5)	0.3019		
J123705.6+302002	FIRST J123705.5+302005	6.30e-12	12 37 05.58 +30 20 06.3	5.6	1.00	21.1	0.7000		BLLAC
J123739.2+625843	[HB89] 1235+632	3.64e-12	12 37 39.44 +62 58 43.7	12.5	1.00	18.5	0.2970	Opt.var.	BLLAC
J123743.6+114909	MESSIER 058	7.25e-12	12 37 43.74 +11 49 08.0	98.2 ^E	1.00	10.5	0.0051	SAB(rs)b	Sy1.9/LINER
J123802.1+361643	FIRST J123802.5+361639	1.93e-12	12 38 02.49 +36 16 39.8	4.8	1.00	18.3			QSO
J123808.3+532603	87GB 123550.3+534219	2.89e-12	12 38 07.86 +53 25 55.5 ^R	135.7 ^T	1.00	17.2	0.3478 ¹		Sy1
J123859.9+360917	FIRST J123900.6+360924	1.38e-12	12 39 00.68 +36 09 25.1	16.6	1.00	18.1			QSO
J123923.1+413245	FIRST J123922.7+413251	2.30e-12	12 39 22.50 +41 32 53.5	9.1	1.00	(19.5)			
J123939.6-052035	NGC 4593	6.00e-11	12 39 39.31 -05 20 35.9	4.8	1.00	11.7	0.0090	(R)SB(rs)b	Sy1
J123959.8-113725	MESSIER 104	2.70e-12	12 39 59.40 -11 37 23.5	94.0 ^E	1.00	9.0	0.0036	SA(s)a	Sy1.9/LINER
J124046.8-333408	ESO 381- G 007	1.38e-11	12 40 47.13 -33 34 18.1	3.1	0.97	15.9	0.0500	S0	Sy1.5
J124141.2+344032	FIRST J124141.3+344031	2.18e-12	12 41 41.27 +34 40 28.8	10.2	1.00	19.6			BLLAC
J124144.4+350353	NGC 4619	2.81e-12	12 41 44.18 +35 03 49.0	17.5	1.00	13.5	0.0231	SB(r)b pec?	Sy1
J124148.3+063602	RX J1241.8+0636	1.63e-12	12 41 48.24 +06 35 59.2	16.5	1.00	19.4	0.1500		BLLAC
J124149.8-145558	PMN J1241-1456	1.76e-11	12 41 49.41 -14 55 59.0	17.2	0.99	17.9			BLLAC
J124211.3+331703		1.07e-11	12 42 11.05 +33 17 12.3	6.4	1.00	(16.3)			
J124249.7+024117	NGC 4636	1.37e-11	12 42 50.33 +02 41 18.7	78.7 ^E	1.00	10.4	0.0037	E/S0 ₁	LINER
J124252.5+131530	NGC 4639	2.87e-12	12 42 52.32 +13 15 42.1	9.1 ^E	1.00	12.2	0.0034	SAB(rs)bc	Sy1.8
J124312.5+362743	TON 0116	1.64e-11	12 43 12.70 +36 27 45.1	147.9	1.00	17.1	— ²		BLLAC
J124340.6+113309	MESSIER 060	5.92e-12	12 43 40.13 +11 33 09.4	29.9 ^E	1.00	9.8	0.0047	E2	
J124349.3-153320 [†]	PMN J1243-1533	2.59e-12	12 43 48.87 -15 33 06.4	128.2 ^E	0.94	(18.1)			
J124701.3+442325	RGB J1247+443	3.21e-12	12 47 00.77 +44 23 20.2	10.6	1.00	15.3			

Table 2—Continued

(1) <i>ROSAT</i> BSC 1RXS name	(2) NED Common name	(3) S_x ergs s ⁻¹ cm ⁻²	(4) Position α (J2000) δ		(5) $S_{1.4\text{GHz}}$ mJy	(6) ID P	(7) m_B	(8) Redshift	(9) NED Morphology	(10) Sp. Class.
J124818.9+582031	PG 1246+586	5.39e-12	12 48 18.77	+58 20 28.8	245.3	1.00	14.0			BLLAC
J125015.0+315605	FIRST J125015.5+315559	1.82e-12	12 50 15.71	+31 56 04.1	5.6	1.00	19.3			
J125052.5+410713	MESSIER 094	3.29e-12	12 50 53.41	+41 07 08.6	269.9 ^E	1.00	9.0	0.0010	(R)SA(r)ab	LINER
J125135.2-295824	1ES 1248-296	4.29e-12	12 51 34.76	-29 58 43.2	10.4	1.00	19.0	0.4870		BLLAC
J125212.5-132450	NGC 4748	2.30e-11	12 52 12.65	-13 24 52.8	14.3	1.00	14.3	0.0146	Sa	Sy1
J125223.7+645143	87GB[BWE91] 1250+6508	1.69e-12	12 52 24.73	+64 51 38.4	43.2 ^S	1.00	16.6	0.3130		BLRG
J125228.7+563446	[HB89] 1250+568	2.58e-12	12 52 26.33	+56 34 19.7	2288.8	1.00	17.9	0.3210		Sy1
J125233.7-311605 [†]	RX J12525-3116:001	1.71e-11	12 52 34.93	-31 15 56.8	20.0	1.00	(13.9)	0.0535		
J125301.0+382629	FIRST J125300.9+382625	5.26e-12	12 53 00.58	+38 26 29.5	4.8	1.00	(19.0)			
J125303.0-091216	NGC 4761	2.51e-12	12 53 05.65	-09 12 21.1	5.4 ^E	0.95	13.5	0.0142	S0 ⁺	
J125341.2-393200	PMN J1253-3932	1.98e-11	12 53 41.32	-39 31 59.7	50.0	1.00	(17.2)			
J125347.3+032620	CGCG 043-056	2.86e-12	12 53 47.05	+03 26 30.0	107.0	1.00	15.7	0.0654 ¹		Early
J125422.2-290034 [†]	ESO 443- G 004	1.05e-11	12 54 22.34	-29 00 53.2	333.4 ^E	0.83	14.5	0.0548	S0	
J125425.0-282629		7.55e-12	12 54 25.38	-28 26 25.5 ^O	3.8	0.29	(18.7)			
	ESO 443-IG 005		12 54 24.81	-28 26 31.3 ^O	3.8	0.67	17.0	0.0686		Sy1
J125441.7-291342	ESO 443- G 007	1.29e-11	12 54 40.90	-29 13 43.5	1053.6 ^E	1.00	14.2	0.0574	S?	
J125445.9-152556		2.91e-12	12 54 45.00	-15 26 23.4	72.6	1.00	(18.4)			
J125456.3-265653		2.19e-11	12 54 56.26	-26 57 01.0	13.7	1.00	(15.2)			
J125609.5-080906	MCG -01-33-054	3.15e-12	12 56 10.66	-08 09 12.5	3.3 ^E	0.95	14.5	0.0129	Sdm	
J125611.2-054719	3C 279	3.48e-11	12 56 11.15	-05 47 20.1	9711.8	1.00	17.8	0.5362		Blazar/HPQ
J125710.0-172345 [†]	SGC 1254.5-1708	2.06e-11	12 57 11.59	-17 24 33.8 ^O	98.8	0.59	14.3	0.0475	cD;E+4	
J125710.6-133938	NGC 4825	3.15e-12	12 57 12.23	-13 39 55.1	13.2	1.00	12.6	0.0149	SA0 ⁻	
J125721.5+693015	ZwCl 1255.4+6946:001	1.52e-12	12 57 21.57	+69 30 34.8	3.5 ^E	0.95	(19.0)	0.2270	cD	
J125731.7+241245	RX J1257.5+2412	1.16e-11	12 57 31.95	+24 12 38.9	14.7	1.00	15.4	0.1410		BLLAC
J125748.3-235058		3.98e-12	12 57 47.98	-23 50 55.6 ^O	5.6	0.08	(19.1)			
	IRAS F12550-2334		12 57 48.02	-23 50 44.2 ^O	5.6	0.72	17.5	0.1195		Sy1
			12 57 48.79	-23 50 53.7 ^O	5.6	0.19	(18.6)			
J125847.7-044746		8.35e-12	12 58 47.85	-04 47 39.2	4.1	1.00	(18.2)			
J125921.5-041131 [†]		2.47e-11	12 59 22.21	-04 11 44.2	6.6	0.95	15.7	0.0853		
J130058.9-231213	PKS 1258-22	4.31e-12	13 00 58.50	-23 12 15.6	777.4	1.00	17.5	0.1300	Elliptical	
J130113.3-354315		4.24e-12	13 01 13.40	-35 43 12.0	60.2 ^T	1.00	(16.5)			
J130145.3+405628	87GB 125926.5+411221	1.41e-12	13 01 45.61	+40 56 25.6	29.0	1.00	17.5	- ²		BLLAC
J130250.3-023041 [†]		3.56e-12	13 02 52.56	-02 30 59.8	38.8	1.00	(16.1)			
J130255.6+505621	RX J1302.9+5056	6.00e-12	13 02 56.26	+50 56 18.3	2.8	1.00	20.2	0.6880		BLLAC
J130258.8+162423	MRK 0783	4.57e-12	13 02 58.93	+16 24 27.4	33.2	1.00	16.0	0.0672		Sy1.5
J130343.6-241506 [†]	ABELL 1664:[AEF92]	1.27e-11	13 03 42.65	-24 14 42.6	36.8	1.00	(15.5)	0.1276		
J130346.7+191635 [†]	IC 4130	8.39e-12	13 03 46.68	+19 16 17.3	78.7	1.00	14.0	0.0637	cD	
J130350.5-395035		4.96e-12	13 03 50.61	-39 50 43.9	13.3 ^E	0.96	(16.5)			
J130451.3+245446	RX J1304.8+2454	1.96e-12	13 04 52.03	+24 54 45.9	48.2 ^T	1.00	17.1	0.6000 ²		Sy1
J130510.0-033208	LCRS B130235.1-031605	3.10e-12	13 05 10.01	-03 32 09.6 ^O	5.3	0.54	16.4	0.0837 ¹		Sy1
J130533.2-103315	PG 1302-102	8.14e-12	13 05 33.00	-10 33 20.0	711.7	1.00	14.9	0.2784	E4(?);Opt.var.	LPQ
J130552.6+305405 [†]	MCG +05-31-128	2.47e-12	13 05 50.61	+30 54 15.4	75.7	1.00	(17.6)	0.1831		

Table 2—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position		S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	(J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J130730.1+092211	RX J1307.5+0922	1.76e-12	13 07 30.01	+09 22 05.2 ^{OR}	50.6 ^D	1.00	18.0			QSO
J130904.2-040609		2.94e-12	13 09 03.96	-04 06 11.9 ^O	110.2 ^D	1.00	(17.3)			
J130916.1-013658 [†]		2.36e-12	13 09 19.17	-01 37 20.9	11.4	0.98	16.2			
J131012.4-115746		2.30e-12	13 10 12.43	-11 57 46.9	83.3	1.00	(16.5)	0.1397 ¹		BLLAC
J131017.1-072712	MCG -01-34-008	2.58e-12	13 10 17.10	-07 27 16.8	8.6	0.96	14.5	0.0224	S	Sy2
J131106.4+003511	LBQS 1308+0051	1.61e-12	13 11 06.42	+00 35 09.8	17.7	1.00	18.6			
J131129.5-012017 [†]	MCG +00-34-016 NED1	1.37e-11	13 11 30.00	-01 20 28.3 ^F	10.9 ^E	1.00	18.8	0.1990		
J131156.0+085343	RX J1311.9+0853	4.69e-12	13 11 55.80	+08 53 42.7	5.3	1.00	(19.1)			
J131218.0+351524	PG 1309+355	1.50e-12	13 12 17.75	+35 15 21.5	47.7	1.00	15.4	0.1840	Sab	Sy1
J131250.7+554827	87GB 131050.3+560417	1.48e-12	13 12 53.20	+55 48 13.3	586.6	1.00	20.6			
J131327.2+363542	NGC 5033	3.44e-12	13 13 27.28	+36 35 40.4	122.8 ^E	1.00	10.8	0.0029	SA(s)c	Sy1.9
J131443.5+234823	MG2 J131441+2348	1.59e-12	13 14 43.80	+23 48 27.0	184.2 ^A	1.00	16.0	— ²		BLLAC
J131453.0-354313		3.54e-12	13 14 52.47	-35 43 22.0 ^O	4.1	0.87	(16.0)			
			13 14 53.02	-35 42 46.1 ^O	4.1	0.07	(20.1)			
J131506.8+514931 [†]	[CGH98] J1315+5148	2.09e-12	13 15 08.40	+51 48 50.5 ^F	47.1	0.74	18.0	0.2586		
J131523.8-162320	NGC 5044	5.23e-11	13 15 23.99	-16 23 07.6	36.1 ^E	1.00	11.8	0.0090	E0	
J131531.9+113327		2.91e-12	13 15 32.78	+11 33 31.7	20.6	1.00	(18.9)			
J131639.8+205516	NGP9 F379-1207883	1.59e-12	13 16 39.87	+20 55 12.6	12.8	1.00	19.1			
J131654.9+301454	FIRST J131654.5+301454	1.57e-12	13 16 55.28	+30 14 49.3	4.0	1.00	20.3			
J131713.6-382159		4.82e-12	13 17 12.24	-38 21 53.2	22.6	1.00	(17.5)			
J131750.4+601047	RX J1317.8+6010	2.65e-12	13 17 50.16	+60 10 56.6 ^R	2.7	0.99	18.4	0.1372 ¹		Sy1
J131931.7+140524	87GB 131703.8+142108	8.98e-12	13 19 31.67	+14 05 34.4	77.0	1.00	17.8			BLLAC
J131957.2+523533	RX J1319.9+5235	7.55e-12	13 19 56.74	+52 35 34.4	3.8	1.00	18.2	0.0922 ¹		Sy2
J132014.2+043053	RX J1320.2+0430	1.65e-12	13 20 17.85	+04 30 37.5	33.4	0.95	(17.7)		cD?	
J132016.3+330828 [†]	NGC 5098 NED02	1.93e-12	13 20 14.86	+33 08 37.3	83.4	1.00	15.0	0.0362	S?	
J132248.5+545526	RX J1322.8+5455	3.16e-12	13 22 49.19	+54 55 30.4	10.3	1.00	15.6	0.0640		Sy1
J132301.1+044000		4.63e-12	13 23 01.04	+04 39 48.7	37.0	1.00	(17.9)			
J132358.9+140614	87GB 132132.4+142106	1.57e-12	13 23 58.48	+14 06 01.2	31.2	1.00	18.2			
J132400.2+573918	87GB 132204.6+575429	2.29e-12	13 24 00.63	+57 39 18.5	49.5 ^S	1.00	16.5	0.1150		AGN
J132419.9+041913	TXS 1321+045	2.47e-12	13 24 19.71	+04 19 06.8	158.7	1.00	17.6			
J132519.4-382445		6.30e-12	13 25 18.61	-38 24 40.4 ^O	6.8	0.15	(18.9)			
	IRAS 13224-3809		13 25 19.32	-38 24 52.5 ^O	6.8	0.85	(15.3)	0.0667	Starburst	Sy1
J132542.1-022800 [†]		2.39e-12	13 25 41.73	-02 28 09.2	9.0	0.96	0.60			
J132549.2+591937 [†]		2.87e-12	13 25 49.10	+59 19 49.1 ^O	3.7	0.43	(17.5)			
			13 25 49.71	+59 19 48.3 ^O	3.7	0.56	(16.9)	0.1515		
J132615.0+293332	87GB 132354.7+294853	3.58e-12	13 26 14.99	+29 33 31.0 ^F	117.2 ^S	1.00	19.8	0.4310		BLLAC
J132617.4+001329 [†]		4.51e-12	13 26 17.45	+00 13 25.5	21.6	1.00	15.8	0.0825 ¹		Early
J132617.7+123004		6.01e-12	13 26 17.59	+12 30 06.5	58.1	1.00	(17.8)			
J132711.7-245126	GSC 6717 00254	4.43e-12	13 27 12.60	-24 51 46.7 ^R	4.1 ^E	0.97	15.4	0.0410		Sy1
J132758.9-313027 [†]	ESO 444- G 046	6.07E-11	13 27 57.07	-31 29 45.3	4.3	—	13.9	0.0469	cD;E+4	
J132806.6-271959	IRAS F13253-2704	3.12e-12	13 28 10.02	-27 19 54.9 ^O	16.4 ^E	0.62	15.0	0.0418	Sc	
J132833.4+114520		3.85e-12	13 28 33.58	+11 45 21.3	2.8	1.00	(19.5)			

Table 2—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position		S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	(J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J132953.8+471143	MESSIER 051a	1.75e-12	13 29 52.65	+47 11 45.7	430.3	1.00	9.0	0.0015	SA(s)bc pec	Sy2.5
J133107.5-014207		2.39e-12	13 31 04.09	-01 42 29.6 ^F	2.5	0.97	20.1			
J133113.6-252406	ESO 509- G 038	2.20e-11	13 31 13.79	-25 24 09.6	12.9	1.00	14.7	0.0260	S	Sy1
J133226.0-330812 [†]	PGC 047626	7.82e-12	13 32 23.65	-33 08 25.4 ^O	1425.0 ^E	—	15.5	0.0480	E+5	
J133253.5+020047		3.61e-12	13 32 53.66	+02 00 44.0	2666.8 ^T	1.00	19.5	0.2160	N galaxy	Sy1
J133517.2+405958	4C +41.26	3.73e-12	13 35 19.86	+41 00 04.0	858.0	1.00	16.0	0.2278		
J133530.6-295031	MS 1332.6-2935	7.21e-12	13 35 29.75	-29 50 39.0 ^O	10.6	0.94	19.1	0.2500		BLLAC
			13 35 30.19	-29 50 27.4 ^O	10.6	0.06	(16.7)			
J133657.0-295207	MESSIER 083	6.63e-12	13 37 00.52	-29 51 55.1 ^R	405.5 ^E	1.00	8.2	0.0017	SAB(s)c	HII
J133718.8+242306	[HB89] 1334+246	2.82e-11	13 37 18.73	+24 23 02.8	20.0	1.00	15.0	0.1076		Sy1
J133740.3-125727	[HB89] 1334-127	3.96e-12	13 37 39.83	-12 57 24.6	2676.6	1.00	17.2	0.5390	Opt.var.	BLLAC/HPQ
J133938.5+183055	RX J1339.6+1830	2.07e-12	13 39 38.31	+18 31 03.6	3.1	1.00	18.4			
J133950.5+155932	RX J1339.8+1559	1.84e-12	13 39 50.22	+15 59 30.3	4.4	1.00	18.0			QSO
J134029.9+441007	87GB 133822.3+442514	3.67e-12	13 40 29.46	+44 10 06.9	57.2	1.00	18.9			QSO
J134104.8+395942 [†]	FIRST J134105.1+395945	6.98e-12	13 41 05.25	+39 59 48.5	88.8	0.98	(17.4)			
J134112.5-143836	CTS 0023	9.48e-12	13 41 12.53	-14 38 38.7	5.6	1.00	16.0	0.0418		Sy1
J134134.5+090359		2.41e-12	13 41 33.53	+09 03 55.2	49.4 ^E	1.00	(16.6)			
J134152.6+262230 [†]	UGC 08669 NED02	7.12e-12	13 41 50.62	+26 22 23.7 ^R	307.3 ^S	0.97	15.3	0.0753	Elliptical	
J134545.1+533300	87GB 134352.4+534755	2.84e-12	13 45 45.59	+53 32 54.9	421.6 ^E	1.00	15.8			
J134730.5-114455 [†]		8.29e-12	13 47 30.67	-11 45 08.6	47.6 ^E	1.00	(19.0)	0.4500		
J134852.6+263541 [†]	CGCG 162-010	4.38e-11	13 48 52.43	+26 35 33.6	925.0	1.00	15.2	0.0633	cD;S0?	
J134853.8+075704		2.16e-12	13 48 53.08	+07 56 44.2	53.2 ^S	1.00	(18.2)			
J134919.0-301830	IC 4329A	8.05e-11	13 49 19.23	-30 18 33.7	66.8	1.00	14.0	0.0161	SA0 ⁺ : sp	Sy1
J135022.2+094007	PMN J1350+0940	6.25e-12	13 50 22.16	+09 40 10.8	292.9	1.00	(16.3)			
J135045.7+233146	MG2 J135044+2331	1.52e-12	13 50 45.97	+23 31 42.7	164.5 ^T	1.00	16.6	0.5289 ¹		QSO
J135120.9+111458	RGB J1351+112	1.89e-12	13 51 20.91	+11 14 52.8	42.8	1.00	18.0			
J135227.7+490825	87GB 135030.6+492245	1.32e-12	13 52 29.04	+49 08 22.7	70.8	1.00	18.7			
J135300.0+294742	RX J1353.0+2947:046	2.34e-12	13 53 02.00	+29 47 38.3 ^O	3.3	0.64	18.5			AGN
			13 53 02.08	+29 47 43.2 ^O	3.3	0.21	(19.4)			
J135304.8+691832	UGC 08823	4.15e-11	13 53 03.88	+69 18 28.7	24.0	1.00	14.6	0.0294	S0	Sy1.5
J135328.2+560102	RX J1353.4+5601	2.99e-12	13 53 28.08	+56 00 55.3	14.9	1.00	18.8	0.3700		BLLAC
J135336.2+263148	[HB89] 1351+267	2.77e-12	13 53 35.77	+26 31 52.4	246.2 ^T	1.00	17.2	0.3100	Opt.var.	QSO
J135420.2+325547	UGC 08829	9.97e-12	13 54 19.95	+32 55 47.7 ^F	3.7	1.00	14.6	0.0260	SBa	Sy1
J135447.1-104100	[HB89] 1352-104	3.67e-12	13 54 46.93	-10 41 02.4	768.7 ^E	1.00	18.4	0.3320		QSO
J135515.9+561244	RX J1355.2+5612	6.19e-12	13 55 16.87	+56 12 45.9	6.1	1.00	17.1	0.1220		Sy1
J135553.3+383427	MRK 0464	1.83e-12	13 55 53.39	+38 34 28.0	16.4	1.00	16.5	0.0510	Compact	Sy1.5
J135703.6+191915	[HB89] 1354+195	1.78e-12	13 57 04.37	+19 19 08.1	2586.3 ^E	1.00	16.0	0.7200	Opt.var.	QSO
J135738.7+012813	RX J1357.6+0128	2.61e-12	13 57 38.69	+01 28 13.6	61.3	1.00	(17.3)			
J135851.7+251148	RX J1358.8+2511	2.66e-12	13 58 52.77	+25 11 46.0 ^R	4.5 ^E	1.00	(16.6)	0.0888 ¹		Sy1
J140102.1+025249 [†]	ABELL 1835:[AEF92]	1.23e-11	14 01 02.05	+02 52 41.0	41.4	1.00	19.3	0.2523		
J140137.0-110735	GSC 5557 00266	1.02e-11	14 01 36.20	-11 07 47.5	4.8	0.99	14.6	0.0691		
J140337.0-335840 [†]	NGC 5419	7.07e-12	14 03 38.71	-33 58 46.9	350.0 ^E	1.00	11.9	0.0139	E	

Table 2—Continued

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position	S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α (J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J140450.2+655433	RX J1404.8+6554	2.85e-12	14 04 50.06 +65 54 28.4	15.4	1.00	19.4	0.3640		QSO
J140451.2+040206	[HB89] 1402+042	3.75e-12	14 04 50.92 +04 02 02.4	42.3	1.00	17.5	0.2000	Opt.var.	BLLAC
J140622.2+222350	PG 1404+226	7.65e-12	14 06 21.40 +22 23 44.3	3.3	1.00	15.8	0.0980		Sy
J140630.1+123632		2.29e-12	14 06 29.84 +12 36 21.3	6.2	1.00	(19.8)			
J140630.3-393508		4.09e-12	14 06 30.29 -39 35 10.4	8.6	1.00	(19.7)			
J140640.3+481824	RX J1406.6+4818	1.46e-12	14 06 41.26 +48 18 28.2	7.0 ^E	1.00	(19.0)			
J140659.1+164204		4.89e-12	14 06 59.10 +16 42 09.3	8.3	1.00	(18.1)			
J140728.4-270055 [†]	IC 4374	3.17e-11	14 07 29.83 -27 01 04.2	646.2	1.00	13.6	0.0218	SA(rs)0 ⁻ :	
J140827.9+240934	RX J1408.4+2409	3.68e-12	14 08 27.82 +24 09 24.4 ^F	3.9	1.00	16.6	0.1310 ¹		Sy1
J140918.9+135250		2.93e-12	14 09 19.07 +13 52 41.3	7.2	1.00	(19.0)			
J140923.6+593943	MS 1407.9+5954	1.68e-12	14 09 23.46 +59 39 37.3	36.5	1.00	19.7	0.4950		BLLAC
J140924.1+261827	PG 1407+265	2.24e-12	14 09 23.96 +26 18 22.4	9.6 ^E	1.00	15.7	0.9400		QSO
J141029.6+282112	RX J1410.4+2821	1.66e-12	14 10 29.43 +28 20 55.5	28.8	1.00	18.9			QSO
J141031.6+610020	RX J1410.5+6100	2.74e-12	14 10 31.67 +61 00 09.7	11.4	1.00	20.1	0.3840		QSO
J141315.2-031223	NGC 5506	2.73e-12	14 13 14.84 -03 12 27.0	339.4	1.00	13.4	0.0062	Sa pec sp	Sy1.9
J141342.6+433938 [†]	87GB 141144.3+435404	4.41e-12	14 13 43.68 +43 09 45.5	48.5	1.00	18.4	0.0890	cD	BLRG
J141357.3+711751 [†]		1.76e-12	14 13 55.56 +71 17 19.6	9.1 ^A	0.96	(19.2)			
J141358.3+764456		3.73e-12	14 13 57.96 +76 44 57.2	7.1	1.00	(16.5)			
J141441.9+165812		1.66e-12	14 14 41.64 +16 58 12.3 ^O	2.6	0.39	(16.6)			
	RX J1414.6+1658		14 14 42.01 +16 58 07.6 ^O	2.6	0.58	16.2			QSO
J141554.4-070516		2.31e-12	14 15 54.87 -07 05 34.2	5.6	1.00	(16.8)			
J141628.4+124215	MG1 J141630+1242	2.30e-12	14 16 28.64 +12 42 13.5	110.6	1.00	17.6			
J141650.6-115845	LCRS B141408.4-114506	5.72e-12	14 16 49.63 -11 59 00.3	3.5	1.00	15.7	0.0986		
J141659.6+135443		1.61e-12	14 17 01.90 +13 54 16.5 ^{OR}	31.5 ^D	0.56	(21.5)			
J141756.8+254329	[HB89] 1415+259	2.34e-11	14 17 56.27 +25 43 23.1	89.6	1.00	16.0	0.2370	Opt.var.	BLLAC
J141759.6+250817	NGC 5548	7.74e-11	14 17 59.32 +25 08 13.5	29.1	1.00	13.3	0.0172	(R')SA(s)0/a	Sy1.5
J141901.8+773229		2.32e-12	14 19 00.47 +77 32 26.3	8.1	1.00	(17.2)			
J141909.5+062836	[HB89] 1416+067	5.62e-12	14 19 08.18 +06 28 36.3	6100.8	1.00	16.8	1.4360		QSO
J141922.5-263842	ESO 511- G 030	3.51e-11	14 19 22.29 -26 38 40.9	13.4 ^E	1.00	13.3	0.0224	SA(rs)c pec	Sy1
J141946.4+542324	PG 1418+546	1.19e-12	14 19 46.50 +54 23 15.1	788.7	1.00	15.6	0.1510	Opt.var.	BLLAC
J141950.3-192811	EC 14170-1914	1.31e-11	14 19 49.65 -19 28 35.1	1974.1 ^D	0.98	17.5	0.1200	N galaxy	Sy1
J142136.4+493305 [†]	MCG +08-26-021	5.05e-12	14 21 36.35 +49 33 05.9	4.0	0.98	16.0	0.0710	cD	
J142139.7+371743	FIRST J142140.4+371731	3.13e-12	14 21 40.50 +37 17 31.2 ^F	3.3	1.00	(16.5)	0.1600	cD	
J142219.8+294257	IRAS 14201+2956	1.29e-12	14 22 20.23 +29 42 55.0 ^F	3.0	1.00	15.9	0.0530		Sy1
J142239.1+580159	RGB J1422+580	2.44e-11	14 22 38.55 +58 01 57.6	13.2	1.00	19.0	0.6380		BLLAC
J142309.0+261518	NGC 5594	1.56e-12	14 23 10.31 +26 15 56.8 ^F	14.9	0.98	15.1	0.0375		
J142313.4+505537	87GB 142127.2+510856	4.64e-12	14 23 14.17 +50 55 38.7	178.3	1.00	16.0	0.2740		Sy1
J142337.3-173057		5.02e-12	14 23 37.29 -17 31 24.7	3.3	0.99	(17.4)			
J142353.6+401533	FIRST J142351.5+401531	2.16e-12	14 23 51.55 +40 15 28.9	12.5	1.00	17.2	0.0820 ¹		Early
J142356.0+262630	RX J1423.9+2626:074	2.22e-12	14 23 55.38 +26 26 22.9	22.5	1.00	19.0			
J142422.9+343407	FIRST J142422.9+343356	1.69e-12	14 24 22.75 +34 33 57.2	10.0	1.00	(18.7)			
J142425.2+595254		2.41e-12	14 24 24.08 +59 53 08.9	2.4	0.99	(16.5)	0.1348 ¹		Sy1

Table 2—Continued

(1) <i>ROSAT</i> BSC 1RXS name	(2) NED Common name	(3) S_x ergs s ⁻¹ cm ⁻²	(4) Position α (J2000) δ	(5) $S_{1.4\text{GHz}}$ mJy	(6) ID P	(7) m_B	(8) Redshift	(9) NED Morphology	(10) Sp. Class.
J142601.3+374935	ABELL 1914:001	9.44e-12	14 25 59.60 +37 49 39.8 ^O 14 26 01.78 +37 49 27.8 ^O 14 26 02.61 +37 49 51.9 ^O	19.6 ^A 19.6 ^A 19.6 ^A	0.07 0.49 0.34	17.2 (20.5) (18.0)	0.1712	cD?	
J142700.5+234803	RX J1426.0+3749:008 PG 1424+240	6.82e-12	14 27 00.38 +23 48 00.7	430.1	1.00	16.5			BLLAC
J142715.0+440726	87GB 142522.4+442128	1.52e-12	14 27 16.14 +44 07 32.1	48.6	1.00	(21.5)			
J142740.6-252106		5.08e-12	14 27 38.95 -25 20 54.3	3.5	0.91	(17.1)			
J142832.6+424028	87GB 142634.5+425353	5.21e-11	14 28 32.54 +42 40 22.3	58.8	1.00	16.5	0.1290		BLLAC
J142906.7+011708	PG 1426+015	2.06e-11	14 29 06.93 +01 17 06.3	3.6	1.00	17.5	0.0865	Compact	Sy1
J143104.8+281716	PG 1428+285	7.67e-12	14 31 05.05 +28 17 21.4	3.7	1.00	15.2	0.0458	S?	Sy1
J143106.2+253815	ABELL 1927:[HO98] 201	4.21e-12	14 31 06.64 +25 38 00.5	8.7	1.00	(16.3)	0.0959 ¹		Early
J143150.6+133256		1.40e-12	14 31 49.67 +13 31 59.1 ^{OR} 14 31 50.08 +13 32 05.2 ^{OR}	10.4 ^E 10.4 ^E	0.35 0.29	(19.4) (16.8)			
J143236.0+313855 [†]	RX J1432.5+3138:062	2.21e-12	14 32 37.95 +31 38 49.2 ^F	10.0	1.00	18.1			
J143509.2+313203	FIRST J143509.5+313148	1.15e-12	14 35 09.57 +31 31 49.5	43.2	1.00	(18.7)			
J143527.9+550747 [†]	RGB J1435+551	1.66e-12	14 35 27.97 +55 07 56.0	475.7 ^D	1.00	15.2	0.1402		
J143622.2+584750	UGC 09412	1.38e-12	14 36 22.07 +58 47 41.6	11.7	1.00	14.5	0.0315	SBc	Sy1.5
J143649.6-161344	EC 14340-1600	2.08e-11	14 36 49.57 -16 13 39.0	99.7 ^S	1.00	15.8	0.1440		QSO
J143658.7+563934	RX J1436.9+5639	2.74e-12	14 36 57.77 +56 39 25.4	21.3	1.00	18.8			BLLAC
J143830.1+353922	FIRST J143830.2+353915	1.26e-12	14 38 30.51 +35 39 13.7	4.8	1.00	(18.4)			
J143917.7+393248	PG 1437+398	1.53e-11	14 39 17.46 +39 32 42.5	42.8	1.00	16.0			BLLAC
J144011.6+615633	IRAS F14390+6209	2.64e-12	14 40 13.25 +61 56 43.4	3.7	0.95	16.4	0.2760		Sy1
J144037.4-384658		3.59e-11	14 40 37.88 -38 46 52.8	23.4	1.00	(17.3)			
J144157.2+160439		1.52e-12	14 41 57.73 +16 04 38.5	2.6	0.96	(16.4)			
J144207.7+352632	PG 1440+356	6.20e-11	14 42 07.39 +35 26 24.9	5.4	1.00	14.9	0.0791	Compact	Sy1
J144218.9+221820	UGC 09480 NOTES01	5.84e-12	14 42 19.37 +22 18 11.4	90.1 ^S	1.00	(16.2)			
J144248.5+120042	87GB 144023.9+121335	1.20e-11	14 42 48.26 +12 00 40.4	69.2	1.00	17.1	0.1620		BLLAC
J144302.8+520141	[HB89] 1441+522	5.12e-12	14 43 02.58 +52 01 37.0 ^F	2543.9	1.00	20.0	1.570		QSO
J144357.0-390847		1.85e-11	14 43 57.09 -39 08 36.7	110.3	1.00	(16.3)			
J144428.4+311304 [†]	87GB 144222.3+312626	1.45e-12	14 44 27.99 +31 13 14.5	63.1	1.00	22.0			
J144433.9+633604	WN B1443+6348	2.09e-12	14 44 34.73 +63 36 05.4	18.9	1.00	19.6	0.2990		BLLAC
J144505.9-032613	RBS 1424	7.90e-12	14 45 06.23 -03 26 13.4	21.6	1.00	17.2			BLLAC
J144652.9-341323		4.63e-12	14 46 51.79 -34 13 17.7 ^O 14 46 51.95 -34 13 11.9 ^O	3.3 3.3	0.39 0.55	(18.4) (16.9)			
J144732.7+345507	IRAS F14455+3507	2.87e-12	14 47 33.01 +34 55 07.0	16.3	1.00	(17.1)	0.6610 ²		Sy1
J144801.0+360833	[WB92] 1446+3620	6.37e-12	14 48 00.62 +36 08 31.2	36.2	1.00	16.3	- ²		BLLAC?
J144932.3+274630	RBS 1434	6.84e-12	14 49 32.96 +27 46 21.0	90.7	1.00	19.1	0.2250		BLLAC
J145005.5+463520	CBS 0287	1.84e-12	14 50 05.15 +46 35 21.3 ^F	4.5	1.00	18.0	0.2920		
J145054.0-071423		4.29e-12	14 50 54.48 -07 14 14.0	3.8	0.95	16.6	0.0722 ¹		Sy1.9
J145108.2+143654		2.56e-12	14 51 08.25 +14 37 05.2 ^O 14 51 09.43 +14 36 43.3 ^O	3.3 3.3	0.15 0.84	(19.0) (16.8)		cD	
J145108.5+270933	PG 1448+273	1.54e-11	14 51 08.81 +27 09 26.7 ^F	3.4	1.00	15.0	0.0650		Sy1
J145127.5+635421	RX J1451.4+6354	2.67e-12	14 51 27.85 +63 54 26.2	10.0	1.00	19.6	0.6500		QSO

Table 2—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position		S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	(J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J145224.0+452234	B3 1450+455	2.24e-12	14 52 24.66	+45 22 23.8 ^{RF}	413.6 ^D	1.00	17.3	0.4690		Sy1
J145307.8+215333 [†]	RX J1453.1+2153:019	2.64e-12	14 53 09.50	+21 54 05.6	42.9 ^E	0.99	18.6			
J145323.4-182609		4.88e-12	14 53 24.34	-18 26 19.1	4.2	0.97	(15.5)			
J145427.2-374738	[HB89] 1451-375	8.47e-12	14 54 27.59	-37 47 34.6	942.7 ^D	1.00	16.7	0.3140		QSO
J145431.4+183834 [†]	NGC 5778	8.93e-12	14 54 31.55	+18 38 32.9	39.4	1.00	14.8	0.0589	D	
J145434.1+080250 [†]		2.64E-12	14 54 34.29	+08 03 36.6	46.7	—	(16.0)			
J145507.9+192025 [†]	RGB J1455+193B	2.68e-12	14 55 08.23	+19 20 15.2 ^R	40.3 ^E	1.00	(18.2)			
J145603.4+504824	RX J1455.0+1920	2.14e-11	14 56 03.62	+50 48 26.0 ^{RF}	245.3 ^D	1.00	15.7			
J145613.0-382628		7.05e-12	14 56 15.60	-38 26 59.9 ^O	6.4	0.52	(15.7)			
J145715.4+222026 [†]	MS 1455.0+2232:[AEF92]	6.04e-12	14 57 15.18	+22 20 36.0	17.6	1.00	(17.8)	0.2578		
J145827.3+483250	RX J1458.4+4832	6.11e-12	14 58 27.35	+48 32 46.2 ^F	3.1 ^E	1.00	20.2	0.5390		QSO
J145904.1-084254 [†]		4.80e-12	14 59 05.22	-08 42 36.0 ^O	28.4 ^E	—	(18.7)			
J145928.8-181034	PMN J1459-1810	9.36e-12	14 59 28.84	-18 10 44.8	105.7	1.00	(16.3)			
J145958.6+333704	87GB 145756.1+334845	1.51e-12	14 59 58.49	+33 37 00.9	15.6	1.00	17.1	0.6450 ²		Sy1
J150020.7+212213 [†]	RX J1500.3+2122	9.33e-12	15 00 19.54	+21 22 10.7	26.9	1.00	15.6	0.1532	S0	
J150101.7+223812	MS 1458.8+2249	8.02e-12	15 01 01.89	+22 38 05.8	32.4	1.00	16.8	0.2350		BLLAC
J150111.6+014151	NGC 5813	1.43e-11	15 01 10.98	+01 42 06.9	15.8 ^E	1.00	11.4	0.0064	E1-2	
J150343.0-154107		2.45e-11	15 03 40.67	-15 41 13.7 ^O	5.7	0.84	(17.1)			
J150407.4-024820	LCRS B150131.5-023636	3.06e-11	15 04 07.55	-02 48 16.3	61.7	1.00	16.4	0.2169		
J150413.1+685610	[HB89] 1503+691	4.78e-12	15 04 12.51	+68 56 14.8 ^R	459.4 ^T	1.00	17.0	0.3180		Sy1
J150629.3+013628	NGC 5846	1.24e-11	15 06 29.32	+01 36 19.0	22.1 ^E	1.00	11.1	0.0061	E0-1	LINER/HII
J150636.4-054011	RHS 45	4.65e-12	15 06 37.02	-05 40 06.4	15.2	1.00	20.0			
J150644.5+081342	RX J1506.7+0813	2.54e-12	15 06 44.48	+08 13 58.7	84.6	1.00	18.5			QSO
J150744.6+512709	MRK 0845	3.65e-12	15 07 44.43	+51 27 02.7	3.8	0.98	15.6	0.0461		Sy1
J150842.2+270910	RBS 1467	6.25e-12	15 08 42.69	+27 09 08.6	39.9 ^E	1.00	16.2			BLLAC
J151040.8+333515	FIRST J151041.1+333504	3.99e-12	15 10 41.97	+33 35 08.8	8.8 ^E	1.00	(16.8)			
J151056.3+054431 [†]	IC 1101	7.01e-11	15 10 55.87	+05 44 39.3	540.9	1.00	14.7	0.0780	SO ⁻ ;cD	
J151127.2+062153 [†]	UGC 09756 NED02	1.09E-11	15 11 26.48	+06 20 55.5	566.2	—	15.0	0.0805		
J151159.4-211903	IRAS 15091-2107	1.35e-11	15 11 59.82	-21 19 00.1	47.3	1.00	15.5	0.0446		Sy1
J151213.1-225515		5.34e-12	15 12 12.84	-22 55 08.3	19.2	1.00	(18.4)			
J151420.8+424445	IRAS F15125+4255	1.98e-12	15 14 20.53	+42 44 45.3 ^F	6.6 ^E	1.00	16.0			
J151443.1+365059	PG 1512+370	3.24e-12	15 14 43.20	+36 50 49.9	989.7 ^D	1.00	15.5	0.3707		Sy1?
J151523.3+553057	NGC 5905	4.06e-12	15 15 23.30	+55 31 00.6	21.4 ^E	1.00	12.5	0.0113	SB(r)b	Sy1
J151601.5+020055	GNZ 095	3.56e-12	15 16 01.56	+02 01 01.7	13.9	1.00	16.9			
J151618.7-152347	RBS 1478	1.52e-11	15 16 18.08	-15 23 43.9	8.0 ^E	0.97	19.6			BLLAC
J151642.1+291803	FIRST J151641.5+291809	2.18e-12	15 16 41.55	+29 18 06.5	135.3	1.00	16.2	0.1300		BLLAC?
J151642.4+070058 [†]	UGC 09799	5.30e-11	15 16 44.55	+07 01 17.8 ^O	5500.3	0.87	14.0	0.0342	cD;E	
J151747.3+652522	87GB 151706.4+653539	1.99e-11	15 17 47.55	+65 25 23.6	37.7	1.00	15.5			BLLAC
J151750.8+050615	CGCG 049-106	2.45e-12	15 17 51.69	+05 06 23.9	12.6	0.97	15.6	0.0387		
J151845.3+061340 [†]	PMN J1518+0613	1.20e-11	15 18 45.75	+06 13 55.8 ^R	498.0 ^E	1.00	15.2	0.1021		AGN
J152053.7+484006	MCG +08-28-020	5.86e-12	15 20 52.34	+48 39 38.4	82.5	1.00	16.0	0.0738 ¹		Early
J152151.0+074221 [†]	NGC 5920	3.57e-11	15 21 51.76	+07 42 29.2 ^R	117.4 ^S	0.97	14.7	0.0453	cD;S0:	

Table 2—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position		S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	(J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J152225.9+274232		2.63e-11	15 22 27.46	+27 42 23.1 ^O	13.4 ^A	0.13	(19.8)			
	ABELL 2065:[PGH88] 166		15 22 27.64	+27 42 35.0 ^O	13.4 ^A	0.54	16.1	0.0707		
J152228.1-064438	[HB89] 1519-065	5.34e-12	15 22 28.65	-06 44 39.6	15.0	1.00	14.9	0.0830		Sy
J152305.7+083550 [†]	CGCG 077-097	4.51e-11	15 23 05.06	+08 36 36.2 ^O	15.1	0.56	13.6	0.0344	cD;S	
J152346.0+633930	4C +63.22	5.63e-12	15 23 45.91	+63 39 24.1 ^O	695.5 ^T	1.00	15.9	0.2040		BLRG
J152412.7-315427		1.71e-11	15 24 12.93	-31 54 24.7	50.1	1.00	(16.1)			
J152502.6+110743	[HB89] 1522+113	3.43e-12	15 25 02.88	+11 07 45.0	408.1	1.00	18.0	0.3310		QSO
J152558.6+181423		5.31e-12	15 25 58.52	+18 14 10.7 ^O	3.9 ^E	0.36	(18.5)			
	RX J1525.9+1814		15 25 58.52	+18 14 16.1 ^O	3.9 ^E	0.63	16.3			QSO
J152907.5+561604	IRAS F15279+5626	8.41e-12	15 29 07.53	+56 16 09.1	5.4	1.00	15.8	0.0990		Sy1
J152957.7-130042		4.40e-12	15 29 57.98	-13 00 34.3 ^O	3.2 ^A	0.45	(19.1)			
	EC 15272-1250		15 29 58.30	-13 00 39.3 ^O	3.2 ^A	0.32	15.7			
J153118.2+072713	NGC 5940	6.27e-12	15 31 17.94	+07 27 29.1	9.3	0.96	14.3	0.0337	SBab	Sy1
J153202.3+301631	87GB 152959.0+302636	6.17e-12	15 32 02.27	+30 16 29.4	54.4	1.00	12.1	0.0640 ¹		BLLAC?
J153253.7+302103	RBS 1509	3.90e-12	15 32 53.77	+30 20 59.8	23.3	1.00	15.8	0.3620		LINER
J153311.7+185427	87GB 153052.7+190538	1.47e-11	15 33 11.40	+18 54 29.0	22.9	1.00	17.1			
J153324.9+341640	87GB 153121.5+342710	4.81e-12	15 33 24.26	+34 16 40.3	30.0	1.00	17.1	— ²		BLLAC
J153501.1+532042	RX J1535.0+5320	1.75e-11	15 35 00.66	+53 20 36.2	18.2	1.00	17.6	0.8900		BLLAC
J153528.7+392242	FIRST J153529.0+392246	2.20e-12	15 35 29.24	+39 22 45.3	19.7	1.00	18.7	0.2570 ²		BLLAC?
J153548.0+432251	RX J1535.8+4322	1.76e-12	15 35 47.21	+43 22 43.2 ^R	25.8 ^T	1.00	(19.0)			
J153552.0+575404	PG 1534+580	1.31e-11	15 35 52.56	+57 54 16.7	5.2	1.00	15.2	0.0296	E1?	Sy1
J153646.8+013756	87GB 153415.4+014705	1.14e-11	15 36 46.73	+01 38 00.0	72.9	1.00	18.7	0.3120		BLLAC
J153846.0-102641		5.47e-12	15 38 46.71	-10 26 21.3	3.8	0.99	(16.7)	0.1958 ¹		Sy1
J153941.8-112838	PMN J1539-1128	7.29e-12	15 39 41.26	-11 28 36.2	78.7	1.00	(16.9)			
J153950.3+304305 [†]	FIRST J153950.7+304303	7.92e-12	15 39 50.76	+30 43 06.3	10.2	1.00	18.1	0.0968		
J154009.4+141116 [†]	87GB 153746.8+142023	3.24e-12	15 40 07.85	+14 11 36.9 ^R	111.7 ^D	1.00	16.1		* or G?	
J154016.4+815504	RBS 1524	1.31e-11	15 40 15.69	+81 55 05.6	69.9	1.00	15.3			BLLAC
J154127.5+044408	SHK 360:[STC97c] 01	2.74e-12	15 41 26.54	+04 43 55.9	27.2	1.00	(16.6)	0.1103		
J154203.2-291516		4.84e-12	15 42 03.02	-29 15 09.5	17.4	1.00	(18.4)			
J154348.9+401343	FIRST J154348.5+401324	2.19e-12	15 43 48.10	+40 13 31.3	3.6	1.00	18.0	0.3200 ²		Sy1
J154530.7+484614	PG 1543+489	1.49e-12	15 45 29.84	+48 46 26.6	4.1	0.99	16.0	0.4000		QSO
J154546.3-233933		8.25e-12	15 45 46.72	-23 39 29.9	47.4 ^E	1.00	(16.4)			
J154732.3+102446		5.12e-12	15 47 31.72	+10 24 50.3	3.3	1.00	(16.6)	0.1390 ¹		Sy1
J154744.2+205156	PG 1545+210	7.77e-12	15 47 43.53	+20 52 17.1 ^O	2397.6 ^D	1.00	16.7	0.2643	E3(?);Opt.var.	QSO
J154751.7+025538	MS 1545.3+0305	3.56e-12	15 47 51.98	+02 55 51.3 ^F	3.6	1.00	16.8	0.0980		Sy1
J154824.9-134532	NGC 5995	9.69e-12	15 48 24.90	-13 45 28.5	30.7	1.00	14.5	0.0252	S(B)c	Sy2
J154843.3-320717	ESO 450-PN?016	1.15e-11	15 48 43.51	-32 07 10.9	11.7	1.00	14.5	0.0489	PN or G?	
J154846.4-030012	ABELL 2128:[QR95] 1	5.01e-12	15 48 43.10	-02 59 31.1	11.9	0.99	(16.4)	0.0574		
J154849.7-225107		3.42e-11	15 48 49.80	-22 51 01.9	145.4	1.00	(16.4)			
J155333.4-311841		4.03e-12	15 53 33.57	-31 18 31.6	155.9	1.00	(16.7)			
J155424.3+201116	87GB 155210.3+202028	7.21e-12	15 54 24.17	+20 11 24.2	79.4	1.00	17.7	0.2220		BLLAC
J155539.4-093848	PMN J1555-0939	4.66e-12	15 55 39.40	-09 38 48.1 ^R	391.8 ^T	1.00	(16.1)		asym. dbl.	

Table 2—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position		S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	(J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J155543.2+111114	[HB89] 1553+113	3.67e-11	15 55 43.09	+11 11 24.5	312.4	1.00	15.0	0.3600	Opt.var.	BLLAC
J155611.0+662123 [†]	ABELL 2146:[AEF92]	4.59e-12	15 56 13.96	+66 20 53.8 ^O	15.8 ^A	0.71	(18.5)	0.2337		Sy2
J155625.4+090311	MRK 0863	2.75e-12	15 56 25.95	+09 03 14.8	4.8	1.00	15.3	0.0423	SB nuc.	AGN
J155745.0+353020	RX J1557.7+3530:024	2.86e-12	15 57 42.14	+35 30 24.7	330.8 ^D	0.98	18.2			
J155818.7+255118	MRK 0864	4.75e-12	15 58 18.21	+25 51 23.7	3.1	1.00	16.5	0.0719		Sy2
J155822.4-141005	PKS 1555-140	2.32e-11	15 58 21.95	-14 09 59.3	461.2	1.00	(17.3)	0.0970		
J155909.5+350144	UGC 10120	8.85e-12	15 59 10.07	+35 01 45.6	5.9 ^S	1.00	14.6	0.0319	SB(r)b	Sy1
J160025.5+331411		3.21e-12	16 00 22.74	+33 13 17.8 ^F	9.5	0.87	(16.7)			
J160215.3+155805	UGC 10143	1.09e-11	16 02 17.02	+15 58 29.1 ^{OR}	11.5	0.83	14.2	0.0347	cD;E+	
J160258.4+421202	RGB J1602+422	3.58e-12	16 02 59.11	+42 12 01.9	15.1	1.00	18.9			
J160328.2-093502	IRAS F16007-0926	5.20e-12	16 03 28.29	-09 35 17.4	5.4	1.00	(16.4)			
J160337.5+155353	GIN 463	7.97e-12	16 03 38.05	+15 54 03.2	97.3	1.00	15.5	0.1095	E/S0	
J160435.1+174333 [†]	NGC 6041A	1.23e-11	16 04 35.77	+17 43 39.0 ^R	13.7 ^D	—	13.7	0.0348	cD;SA0 ⁻	
J160456.8+235604 [†]	NGC 6051	1.47e-11	16 04 56.79	+23 55 57.4	666.9 ^T	1.00	14.2	0.0320	cD;E	
J160518.5+542101		5.90e-12	16 05 19.35	+54 20 59.9	7.6	1.00	(19.0)			
J160534.7+162559		2.86e-12	16 05 34.21	+16 25 47.1 ^{OR}	5.2 ^A	0.06	(17.6)			
			16 05 35.23	+16 25 52.9 ^{OR}	5.2 ^A	0.30	(19.1)			
			16 05 35.59	+16 25 56.7 ^{OR}	5.2 ^A	0.37	(18.7)			
			16 05 35.84	+16 26 15.4 ^{OR}	5.2 ^A	0.18	(17.8)			
J160621.0+563017		3.38e-12	16 06 20.61	+56 30 19.0	15.7	1.00	(19.2)			
J160820.9+601827	RGB J1608+603B	1.73e-12	16 08 20.67	+60 18 29.5	83.2 ^T	1.00	(16.2)			
J161004.4+671030	87GB 160948.8+671803	4.66e-12	16 10 04.30	+67 10 27.1	73.1	1.00	16.8	0.0670		BLLAC?
J161047.7+330329	RX J1610.7+3303	3.42e-12	16 10 48.46	+33 03 43.8	5.2	0.99	16.7	0.0970		Sy1
J161124.8+585106	SBS 1610+589	3.62e-12	16 11 24.57	+58 51 03.8	3.6	0.97	15.8	0.0321		Sy1.5
J161204.4-043816		4.35e-12	16 12 04.87	-04 38 19.1	4.1	1.00	(21.5)			
J161301.5+371656	KUG 1611+374B	4.85e-12	16 13 01.69	+37 17 18.4	4.4	1.00	15.5	0.0696		Sy1
J161357.0+654313	PG 1613+658	1.80e-11	16 13 56.88	+65 43 10.4	4.4	1.00	15.2	0.1290	Compact	Sy1
J161413.0+260412	PG 1612+261	1.10e-11	16 14 13.20	+26 04 16.1 ^F	18.2	1.00	15.4	0.1310	Opt.var.	Sy1.5
J161546.9-060841 [†]		3.55e-11	16 15 45.99	-06 08 27.4 ^O	37.2 ^A	0.68	(17.1)			
			16 15 47.46	-06 08 16.6 ^O	31.0 ^A	0.19	(18.3)			
J161633.4+375558	FIRST J161632.8+375603	1.37e-12	16 16 32.87	+37 55 58.0	4.4	1.00	(18.5)			
J161706.7+410634	MG4 J161703+4105	2.00e-12	16 17 06.62	+41 06 45.2	95.8	1.00	16.2	0.2670 ²		BLLAC?
J161711.4+063816 [†]	4C +06.55	3.74E-12	16 17 13.48	+06 37 29.5	699.8 ^D	—	16.6			
J161745.7+060343	RX J1617.7+0603	1.63e-11	16 17 45.74	+06 03 50.8	5.7	1.00	16.5	0.0380		Sy1
J161809.2+361951	RX J1618.1+3619	1.02e-11	16 18 10.55	+36 20 07.9	2.9 ^E	0.98	16.9	0.0340		Sy1
J161849.1+442507	RX J1618.8+4425	1.29e-12	16 18 49.16	+44 25 19.5	3.1	1.00	16.6			QSO
J162032.0+295321 [†]	FIRST J162031.1+295327	6.21e-12	16 20 31.12	+29 53 27.6 ^F	5.2	1.00	18.1	0.0954	cD	
J162100.4+254547 [†]	ABELL 2177:001	2.55e-12	16 20 58.46	+25 46 13.1 ^F	3.2 ^A	0.91	17.5	0.1610		
J162111.0+374556	4C +37.46	1.48e-12	16 21 11.28	+37 46 05.5	631.5	1.00	19.0	0.3400	candidate	BLLAC
J162114.3+181936	RX J1621.2+1819	4.78e-12	16 21 14.40	+18 19 50.2 ^O	2.6	0.11	17.1			QSO
			16 21 15.37	+18 19 22.3 ^O	2.6	0.08	(19.5)			
			16 21 15.51	+18 19 39.5 ^O	2.6	0.79	(19.9)			

Table 2—Continued

(1) <i>ROSAT</i> BSC 1RXS name	(2) NED Common name	(3) S_x ergs s ⁻¹ cm ⁻²	(4) Position α (J2000) δ	(5) $S_{1.4\text{GHz}}$ mJy	(6) ID P	(7) m_B	(8) Redshift	(9) NED Morphology	(10) Sp. Class.
J162128.3+424528	B3 1619+428	1.23E-12	16 21 31.08 +42 45 23.4 ^F	121.9 ^S	0.95	17.9			
J162145.2+542724	SBS 1620+545	2.32e-12	16 21 45.11 +54 27 23.7 ^F	4.7	1.00	18.8	0.0501 ¹		Sy1.5
J162147.0-225306		6.89e-12	16 21 46.88 -22 53 10.6	8.7	1.00	(16.3)			
J162409.7+260421	IRAS F16221+2611	6.21e-12	16 24 09.17 +26 04 34.3	4.4	1.00	16.8	0.0400		Sy1
J162607.6+335902	FIRST J162607.0+335916	1.65e-12	16 26 06.78 +33 59 11.7	5.0	1.00	16.5	0.2040		Sy1
J162612.7+512028	87GB 162455.1+512734	1.95e-12	16 26 11.61 +51 20 39.3	50.3 ^E	1.00	19.5	0.1780		Sy1/BLRG
J162625.5+351331	FIRST J162625.8+351341	1.30e-12	16 26 25.85 +35 13 41.7	24.8	1.00	19.2	0.5000		BLLAC
J162645.0+630047	WN B1626+6307	3.32e-12	16 26 45.98 +63 00 51.6	15.3	1.00	15.9		E	
J162819.4+660556	WN B1628+6612	2.51e-12	16 28 19.97 +66 06 01.4	20.5	1.00	15.1		E	
J162837.7+393249 [†]	NGC 6166	4.50e-11	16 28 38.34 +39 32 60.0	3680.5 ^D	1.00	13.1	0.0310	BCG	
J162839.1+252743		2.77e-12	16 28 39.14 +25 27 56.2	72.8	1.00	(17.8)			
J162901.2+400753	FIRST J162901.3+400759	7.54e-12	16 29 01.51 +40 08 00.6	9.4	1.00	19.0	0.2720		Sy1
J163124.7+421656	RX J1631.4+4217	5.68e-12	16 31 24.81 +42 17 03.4	7.4	1.00	19.8	0.4680		BLLAC
J163208.3+133921		3.57e-12	16 32 05.49 +13 39 10.4 ^O	2.6	0.51	(18.0)			Early
	CGCG 080-045		16 32 05.27 +13 38 50.7 ^O	2.6	0.21	15.4	0.0519 ¹		Early
J163230.3+823214	NGC 6251	3.19e-12	16 32 26.14 +82 32 20.3 ^R	945.4 ^S	1.00	13.6	0.0230	E	Sy2
J163246.8+053423 [†]	TXS 1630+056	2.96e-11	16 32 46.87 +05 34 34.9	70.1	1.00	17.1	0.1523		
J163323.3+471848	IRAS 16319+4725	2.52e-12	16 33 23.64 +47 18 59.9	69.7	1.00	16.5	0.1163		Sy2
J163658.7-124838		8.15e-12	16 36 58.47 -12 48 36.5	26.2	1.00	(21.5)			
J163746.3+114941	[HB89] 1635+119	4.14e-12	16 37 46.28 +11 49 30.9	165.6 ^T	0.97	16.5	0.1460		Sy1/BLRG
J163813.1+572028	[HB89] 1637+574	1.81e-12	16 38 13.41 +57 20 24.2	1199.4	1.00	17.0	0.7506		LPQ
J163830.9-205520	IRAS 16355-2049	2.23e-11	16 38 31.15 -20 55 27.1	7.1	1.00	(14.5)			
J163845.7-141553	PKS 1635-14	6.03e-12	16 38 45.30 -14 15 50.3 ^{OR}	764.9 ^D	1.00	18.0			QSO
J164022.1+464215 [†]	MG4 J164018+4643	8.10e-12	16 40 23.74 +46 42 10.4 ^{RF}	239.9 ^E	1.00	17.0	0.2344		
J164220.4+221132		3.36e-12	16 42 20.34 +22 11 43.4	15.4	1.00	(18.9)			
J164238.9+272621 [†]	RX J1642.6+2726:001	4.63e-12	16 42 38.66 +27 26 39.0	8.7	1.00	(16.5)	0.1030 ²		Early
J164258.9+394822	3C 345	3.43e-12	16 42 58.77 +39 48 37.0	7099.1	1.00	16.0	0.5928	Opt.var.	HPQ
J164419.8+454635	MG4 J164418+4547	2.62e-12	16 44 20.05 +45 46 45.4	183.5	1.00	17.6	0.2230		BLLAC
J164443.2+261909	MG2 J164443+2618	2.94e-12	16 44 42.37 +26 19 13.5	128.9	1.00	17.1	0.1450		Sy1
J164610.3-112405	IRAS 16433-1118	9.76e-12	16 46 10.48 -11 24 05.6	40.2 ^E	1.00	15.9	0.0750		Sy1
J164613.2+693956	RX J1646.2+6939	3.01e-12	16 46 13.32 +69 39 58.4	3.5	0.98	16.6	0.2150		Sy1
J165110.2+045936	Hercules A	6.54e-12	16 51 08.10 +04 59 33.7 ^{OR}	43188.8 ^D	0.99	18.5	0.1540	Radio galaxy	
J165141.2+721824	RX J1651.6+7218	3.05e-12	16 51 39.95 +72 18 25.2	21.3	1.00	16.8			BLLAC
J165217.0+444922		3.08e-12	16 52 14.61 +44 49 23.3 ^{RF}	3.9	0.77	(17.4)			
			16 52 20.23 +44 49 23.6 ^{RF}	4.6	0.29	(16.5)			
J165253.7+400927	FIRST J165253.2+400912	1.62e-12	16 52 53.29 +40 09 13.3	58.9	1.00	(17.0)	0.1480 ²		BLLAC?
J165352.6+394538	MRK 0501	6.17e-11	16 53 52.24 +39 45 36.6	1558.5	1.00	14.1	0.0337	E?	BLLAC
J165358.1+441042	RX J1653.9+4410:001	2.50e-12	16 53 55.23 +44 10 22.9	37.7	1.00	18.1			
J165423.8+233322		2.94e-12	16 54 23.86 +23 33 57.6 ^F	29.7 ^E	0.99	(16.4)			
J165551.7+214559		6.98e-12	16 55 51.48 +21 46 02.4	6.9	1.00	(16.5)			
J165833.5+051521	[HB89] 1656+053	3.76e-12	16 58 33.47 +05 15 15.6	1534.3	1.00	16.5	0.8790		HPQ
J165945.3+323651	PGC 059392	3.62e-12	16 59 43.42 +32 36 50.8 ^R	40.0 ^E	0.99	16.5	0.0984		

Table 2—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position		S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	(J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J170242.5+340336 [†]	FIRST J170242.5+340337	1.65e-11	17 02 42.51	+34 03 37.3 ^{RF}	3.5	1.00	14.5	0.0984 ¹		Early
J170245.5+725330	UGC 10697	1.04e-11	17 02 44.14	+72 53 28.5	4.3	0.99	15.7	0.0530	S	Sy1
J170330.5+454040	IRAS 17020+4544	1.60e-11	17 03 30.34	+45 40 47.5	121.8	1.00	15.1	0.0604		Sy2
J170500.2-013225	UGC 10683 NOTES02	4.61e-12	17 05 00.83	-01 32 27.7	3.8 ^E	0.97	15.6	0.0308	SBa	Sy1
J170631.6-064645		6.21e-12	17 06 31.65	-06 46 36.3	7.8	1.00	(16.4)			
J170745.1+133102	MG1 J170746+1331	3.52e-12	17 07 45.67	+13 31 05.7	302.0	1.00	(17.2)			
J170817.7-034938		5.18e-12	17 08 17.48	-03 49 13.2 ^O	4.6 ^E	0.07	(17.5)			
			17 08 18.09	-03 49 26.2 ^O	4.6 ^E	0.87	(17.8)			
J170941.2+342529 [†]	4C +34.45	2.63e-12	17 09 39.03	+34 25 50.3	642.9 ^E	0.98	15.3	0.0808	E	
J171013.2+334410	FIRST J171013.4+334402	2.59e-12	17 10 13.43	+33 44 02.8 ^F	5.0	1.00	16.4	0.2080		Sy1
J171100.5+394128	NPM1G +39.0436	3.21e-12	17 10 56.34	+39 41 31.8 ^{RF}	497.2 ^S	0.73	16.5	0.0623		AGN
J171304.5+352335	FIRST J171304.4+352333	2.18e-11	17 13 04.37	+35 23 32.9	12.0	1.00	(16.2)	0.0850 ²		QSO/HII?
J171322.8+325631	87GB 171131.7+325957	3.91e-12	17 13 22.34	+32 56 33.3	73.3 ^E	1.00	15.4	0.1010 ²		Sy2
J171511.5+030941	PMN J1715+0309	6.77e-12	17 15 12.27	+03 09 39.0	163.3	1.00	(21.5)			
J171519.5+572430 [†]	NGC 6338	1.41e-11	17 15 22.94	+57 24 40.5	57.5	1.00	13.4	0.0280	cD;S0	
J171601.3+311215	IRAS F17141+3115	6.52e-12	17 16 02.63	+31 12 08.6	3.4	0.99	16.0	0.1110		Sy1
J171614.5+683636	[HB89] 1716+686	3.21e-12	17 16 13.94	+68 36 38.3	489.3	1.00	18.5	0.7770		QSO
J171618.7+202123		3.56e-12	17 16 18.50	+20 21 37.6 ^{OR}	74.3	0.25	(17.0)			
			17 16 19.14	+20 21 42.6 ^{OR}	74.3	0.74	(18.2)			
J171706.8+293117 [†]	RBS 1634:001	6.55e-12	17 17 06.46	+29 31 17.7	3.2	0.97	19.6	0.2750		
J171718.9+422652 [†]	LEDA 093954	3.46e-12	17 17 19.18	+42 26 59.6	134.0	1.00	14.8	0.1829		
J171746.9+194057 [†]		3.86e-12	17 17 45.55	+19 40 40.5 ^O	21.9 ^A	0.18	(17.2)			
	ABELL 2254:[CEF95]		17 17 45.88	+19 40 48.4 ^O	21.9 ^A	0.35	(17.3)	0.1780	cD	
			17 17 46.55	+19 40 57.0 ^O	21.9 ^A	0.08	(18.4)			
J171810.9+563955 [†]	7C 1717+5643	6.25e-12	17 18 11.92	+56 39 55.9	38.8	1.00	(16.5)	0.1135		
J171937.9+480409	PG 1718+481	2.11e-12	17 19 38.31	+48 04 12.2	63.7	1.00	15.3	1.0840		QSO
J172000.8+111500		3.93e-12	17 20 01.06	+11 15 03.3	3.6 ^E	0.95	(16.2)			
J172009.3+263727 [†]	RGB J1720+266	1.39e-11	17 20 09.89	+26 37 30.8 ^O	89.0 ^E	1.00	12.3			
J172015.5+353622	FIRST J172016.7+353625	2.97e-12	17 20 16.80	+35 36 28.4	18.8	1.00	(21.5)			
J172019.7+025328	PMN J1720+0253	3.46e-12	17 20 19.45	+02 53 29.1 ^O	334.0 ^D	0.17	(18.9)			
	PMN J1720+0253		17 20 19.57	+02 53 31.2 ^O	334.0 ^D	0.83	(17.0)			
J172202.0+431523		5.35e-12	17 22 01.83	+43 15 26.8	8.8	1.00	(18.7)			
J172226.7+320752 [†]	FIRST J172227.0+320758	8.53e-12	17 22 27.08	+32 07 58.0 ^F	7.2 ^E	1.00	19.2	0.2240	cD	
J172239.8+305252	MRK 0506	3.42e-11	17 22 40.10	+30 52 55.2 ^R	11.4 ^E	0.99	15.1	0.0430	SAB(r)a	Sy1.5
J172241.0+243625	[HB89] 1720+246	3.75e-12	17 22 41.74	+24 36 21.0	13.4	1.00	16.4	0.1750	Opt.var.	BLRG
J172320.3+235053		3.30e-12	17 23 21.61	+23 50 39.2 ^O	3.9	0.70	(17.4)			
			17 23 21.87	+23 50 46.3 ^O	3.9	0.19	(19.6)			
			17 23 22.56	+23 50 41.7 ^O	3.9	0.06	(19.2)			
J172320.5+341756	[HB89] 1721+343 NED02	2.17e-11	17 23 20.85	+34 17 57.3	1614.1 ^T	1.00	16.5	0.2060	Opt.var.	QSO
J172323.3+363010	IRAS 17216+3633	5.10e-12	17 23 23.89	+36 30 13.3	4.7	1.00	16.0	0.0400		Sy1
J172504.4+115218	87GB 172245.8+115500	3.30e-11	17 25 04.33	+11 52 15.5	120.8	1.00	15.8	0.0180		BLLAC
J172520.6+525508	RX J1725.3+5255	2.22e-12	17 25 20.11	+52 54 55.6	15.6 ^E	1.00	17.3			

Table 2—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position		S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	(J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J172658.4−031729	TXS 1724−032	1.01e−11	17 26 57.60	−03 17 37.9	188.3 ^E	0.96	(17.7)		sym. dbl.	
J172818.3+501310	[HB89] 1727+502 NED01	3.73e−11	17 28 18.58	+50 13 11.3	201.1	1.00	16.4	0.0554	Opt.var.	BLLAC
J172821.0+014137		4.28e−12	17 28 21.45	+01 41 29.8 ^O	6.4 ^E	0.35	(16.6)			
			17 28 21.25	+01 41 42.5 ^O	6.4 ^E	0.60	(16.5)			
J173113.6+144259		6.15e−12	17 31 13.46	+14 42 51.6	5.5	1.00	(16.1)			
J173114.5+323250	RGB J1731+325	4.77e−12	17 31 13.91	+32 32 43.8 ^{OR}	709.8 ^D	0.23	(16.5)			
			17 31 14.50	+32 32 48.5 ^{OR}	709.8 ^D	0.60	(16.8)			
			17 31 15.20	+32 32 59.4 ^{OR}	709.8 ^D	0.12	(16.1)			
J173140.1+225239		3.84e−12	17 31 39.87	+22 52 02.0	20.8 ^A	0.96	(19.0)			
J173257.0+403635 [†]	FIRST J173300.4+403643	3.14e−12	17 33 00.48	+40 36 43.3 ^F	37.1	0.68	(16.6)	0.1361		
J173301.7+434533 [†]	IC 1262	1.20e−11	17 33 02.07	+43 45 34.1	18.2	1.00	14.7	0.0326	cD;E	
J173416.0+134343		4.46e−12	17 34 15.20	+13 43 40.1 ^O	3.1	0.09	(19.2)			
			17 34 16.39	+13 43 51.5 ^O	3.1	0.90	(16.5)			
J173842.6+382103		2.54e−12	17 38 42.48	+38 21 04.6	13.3	1.00	(18.8)			
J174100.2+400510		2.19e−12	17 40 59.68	+40 05 06.8 ^{OR}	36.3 ^T	0.43	(18.2)			
			17 41 00.16	+40 05 25.1 ^{OR}	36.3 ^T	0.56	(19.6)			
J174128.1+034848	IRAS F17389+0350	3.25e−11	17 41 28.20	+03 48 52.4	14.2	0.95	15.3	0.0300		Sy1
J174207.1+182720	[HB89] 1739+184	6.55e−12	17 42 06.85	+18 27 29.9	1136.4 ^T	0.99	16.4	0.1860		Sy1
J174248.6+390027	PGC 060736	3.91e−12	17 42 46.36	+39 00 31.7 ^O	325.7 ^D	0.99	16.4			
J174357.3+193505	NPM1G +19.0510	1.02e−11	17 43 57.96	+19 35 08.6	565.5 ^T	1.00	16.8	0.0830		BLLAC
J174414.3+325925 [†]	PGC 060777	1.77e−11	17 44 14.46	+32 59 29.8	92.0	1.00	13.5	0.0757		
J174538.1+290823		1.48e−11	17 45 38.34	+29 08 21.5	13.3	1.00	(16.3)			
J174702.0+493803	RX J1747.0+4938	2.54e−12	17 47 02.46	+49 38 03.4	7.8	1.00	(19.5)			
J175004.4+470037 [†]	B3 1748+470	3.95e−12	17 50 05.00	+47 00 43.7 ^R	109.8 ^S	1.00	18.0	0.1600		BLLAC?
J175016.5+350452	87GB 174830.8+350607	4.91e−12	17 50 16.82	+35 04 58.7	69.5	1.00	11.6			
J175116.3+504537	IRAS 17500+5046	9.36e−12	17 51 16.60	+50 45 36.4	11.0	1.00	15.4	0.2997		Sy1
J175441.9+680334 [†]		4.08e−12	17 54 38.61	+68 03 17.3 ^O	4.8	0.28	(16.1)			
	VII Zw 751 NOTES02		17 54 38.14	+68 03 34.6 ^O	4.8	0.72	17.4			
J175607.5+545536		5.78e−12	17 56 08.70	+54 55 37.3	3.5	1.00	(15.5)			
J175615.5+552217	RGB J1756+553	1.40e−11	17 56 15.96	+55 22 17.8	16.8	1.00	17.3			BLLAC
J175706.9+535130 [†]	87GB 175603.7+535155	2.91e−12	17 57 06.74	+53 51 37.3	59.7	1.00	13.4	0.1190	cD	
J175712.8+703339	MS 1757.7+7034	9.18e−12	17 57 12.62	+70 33 37.8	10.8	1.00	18.3	0.4070		BLLAC
J175747.8+194650		1.23e−11	17 57 47.83	+19 46 57.3 ^O	3.0	0.62	(16.5)			
			17 57 47.86	+19 46 46.3 ^O	3.0	0.36	(19.7)			
J175949.4+703719		2.81e−12	17 59 49.01	+70 37 21.3	11.3	1.00	(21.5)			
J180208.6+183043	87GB 175956.0+183122	4.57e−12	18 02 07.51	+18 31 04.0 ^{OR}	101.1	0.68	(17.7)			
J180216.3+641546		3.46e−12	18 02 16.24	+64 15 42.9	2.8	1.00	(16.1)			
J180650.2+694923	3C 371	4.32e−12	18 06 50.46	+69 49 28.1	1887.1 ^E	1.00	14.4	0.0510	E	BLLAC
J181100.0+495430	NGC 6582 NED01	6.64e−12	18 11 01.93	+49 54 43.2	18.4	1.00	16.0	0.0481	E	
J181340.8+300143	IRAS F18117+3000	5.06e−12	18 13 41.20	+30 01 38.6 ^O	4.3	0.95	(16.6)			
	IRAS F18117+3000		18 13 41.99	+30 01 37.7 ^O	4.3	0.05	(16.5)			
J181709.7+394703		2.35e−12	18 17 10.20	+39 46 52.1	3.0	0.95	(17.0)			

Table 2—Continued

(1) <i>ROSAT</i> BSC 1RXS name	(2) NED Common name	(3) S_x ergs s ⁻¹ cm ⁻²	(4) Position α (J2000) δ	(5) $S_{1.4\text{GHz}}$ mJy	(6) ID P	(7) m_B	(8) Redshift	(9) NED Morphology	(10) Sp. Class.
J182042.7+381711		5.58e-12	18 20 42.48 +38 17 10.3 ^O	7.2	0.83	(16.9)			
			18 20 44.37 +38 17 08.3 ^O	7.2	0.10	(18.4)			
J182157.4+642051 [†]	[HB89] 1821+643	2.83e-11	18 21 57.22 +64 20 36.4 ^R	93.6 ^S	1.00	14.1	0.2970		Sy1
J182407.7+302933	NPM1G +30.0453	6.55e-12	18 24 06.88 +30 29 31.6	16.7	1.00	16.7			
J182418.7+430954	RGB J1824+431	4.69e-12	18 24 19.21 +43 09 49.6	33.6	1.00	19.1			
J182932.3+484447	[HB89] 1828+487	4.99e-12	18 29 31.80 +48 44 46.7	13753.0	1.00	16.8	0.6920	Opt.var.	LPQ
J183235.1+684805	87GB 183253.8+684546	5.17e-12	18 32 35.52 +68 48 07.2	150.3	1.00	15.7	0.2050	E	BLLAC
J183503.2+324145	CGCG 173-014	7.51e-11	18 35 03.43 +32 41 46.7 ^O	5490.4 ^D	1.00	15.5	0.0579		Sy1
J183849.4+480236	87GB 183734.8+475951	3.36e-12	18 38 49.22 +48 02 35.0	30.1	1.00	16.7			BLLAC
J184147.0+321838	RGB J1841+323	7.29e-12	18 41 47.17 +32 18 38.6	21.9	1.00	(17.5)			
J184209.5+794613	3C 390.3	1.23e-11	18 42 08.89 +79 46 16.7 ^{OR}	11559.2 ^D	1.00	14.4	0.0561	Opt. var.	Sy1
J184402.1+453332	PGC 062332	4.15e-12	18 44 02.44 +45 33 32.1	5597.7	1.00	15.7	0.0910		
J184642.2+842506		3.05e-12	18 46 52.43 +84 25 06.1	7.1	1.00	(16.6)			
J184716.9+732053		4.01e-12	18 47 14.01 +73 21 04.6 ^O	3.0	0.74	(16.6)			
			18 47 15.03 +73 20 43.2 ^O	3.0	0.25	(17.8)			
J184822.6+653700		4.39e-12	18 48 22.66 +65 37 02.3	9.5	1.00	(18.7)			
J184847.6+424536	RGB J1848+427	1.26e-11	18 48 47.17 +42 45 35.0	22.0	1.00	18.4			BLLAC
J185209.4+571133	8C 1851+571	4.42e-12	18 52 08.34 +57 11 42.8	53.3	1.00	15.5		E	
J190749.9-392341		5.38e-12	19 07 50.29 -39 23 27.2	3.0	0.92	(16.0)			
J191928.5-295808		9.10e-12	19 19 27.45 -29 57 52.0 ^{OR}	1690.5 ^E	0.06	(18.8)			
			19 19 27.45 -29 57 52.0 ^{OR}	1902.3 ^D	0.06	(18.8)			
			19 19 27.60 -29 58 04.1 ^{OR}	1902.3 ^D	0.71	(18.6)			
			19 19 27.45 -29 57 52.0 ^{OR}	1805.7 ^D	0.06	(18.8)			
			19 19 27.60 -29 58 04.1 ^{OR}	1805.7 ^D	0.12	(18.6)			
J192006.3-273428		8.28e-12	19 20 06.40 -27 34 50.1 ^O	7.8 ^E	0.06	(17.7)			
			19 20 06.35 -27 34 34.7 ^O	7.8 ^E	0.86	(14.9)			
J192450.8-291437	[HB89] 1921-293	7.00e-12	19 24 51.11 -29 14 31.0	13387.2	1.00	17.0	0.3520	Opt.var.	BLLAC/HPQ
J192649.5+615445	87GB 192614.4+614823	4.35e-12	19 26 49.95 +61 54 41.4	22.4	1.00	(16.9)			
J192719.7+653354	WN B1927+6527	4.48e-11	19 27 19.59 +65 33 54.0	40.3	1.00	15.4	0.0170	S	Sy2
J192728.0-384712		3.82e-12	19 27 25.61 -38 47 11.1 ^O	11.0 ^E	0.75	(17.4)			
			19 27 25.98 -38 47 02.9 ^O	11.0 ^E	0.13	(16.7)			
J192748.0+735757	[HB89] 1928+738	1.02e-11	19 27 48.53 +73 58 02.1	3950.9 ^E	1.00	16.5	0.3021		LPQ
J192833.5+533900		4.98e-12	19 28 33.82 +53 39 05.9	4.5 ^E	1.00	(19.7)			
J193001.6-150929	PMN J1930-1509	1.27e-11	19 30 01.79 -15 09 15.3 ^R	1914.5 ^D	1.00	(16.7)		asym. dbl.	
J193113.8+600001	87GB 193024.4+595349	4.66e-12	19 31 12.76 +60 00 12.3 ^R	81.6 ^S	0.98	(16.3)			
J193139.0-335426	PGC 063333	9.30e-12	19 31 38.53 -33 54 43.2	885.6 ^D	1.00	17.0	0.0980	Elliptical	
J193150.1-263436		6.46e-12	19 31 49.46 -26 34 48.4 ^R	102.1 ^E	0.97	(18.0)			
J193412.6-241923		9.81e-12	19 34 12.84 -24 19 22.4	19.0	1.00	(16.5)			
J193708.3+660821	WN B1936+6601	5.59e-12	19 37 08.25 +66 08 21.3 ^O	15.8 ^E	0.80	(16.2)			
J193818.5+540928		9.33e-12	19 38 17.74 +54 09 51.9	17.0 ^E	0.97	(21.5)			
J194015.0-301552	IRAS 19370-3022	2.40e-11	19 40 15.35 -30 15 56.1	9.9	0.95	(13.8)			
J194241.3-101928	NGC 6814	1.67e-11	19 42 40.52 -10 19 19.7	51.9 ^E	1.00	12.1	0.0052	SAB(rs)bc	Sy1.5

Table 2—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position		S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	(J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J194306.8–351001		1.15e–11	19 43 06.59	–35 10 06.5	67.0	1.00	(16.6)			
J194423.1+655216	4C +65.28	4.19e–12	19 44 24.00	+65 52 23.3	141.6	1.00	(19.1)		sym. dbl.	
J194455.3–214318		3.68e–12	19 44 55.16	–21 43 20.7	42.1	1.00	(18.1)			
J194740.7–054157	CGMW 3–4849	6.81e–12	19 47 40.79	–05 42 16.7	49.4	0.99	(15.9)		E	
J195352.4+624810		5.51e–12	19 53 52.44	+62 48 07.7	4.5	1.00	(15.3)			
J195500.6–160328		4.14e–12	19 55 00.58	–16 03 37.9	24.7	1.00	(17.8)			
J195815.6–301119		1.70e–11	19 58 14.89	–30 11 12.8	128.4 ^S	0.98	(16.3)			
J200000.0+650854	87GB 195929.2+650039	1.09e–10	19 59 59.88	+65 08 54.8	250.1	1.00	14.7	0.0470	E	BLLAC
J200057.3–174856	[HB89] 1958–179	4.06e–12	20 00 57.10	–17 48 57.5	550.5	1.00	17.5	0.6500		HPQ
J200354.5–085707		5.79e–12	20 03 54.73	–08 56 44.6	9.3	0.98	(16.5)	0.0572 ¹		Early
J200658.1–343301	ESO 399–IG 020 NED02	1.32e–11	20 06 57.90	–34 32 55.3	8.8	1.00	17.0	0.0250		Sy1
J201020.0–252356		4.60e–12	20 10 19.79	–25 23 58.8	58.9	1.00	(19.2)			
J201116.7–154631	[HB89] 2008–159	4.10e–12	20 11 15.72	–15 46 40.3	546.5	1.00	17.2	1.1800		QSO
J201449.9–243043	PMN J2014–2430	2.22e–11	20 14 51.75	–24 30 22.7	229.5	1.00	(16.3)			
J202253.6–205642		6.51e–12	20 22 49.30	–20 57 16.5 ^{OR}	6.1	0.15	(17.3)			
	ESO 596– G 024		20 22 59.47	–20 56 57.5 ^{OR}	229.3 ^E	0.74	15.6	0.0540	S...	
J202428.9–084810		5.21e–12	20 24 29.38	–08 48 04.7	37.0	1.00	(18.4)			
J202702.9–303109		5.93e–12	20 27 02.01	–30 30 59.9 ^O	3.8	0.23	(16.6)			
	MCG –05–48–013		20 27 02.72	–30 31 11.6 ^O	3.8	0.77	15.5			
J202829.1–332520		3.05e–12	20 28 26.90	–33 25 34.7 ^O	19.2 ^E	0.24	(18.5)			
			20 28 27.70	–33 25 39.5 ^O	19.2 ^E	0.73	(16.0)			
J203252.6–025905		3.88e–12	20 32 51.90	–02 59 16.0	4.5	1.00	(16.5)			
J203433.1–303731	PGC 064989	5.02e–12	20 34 31.49	–30 37 26.6 ^R	3.4	0.94	(13.3)	0.0190		Sy1
J203442.0–340427		9.03e–12	20 34 41.43	–34 03 56.2	41.7	0.99	(15.3)			
J203445.3–354921 [†]	ESO 400–IG 040 NED01	1.09e–11	20 34 44.47	–35 48 51.9	1717.9	0.94	17.0	0.0884	E1+	
J203650.9–332817		3.96e–12	20 36 49.49	–33 28 30.8 ^O	3.3	0.72	(18.9)			
			20 36 50.09	–33 28 19.2 ^O	3.3	0.24	(20.4)			
J203828.4–201059	PMN J2038–2011	2.87e–12	20 38 27.78	–20 11 07.9 ^{OR}	760.5 ^D	1.00	16.4	0.5200		QSO
J203845.1–263625		4.42e–12	20 38 44.98	–26 36 31.7	5.5	1.00	(19.4)			
J203910.6–195848	PMN J2039–1958	3.58e–12	20 39 10.97	–19 58 51.0 ^{OR}	309.1 ^D	1.00	(16.7)		asym. dbl.	
J203927.2–301844	RBS 1688	1.26e–11	20 39 27.24	–30 18 51.2	5.9	0.99	16.0	0.0800	NLS1	Sy1
J204149.8–373346		9.25e–12	20 41 50.52	–37 33 41.3	88.4 ^E	1.00	(16.1)			
J204235.3+750758	4C +74.26	2.57e–11	20 42 37.93	+75 08 02.1	1323.0 ^T	1.00	15.1	0.1040		QSO
J204314.1–214417	PMN J2043–2144	4.45e–12	20 43 14.66	–21 44 33.1	295.4	1.00	(17.2)	0.2040		
J204402.5–075812		3.15e–12	20 44 02.01	–07 58 09.6 ^O	5.8 ^E	0.32	(16.4)			
			20 44 02.76	–07 58 19.7 ^O	5.8 ^E	0.68	(16.3)			
J204409.9–104330	MRK 0509	9.80e–11	20 44 09.68	–10 43 23.0	19.2	1.00	13.0	0.0344	Compact	Sy1.2
J204737.0–290901		4.12e–12	20 47 36.05	–29 08 57.3	10.6	1.00	(18.7)			
J204745.9–024609	PMN J2047–0246	8.17e–12	20 47 45.66	–02 46 05.0	73.3	1.00	(16.3)			
J204921.6–003930		5.27e–12	20 49 21.83	–00 39 32.8	5.8	1.00	(18.5)			
J205242.6+081039	RX J2052.7+0810	5.37e–12	20 52 43.13	+08 10 36.0 ^O	6.0	0.28	(19.0)			
	RX J2052.7+0810		20 52 42.70	+08 10 38.2 ^O	6.0	0.72	(21.5)			

Table 2—Continued

(1) <i>ROSAT</i> BSC 1RXS name	(2) NED Common name	(3) S_x ergs s ⁻¹ cm ⁻²	(4) Position α (J2000) δ	(5) $S_{1.4\text{GHz}}$ mJy	(6) ID P	(7) m_B	(8) Redshift	(9) NED Morphology	(10) Sp. Class.
J205456.9+001533	PMN J2054+0016	3.71e-12	20 54 56.83 +00 15 38.1	57.0	1.00	(17.7)			
J205528.2-002123		1.33e-11	20 55 28.23 -00 21 15.8	40.9	1.00	(18.3)			
J205941.5-385847	IRAS F20564-3910	3.22e-12	20 59 40.65 -38 58 30.4 ^R	3.1 ^E	0.95	15.7			
J210027.7-170913 [†]	PMN J2100-1708	3.45e-12	21 00 27.76 -17 09 08.7 ^{OR}	307.4 ^E	—	(16.5)	0.2436 ¹		Sy1
J210210.4-243202	QW 2059-247	6.14e-12	21 02 09.89 -24 31 59.4	69.4	1.00	(16.5)	0.1899		
J210221.7+105812	CGCG 425-034	7.07e-12	21 02 21.43 +10 58 18.3	6.9	1.00	15.4	0.0287 ¹		Sy1.5
J210322.2-214743	TXS 2100-219	3.48e-12	21 03 22.19 -21 47 56.9	234.5 ^T	1.00	(16.7)		sym. dbl.	
J210352.6-204701	PMN J2103-2046	3.20e-12	21 03 52.72 -20 46 51.1	38.6	1.00	(16.6)			
J210416.4+211816		7.14e-12	21 04 15.87 +21 18 05.2	14.6	1.00	(18.7)			
J210707.7-252643 [†]		4.99e-12	21 07 08.14 -25 26 22.3 ^R	14.3	1.00	(16.7)			
J210910.1-094011	H 2106-099	5.18e-11	21 09 09.82 -09 40 09.8	5.9	1.00	14.3	0.0268		Sy1.2/LINER
J210928.6-381310		2.92e-12	21 09 29.82 -38 13 13.8 ^O	2.7	0.08	(18.2)			
	AM 2106-382		21 09 28.72 -38 13 11.3 ^O	2.7	0.86	15.3	0.0465	Sa	LINER
J211242.5+081831		7.29e-12	21 12 42.98 +08 18 36.2	23.4	1.00	(21.5)			
J211400.0+820447	IRAS F21166+8152	1.62e-11	21 14 00.33 +82 04 52.0	482.1 ^T	1.00	15.7	0.0840		Sy1
J211843.5-063617	PMN J2118-0636	3.61e-12	21 18 43.18 -06 36 19.1	89.3	1.00	(17.0)	0.3280 ²		Sy1
J211926.1+152105		4.76e-12	21 19 26.43 +15 21 07.0	3.3	1.00	(19.1)			
J212307.4-103650	PMN J2123-1036	1.37e-11	21 23 07.24 -10 36 49.0	119.1	1.00	(1756)	0.0227 ¹		Early?
J212516.7-255530		2.67e-12	21 25 16.19 -25 55 30.6	5.0	1.00	(18.8)			
J212522.6+174250		3.53e-12	21 25 21.97 +17 43 06.4	16.7	0.99	(18.2)			
J212706.7-120927	MRC 2124-123	5.90E-12	21 27 09.86 -12 09 58.8	265.0 ^D	—	19.2			
J212912.9-153846	[HB89] 2126-158	3.50e-12	21 29 12.23 -15 38 41.5	590.1	1.00	17.3	3.2680	GPS	QSO
J212940.3+000544	[CGH98] J2129+0005	8.96e-12	21 29 39.99 +00 05 23.0	26.2	1.00	(18.3)			
J213103.5-274702		5.01e-12	21 31 03.29 -27 46 58.0	37.3	1.00	(17.6)			
J213135.5-091525		1.53e-11	21 31 35.45 -09 15 22.7	43.5	1.00	(17.0)			
J213135.7-120719	[HB89] 2128-123	6.11e-12	21 31 35.26 -12 07 06.2	1770.0	1.00	15.5	0.5010	Opt.var	Sy1/LPQ
J213151.7-251602		5.67e-12	21 31 51.50 -25 15 58.5	10.9	1.00	(18.8)			
J213202.3-334255		4.44e-11	21 32 02.50 -33 43 01.9 ^O	3.5	0.57	(17.6)			
	CTS 0109		21 32 02.23 -33 42 54.0 ^O	3.5	0.40	15.7			Sy1
J213228.1+100812		3.31e-11	21 32 27.33 +10 08 15.0 ^O	6.5	0.13	(18.3)			
			21 32 28.33 +10 08 23.9 ^O	6.5	0.11	(18.4)			
	UGC 11763		21 32 27.87 +10 08 19.8 ^O	6.5	0.76	14.6	0.0621		Sy1
J213231.0-262611		2.76e-12	21 32 32.09 -26 25 46.9 ^{OR}	66.5 ^D	1.00	(17.2)			
J213414.4-311737 [†]		3.10e-12	21 34 14.36 -31 17 41.5 ^O	32.1 ^E	—	(18.5)			
J213445.2-272551		2.82e-12	21 34 44.66 -27 25 39.9 ^O	9.3	0.61	(16.2)			
	IRAS F21318-2739		21 34 45.14 -27 25 55.5 ^O	9.3	0.34	16.4		S0	
J213504.7+085807 [†]		3.08e-12	21 35 04.87 +08 57 52.2	3.1	0.95	(15.3)			
J213745.3-143250	[HB89] 2135-147	1.32e-11	21 37 45.21 -14 32 55.6 ^O	3866.2 ^D	1.00	15.9	0.2003	E1;opt.var.	Sy1
J213833.0+320507	CGCG 493-002	8.56e-12	21 38 33.32 +32 05 03.4	7.0	1.00	15.6	0.0248 ¹		Sy1.5
J213852.9-205354		1.19e-11	21 38 52.77 -20 53 46.5	9.8	1.00	(18.1)			
J214015.3-233946 [†]	NGC 7099:[JVH94] 12	3.86e-12	21 40 14.99 -23 39 39.5	4.0	1.00	18.5	0.3130		
J214237.3-043757	[HB89] 2140-048	3.26e-12	21 42 36.86 -04 37 43.0	806.1	1.00	18.0	0.3440		QSO

Table 2—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position		S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	(J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J214533.6−043434	NPM1G −04.0578	3.74e−12	21 45 33.34	−04 34 39.0	68.0	1.00	16.2	0.0697 ¹		Early
J214552.3+071923	PMN J2145+0720	1.06e−11	21 45 52.30	+07 19 26.7	101.9	1.00	18.0	0.2370		BLLAC
J214636.3−305132	DUKST 466−078	1.48e−11	21 46 35.47	−30 51 37.1	2.8	1.00	17.0			
J214703.0−101931		4.30e−12	21 47 00.41	−10 19 12.0	20.0	1.00	(16.1)	0.0797 ¹		Early
J214805.4+065737	[HB89] 2145+067	3.29e−12	21 48 05.48	+06 57 39.4	2589.9	1.00	16.5	0.9900	Opt.var.	LPQ
J214906.4−304157	ABELL 3814:001	6.67e−12	21 49 07.43	−30 42 04.6	31.8	1.00	17.2	0.1177	cD?	
J214958.2−185926	PMN J2149−1859	1.14e−11	21 49 58.06	−18 59 23.8 ^{OR}	419.7 ^D	1.00	(16.2)			
J215015.6−141047	RBS 1787	1.51e−11	21 50 15.82	−14 10 47.1	68.3	1.00	18.5	0.2290		BLLAC
J215101.0−073546		3.04e−12	21 51 00.77	−07 36 31.2	8.0	0.97	(21.5)			
J215155.4−302754	[HB89] 2149−307	7.17e−12	21 51 55.54	−30 27 53.6	1243.6	1.00	18.4	2.3450		QSO
J215213.9−194256	ESO 600− G 014	3.02e−12	21 52 09.59	−19 43 23.7 ^O	56.2	0.55	16.2		S...	
J215214.4−120544		5.35e−12	21 52 14.10	−12 05 41.5	53.4	1.00	(18.2)			
J215305.2−004227		7.42e−12	21 53 05.41	−00 42 31.4	24.7	1.00	(18.3)	− ²		BLLAC?
J215336.1+174111 [†]	ABELL 2390:341	1.58e−11	21 53 36.81	+17 41 44.8	236.0 ^E	1.00	19.6	0.2302	Sb	
J215541.5+123144 [†]	4C +12.76 NED02	5.25e−12	21 55 42.23	+12 31 29.4	494.9	1.00	(17.1)		VLA−reslvdcmpt	
J215602.5+181839	RX J2156.0+1818	5.04e−12	21 56 01.63	+18 18 39.2	33.2	1.00	17.4			QSO
J215623.8+331829		2.09e−11	21 56 23.18	+33 18 37.9	161.6	1.00	16.2	0.0784 ¹		Early
J215816.5−200617	APMBGC 601+092+014	2.42e−12	21 58 22.45	−20 06 12.9	35.3	0.78	15.5	0.0578	E	
J215830.1−094759	ABELL 2402:[OWB92]	6.36e−12	21 58 28.96	−09 47 48.1	32.3	1.00	13.8	0.0803		
J215852.2−301338	[HB89] 2155−304	5.27e−10	21 58 52.12	−30 13 30.2	525.4 ^T	1.00	14.0	0.1160	Opt.var.	BLLAC
J220040.1−230731	PMN J2200−2307	2.29e−12	22 00 40.30	−23 07 23.2	170.7	1.00	(16.9)			
J220156.0−170656		4.33e−12	22 01 55.73	−17 06 52.9	4.6	1.00	(16.7)	0.1686 ¹		Early?
J220246.0−130458		2.94e−12	22 02 44.62	−13 04 46.1 ^O	2.9	0.08	(16.3)			
			22 02 45.17	−13 04 42.3 ^O	2.9	0.23	(17.2)			
			22 02 45.50	−13 04 39.1 ^O	2.9	0.06	(18.5)			
	GSC 5808 00770		22 02 45.12	−13 04 52.6 ^O	2.9	0.62	15.0	0.0389 ¹		Sy1.9
J220315.6+314535	[HB89] 2201+315	1.08e−11	22 03 14.92	+31 45 38.7	2879.2 ^E	1.00	15.5	0.2950		LPQ
J220408.6−085710		3.73e−12	22 04 07.54	−08 57 11.2	2.8	0.99	(16.1)	0.1154 ¹		Sy1.8
J220610.3−300512	RX J2206.1−3005	1.48e−12	22 06 10.51	−30 05 10.4	11.4	0.99	19.1	0.1650		Sy1
J220907.9−274839	NGC 7214	1.01e−11	22 09 07.94	−27 48 31.1	29.2	1.00	13.1	0.0231	SB(s)bc pec:	Sy1.2
J221020.8−121040 [†]	ABELL 2420:[B90]	2.15e−11	22 10 18.64	−12 10 14.7	20.2	0.97	13.8		cD	
J221108.1−000315		3.44e−12	22 11 08.22	−00 03 01.2	25.5	1.00	(17.2)	− ²		BLLAC
J221124.2−132801	[HB89] 2208−137	3.23e−12	22 11 24.18	−13 28 10.1	1330.3	1.00	17.0	0.3920		BLLAC/HPQ
J221150.5−370714	CTS 0377	1.16e−12	22 11 50.57	−37 07 05.9	186.9	1.00	16.1	0.3200		QSO
J221154.5+184157	PG 2209+184	6.97e−12	22 11 53.90	+18 41 50.3	81.8	1.00	15.9	0.0700		Sy1
J221300.5−171015		7.87e−12	22 13 00.26	−17 10 17.9 ^O	3.3 ^E	0.91	(16.7)	0.1460 ¹		Sy2
			22 13 00.91	−17 10 09.1 ^O	3.3 ^E	0.09	(19.3)			
J221306.6−275405	PMN J2213−2754	2.97e−12	22 13 05.99	−27 54 20.0	76.1	1.00	(15.1)			
J221413.9−054223		3.24e−12	22 14 12.93	−05 42 11.3	7.7 ^E	0.92	(18.2)			
J221426.3−170124	PGC 068361	2.47e−12	22 14 25.71	−17 01 36.2 ^O	8915.4 ^D	1.00	17.8	0.1530		
J221442.4−384827		5.53e−12	22 14 42.98	−38 48 35.0 ^O	3.8 ^E	0.07	(16.4)			
	ESO 344− G 016		22 14 42.10	−38 48 22.3 ^O	3.8 ^E	0.92	15.1	0.0397	Sb	Sy1.5

Table 2—Continued

(1) <i>ROSAT</i> BSC 1RXS name	(2) NED Common name	(3) S_x ergs s ⁻¹ cm ⁻²	(4) Position α (J2000) δ	(5) $S_{1.4\text{GHz}}$ mJy	(6) ID P	(7) m_B	(8) Redshift	(9) NED Morphology	(10) Sp. Class.
J221513.4+171524		3.27e-12	22 15 13.49 +17 15 22.0	3.3	0.99	(16.9)			
J221537.8+290239		3.80e-12	22 15 36.66 +29 02 29.0 ^{OR}	603.5 ^T	0.20	(19.5)			
	4C +28.53		22 15 36.83 +29 02 36.2 ^{OR}	603.5 ^T	0.80	16.5	0.2288		QSO
J221601.2-280339		1.66e-12	22 16 02.74 -28 03 29.8 ^R	417.5 ^D	0.99	(21.5)			
J221614.9-092033 [†]	NPM1G -09.0687	8.45e-12	22 16 15.66 -09 20 00.8	12.6	—	16.7	0.0846		
J221656.7-172518		4.28e-12	22 16 57.46 -17 25 28.3 ^O	14.3 ^E	0.53	(18.3)			
			22 16 57.95 -17 25 18.4 ^O	14.3 ^E	0.34	(18.5)			
			22 16 58.07 -17 25 20.5 ^O	14.3 ^E	0.07	(20.0)			
			22 16 56.85 -17 25 25.6 ^O	14.3 ^E	0.05	(16.2)			
J221729.2-310619	RX J2217.4-3106	4.68e-12	22 17 28.27 -31 06 19.4	27.7	1.00	19.5			BLLAC
J221838.4-385342	LCRS B221542.3-390904	6.01e-12	22 18 39.36 -38 54 00.7 ^O	187.0 ^E	0.54	15.2			
J221839.1+080216	RX J2218.6+0802	3.77e-12	22 18 39.07 +08 02 13.3	4.0	1.00	16.6	0.1200		Sy1
J222033.5-350955		7.13e-12	22 20 32.65 -35 09 48.4 ^{OR}	9.3 ^D	0.33	(16.2)			
			22 20 32.97 -35 09 49.9 ^{OR}	9.3 ^D	0.66	(16.5)			
J222253.9-175318		2.81e-12	22 22 54.16 -17 53 24.0	5.6 ^E	0.98	(18.9)			
J222345.4+115012		6.53e-12	22 23 45.92 +11 50 11.5 ^O	3.7	0.21	(19.4)			
	MCG +02-57-002		22 23 45.01 +11 50 09.5 ^O	3.7	0.78	16.0	0.0297 ¹		Sy1.9
J222538.6+211806	[HB89] 2223+210	3.52e-12	22 25 38.03 +21 18 06.6	1837.6	1.00	18.2	1.9590		QSO
J222602.8+172245	ABELL 2443:[SPS89] 09	5.08e-12	22 26 02.34 +17 22 54.1 ^R	208.0 ^D	1.00	16.2	0.1096 ¹		Early
J222657.0+254946	87GB 222433.6+253457	3.07e-12	22 26 56.09 +25 50 09.6	77.2	1.00	(18.5)	0.1678 ¹		Early
J222846.3+333500		4.66e-12	22 28 46.62 +33 35 01.6	3.6	1.00	(16.4)	0.0897 ¹		Sy1
J222944.5-275538		2.35e-12	22 29 45.42 -27 55 34.7	7.6	1.00	(17.7)			
J223039.2-394246	[HB89] 2227-399	6.34e-12	22 30 40.36 -39 42 49.5	584.1 ^T	1.00	18.5	0.3230	Opt.var.	QSO
J223249.1-052954		3.39e-12	22 32 49.13 -05 29 58.1 ^{OR}	221.2 ^D	1.00	(17.9)			
J223249.5-202232		2.68e-12	22 32 48.80 -20 22 26.3	12.4	0.98	(18.0)			
J223301.0+133554	RX J2233.0+1335	3.47e-12	22 33 01.14 +13 36 01.6	25.1	1.00	19.4			BLLAC
J223546.0-260308	NGC 7314	3.09e-12	22 35 46.43 -26 03 01.7	33.9	1.00	11.9	0.0047	SAB(rs)bc	Sy1.9
J223647.3-123228	MRK 0915	3.56e-12	22 36 46.54 -12 32 43.1	70.6	1.00	14.8	0.0241	Sb	Sy1
J223656.5-221321	ESO 602- G 031	7.10e-12	22 36 56.02 -22 13 16.1	7.1	1.00	14.9	0.0337	(R'/SAB(rs)b	Sy1.8
J223812.7-394019	ESP 16713	5.00e-12	22 38 12.68 -39 40 18.5	20.3	1.00	19.2			
J224017.7+080316	UGC 12138	1.64e-11	22 40 17.03 +08 03 15.1	15.6	1.00	14.2	0.0246	SBa	Sy1.8
J224239.6+294333	UGC 12163	1.30e-10	22 42 39.40 +29 43 33.6	29.1	1.00	14.6	0.0240	SB	Sy1
J224329.6-303453		1.23e-12	22 43 27.66 -30 34 56.6 ^O	3.4	0.91	(19.0)			
J224341.9-123106		8.66e-12	22 43 40.50 -12 31 01.2	10.6	1.00	(18.2)			
J224505.2+263758		4.52e-12	22 45 04.72 +26 38 04.0	6.3	1.00	(19.3)			
J224621.7+314206	MG3 J224621+3142	5.28e-12	22 46 21.68 +31 42 08.0 ^O	361.3 ^D	0.75	15.0	0.1470		Sy1
			22 46 23.01 +31 42 06.4 ^O	361.3 ^D	0.25	(17.2)			
J224634.4-285400		1.77e-12	22 46 33.82 -28 53 40.2	8.6	1.00	(17.1)			
J224713.6+020410		3.45e-12	22 47 14.07 +02 04 09.2 ^{OR}	5.6	0.22	(16.9)			
			22 47 14.60 +02 04 08.3 ^{OR}	5.6	0.47	(16.1)			
			22 47 15.02 +02 04 15.4 ^{OR}	5.6	0.25	(18.3)			
			22 47 15.10 +02 03 54.9 ^{OR}	5.6	0.06	(16.4)			

Table 2—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position		S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	(J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J224805.1–172814	IRAS 22453–1744	3.59e–12	22 48 04.27	–17 28 30.7	16.0	1.00	(15.9)	0.1173		Sy1.8
J224818.7–101557		3.32e–12	22 48 17.54	–10 15 46.3	19.2	1.00	(17.0)	0.2910 ²		Sy1
J224911.1–130005		9.83e–12	22 49 10.84	–12 59 57.1	7.4	1.00	(17.7)			
J224919.5–372724 [†]	ESO 346– G 003	1.80e–12	22 49 22.84	–37 28 16.9	341.5	0.99	13.7	0.0287	E+2 pec:	
J225005.8+382439	MG3 J225005+3825	8.24e–12	22 50 05.86	+38 24 37.2	103.9	1.00	14.8	0.1184 ¹		Early
J225019.2+105406 [†]	MCG +02–58–021	1.28e–11	22 50 19.68	+10 54 11.0	15.2	1.00	16.0	0.0800		
J225119.5–382710	LCRS B224829.3–384303	4.39e–12	22 51 19.05	–38 27 07.7	59.7	1.00	16.6	0.1348		
J225146.9–320614		3.61e–12	22 51 47.55	–32 06 16.9	3.4	1.00	(19.5)			
J225207.7+145448	IRAS F22496+1438	4.53e–12	22 52 07.90	+14 54 40.9	2.5	1.00	16.3	0.1300		QSO
J225209.6+264236	IRAS 22497+2626	4.07e–12	22 52 09.36	+26 42 39.7 ^R	2.5	0.99	18.2	0.0670 ¹		HII
J225334.1–334309 [†]	ABELL 3934:001	2.30e–12	22 53 32.62	–33 43 05.4	34.5	1.00	17.4	0.2240		
J225358.0+160855	[HB89] 2251+158	2.06e–11	22 53 57.70	+16 08 53.4	12657.2	1.00	16.1	0.8590		Blazar/HPQ
J225406.7–173451	[HB89] 2251–178	2.03e–11	22 54 05.98	–17 34 55.4	16.6	1.00	14.4	0.0680	Opt.var.	Sy1
J225612.8–330341		2.41e–12	22 56 13.36	–33 03 37.8	21.4	1.00	(18.1)			
J225709.7–362740	IC 1459	1.31e–12	22 57 10.67	–36 27 43.0	1280.6 ^E	1.00	11.0	0.0056	E3	
J225809.9–011510		3.95e–12	22 58 10.02	–01 15 13.1 ^F	3.0	1.00	(16.3)	0.1167 ¹		Sy1
J225818.8–070704		2.72e–12	22 58 18.59	–07 07 08.3	6.0	1.00	(15.8)	0.2150 ¹		Sy1/QSO
J225834.4–094945	PMN J2258–0949	3.94e–12	22 58 33.99	–09 49 29.6	248.6	1.00	(21.5)		sym. dbl.	
J225845.8–260917		1.09e–11	22 58 45.60	–26 09 29.7	2.9	0.98	16.1	0.0760		Sy1
J225932.9+245505	PGC 070195	1.57e–11	22 59 33.05	+24 55 07.7	4.0	1.00	15.9	0.0345	Sa	Sy1
J230012.3+405225		6.02e–12	23 00 12.49	+40 52 24.3	14.9	1.00	(18.9)			
J230048.1–125518	NGC 7450	8.22e–12	23 00 47.58	–12 55 07.7	5.8	1.00	14.0	0.0106	(R)SB(r)a	Sy1.5
J230303.7–184131		9.04e–12	23 03 02.66	–18 41 23.4 ^{OR}	5967.6 ^E	0.36	(17.9)			
	PGC 070334		23 03 02.95	–18 41 26.0 ^{OR}	5967.6 ^E	0.64	18.1	0.1283	N galaxy	Sy1
J230315.7+085226	NGC 7469	5.68e–11	23 03 15.61	+08 52 26.3	181.0	1.00	13.0	0.0163	(R')SAB(rs)a	Sy1.2
J230325.2+075642	TXS 2300+076	4.72e–12	23 03 25.06	+07 56 42.4	41.8	1.00	19.1			
J230402.8+223725	MRK 0315	4.16e–12	23 04 02.61	+22 37 25.5	22.8	1.00	14.8	0.0389	E1 pec?	Sy1.5
J230437.1+370506		1.94e–11	23 04 36.63	+37 05 07.3	22.9	1.00	(17.7)			
J230443.8–084106	MRK 0926	4.03e–11	23 04 43.54	–08 41 07.6	34.5	1.00	15.0	0.0473		Sy1.5
J230636.0–110350		4.23e–12	23 06 35.05	–11 03 49.3	11.3	1.00	(21.5)			
J230654.5–343909		1.44e–12	23 06 53.83	–34 39 11.4 ^R	59.0 ^T	1.00	(17.7)			
J230714.9–151322 [†]	ABELL 2533:[QR95] cD	4.72e–12	23 07 14.01	–15 13 31.5	6.6	0.99	(16.2)	0.1110		
J230722.5–120520		2.44e–12	23 07 22.05	–12 05 18.6	7.1	1.00	(19.1)			
J230840.4–333201	IRAS F23059–3348	3.01e–12	23 08 39.40	–33 32 08.4	2.7	1.00	(14.8)			
J230846.7–221955	MS 2306.1–2236	6.96e–12	23 08 46.91	–22 19 45.5	6.4	1.00	17.5	0.1370		BLLAC
J230919.3–323003		5.48e–12	23 09 19.18	–32 30 16.0 ^O	3.0	0.11	(21.5)			
			23 09 19.20	–32 29 57.4 ^O	3.0	0.87	(15.6)			
J230940.6–363241		2.50e–12	23 09 41.04	–36 32 37.9 ^{OR}	90.8 ^E	0.18	(20.4)			
			23 09 40.03	–36 32 21.8 ^{OR}	90.8 ^E	0.67	(20.7)			
J231027.0–371926		1.90e–12	23 10 27.68	–37 19 12.4 ^O	6.2	0.27	(18.3)			
			23 10 28.05	–37 19 09.8 ^O	6.2	0.73	(17.9)			
J231118.6+100818	PG 2308+098	5.42e–12	23 11 17.73	+10 08 16.2 ^{OR}	579.7 ^D	0.92	16.0	0.4333		Sy1

Table 2—Continued

(1) <i>ROSAT</i> BSC 1RXS name	(2) NED Common name	(3) S_x ergs s ⁻¹ cm ⁻²	(4) Position α (J2000) δ	(5) $S_{1.4\text{GHz}}$ mJy	(6) ID P	(7) m_B	(8) Redshift	(9) NED Morphology	(10) Sp. Class.
	4C +09.72:[KSS94] 11		23 11 17.40 +10 08 22.6 ^{OR}	579.7 ^D	0.08	20.3			
J231235.4–340426	PMN J2312–3403	8.13e–12	23 12 34.14 –34 04 14.2 ^R	135.8	1.00	(16.3)			BLLAC
J231309.7–315816	[HB89] 2310–322	4.66e–12	23 13 10.48 –31 57 53.3 ^O	817.4 ^D	1.00	16.6	0.3370		QSO
J231341.0+140113	NGC 7525	2.95e–12	23 13 40.61 +14 01 13.6	7.5	1.00	15.2	0.0409	E	Sy1.5
J231357.3+144424	RGB J2313+147	3.30e–12	23 13 57.27 +14 44 22.7	41.0	1.00	15.2			AGN
J231456.0+224325	IRAS F23124+2226	3.53e–12	23 14 55.94 +22 43 22.4	18.7	1.00	16.3	0.1692		AGN
J231546.4–022246	NGC 7556	5.70e–12	23 15 43.94 –02 23 00.8	3.8	0.95	13.8	0.0250	S0 [–]	
J231607.5–202739 [†]	ABELL 2566:[OWG93]	8.07e–12	23 16 04.95 –20 27 48.2	26.4	0.95	15.2	0.0871		
J231712.8+184237 [†]	NGC 7578B	8.56e–12	23 17 13.63 +18 42 29.8	30.1	1.00	15.0	0.0402	E1:	
J231829.5+184246 [†]	NGC 7597	7.19E–12	23 18 29.96 +18 41 26.7	5.4	–	14.9	0.0369	cD;S?	
J231839.4+425745	UGC 12491	1.10e–11	23 18 39.20 +42 57 19.1	3.8 ^E	0.97	14.9	0.0169 ¹		Early
J231856.8+001436	NGC 7603	1.06e–11	23 18 56.68 +00 14 37.2	24.9	1.00	14.0	0.0295	SA(rs)b: pec	Sy1.5
J231947.4+425117	NGC 7618	5.53e–12	23 19 47.95 +42 51 10.8	40.0 ^E	1.00	14.0	0.0175	E	
J232011.5+081049	NGC 7619	3.78e–12	23 20 14.53 +08 12 22.4 ^{OR}	21.1	0.57	12.1	0.0127	E	
J232013.5+414643		6.85e–12	23 20 12.20 +41 46 05.3 ^O	9.7	0.20	(18.5)	0.1520 ¹		Early
			23 20 13.63 +41 46 19.1 ^O	9.7	0.45	(17.0)			
			23 20 14.14 +41 46 16.2 ^O	9.7	0.13	(18.1)			
J232111.3–121617		2.01e–12	23 21 11.36 –12 15 55.4	14.6 ^E	1.00	(19.2)			
J232125.9–231230 [†]		3.54e–12	23 21 26.93 –23 12 31.6 ^O	48.7 ^E	0.07	(17.5)			
	DUKST 536–007		23 21 26.34 –23 12 28.1 ^O	48.7 ^E	0.93	16.3		cD	
J232245.1–173852		4.69e–12	23 22 45.25 –17 38 55.6 ^O	238.1 ^D	1.00	(16.6)	0.1674 ¹		Sy1.5
J232352.2+421057	H 2321+419	8.48e–12	23 23 52.11 +42 11 00.6	11.4	1.00	17.0	0.0590		BLLAC
J232512.2–323637	PMN J2325–3236	6.63e–12	23 25 11.79 –32 36 34.5 ^O	253.0 ^D	1.00	17.0	0.2160		Sy1
J232519.4–120741	NPM1G –12.0625	2.35e–11	23 25 19.82 –12 07 28.6	1874.9	1.00	15.4	0.0822	Elliptical	
J232524.6–382653	LCRS B232242.2–384320	1.77e–11	23 25 24.15 –38 26 48.7	4.3	0.99	14.9	0.0359		Sy1
J232554.6+215310	RBS 2005	1.60e–11	23 25 54.25 +21 53 13.9 ^O	6.0 ^E	0.83	15.9	0.1202 ¹		Sy1
			23 25 54.90 +21 53 21.7 ^O	6.0 ^E	0.17	(18.2)			Sy1
J233315.8+225539		3.26e–12	23 33 15.59 +22 55 49.8 ^O	18.0 ^E	0.91	(17.7)			
J233340.1–252708		1.36e–12	23 33 39.49 –25 27 07.8	7.1	1.00	(19.2)			
J233355.5–234336	PGC 071756	6.36e–12	23 33 55.24 –23 43 40.0	782.5	1.00	16.5	0.0477	Elliptical	
J233500.0+272208	PGC 071807	3.17e–12	23 35 01.60 +27 22 18.0	78.3	1.00	13.5	0.0613	cD	
J233631.0+210848 [†]	IC 5338	1.23e–11	23 36 30.42 +21 08 45.1	56.1 ^E	1.00	15.0	0.0549	E?	
J233641.8+235526 [†]	MG3 J233645+2354	6.08e–12	23 36 42.12 +23 55 29.1 ^O	434.0 ^E	0.71	(16.4)	0.1267	cD	
			23 36 43.51 +23 55 26.6 ^O	434.0 ^E	0.28	(19.1)			
J233827.6+270045 [†]	NGC 7720 NED01	6.23e–12	23 38 29.40 +27 01 54.5 ^{OR}	6268.2 ^D	–	13.9	0.0309	E	
J233856.8+212432	87GB 233624.8+210840	7.63e–12	23 38 56.38 +21 24 41.3	45.8	1.00	17.0	0.2910		
J234050.2–284834	PKS 2338–290	1.56e–12	23 40 51.00 –28 48 41.6 ^O	501.6 ^D	1.00	19.0	0.4460	asym. dbl.	QSO
J234051.4+801513		1.15e–11	23 40 53.24 +80 15 14.7	48.9	0.99	(17.0)			
J234106.1+001836	PKS 2338+000	2.72e–12	23 41 06.94 +00 18 33.3	434.6	1.00	19.0	0.2766	Elliptical	
J234106.5+093805	NPM1G +09.0618	6.44e–12	23 41 06.65 +09 38 04.0	5.1	1.00	16.2	0.0421 ¹		Sy1.5
J234256.7–032227	[HB89] 2340–036	4.44e–12	23 42 56.68 –03 22 26.3	361.5 ^D	1.00	16.0	0.8960		QSO
J234312.3–363807	RBS 2040	2.74e–12	23 43 13.68 –36 37 55.7	10.3	1.00	16.0	0.6220		Sy1

Table 2—Continued

(1)	(2)	(3)	(4)		(5)	(6)	(7)	(8)	(9)	(10)
<i>ROSAT</i> BSC	NED	S _x	Position		S _{1.4GHz}	ID			NED	
1RXS name	Common name	ergs s ⁻¹ cm ⁻²	α	(J2000) δ	mJy	P	m _B	Redshift	Morphology	Sp. Class.
J234332.5+343957		1.41e-11	23 43 33.60	+34 39 51.2 ^O	34.9 ^E	0.14	(18.7)			
			23 43 33.87	+34 40 06.2 ^O	34.9 ^E	0.85	(21.5)			
J234538.0-144924	1ES 2343-151	1.95e-12	23 45 38.62	-14 49 29.3	10.7	1.00	19.2	0.2240		BLLAC
J234554.2+341419		7.47e-12	23 45 55.16	+34 14 03.6	28.8	1.00	(18.0)			
J234741.2-280829	IC 5358	2.91e-11	23 47 44.85	-28 08 22.9	89.6 ^S	—	13.6	0.0288	cD;E+4 pec	
J234907.7-331148	CTS 0120	4.91e-12	23 49 07.35	-33 11 46.6	5.0	1.00	16.9	0.1350		Sy1
J235013.4+015146		5.27e-12	23 50 13.75	+01 51 47.9	15.0 ^S	1.00	(21.5)			
J235018.0-055928	RBS 2051	6.79e-12	23 50 17.96	-05 59 29.4	25.1	1.00	19.5	0.5150		BLLAC
J235023.6-243552	[CM98] R843053	1.94e-12	23 50 23.41	-24 36 05.3	6.6	1.00	(16.8)	0.1930	cD?	
J235034.7+292924 [†]		3.68e-12	23 50 35.67	+29 29 40.4	66.8 ^E	1.00	(17.4)			
J235050.3+060921	MCG +01-60-039	1.23e-11	23 50 50.51	+06 08 58.2	56.3	1.00	15.0	0.0561		
J235140.7-260502		8.72e-12	23 51 39.37	-26 05 02.7	20.1	1.00	(17.1)			
J235156.2-010911	PG 2349-014	1.28e-11	23 51 56.21	-01 09 16.3	1596.8	1.00	15.3	0.1740	N galaxy	Sy1
J235321.1-145902		2.55e-12	23 53 20.97	-14 58 59.2	15.0	1.00	(21.5)			
J235525.6-335800	[HB89] 2352-342	1.61e-12	23 55 25.85	-33 57 57.2	331.4	1.00	16.4	0.7060	Opt.var.	QSO
J235555.1+143659		3.12e-12	23 55 55.07	+14 37 29.9 ^O	12.6	0.93	(21.5)			
J235700.0-344531	ESO 349- G 010	2.51e-11	23 57 00.67	-34 45 31.7	1286.3 ^E	1.00	13.9	0.0491	E+4;cD	LINER?
J235728.4-302739	AM 2354-304 NED02	7.21e-12	23 57 27.63	-30 27 47.2	5.1	1.00	15.0			
J235730.1-171801		2.13e-11	23 57 29.68	-17 18 05.3	44.4	1.00	(17.4)	0.3899? ¹		Sy?
J235908.0-303740	EUVE J2359-30.6	6.53e-11	23 59 07.56	-30 37 40.2	64.8	1.00	16.8	0.1650		BLLAC
J235910.6-040737	IC 1524	9.04e-12	23 59 10.82	-04 07 32.4	3.8	0.98	13.9	0.0193	SA(rs)b pec:	Sy1

[†]See notes.

^OOptical position given: because identification is more than 3σ from radio position or there are multiple optical candidates

^RNVSS position and/or flux density remeasured.

^FHigh resolution position and flux density taken from the FIRST 1.4 GHz survey (White et al. 1997).

^SExtended radio emission – central point source and single lobe; using central point source position and total emission.

^EExtended radio emission.

^DExtended radio emission – double; using centroid and total emission.

^TExtended radio emission – triple; using central source position and total emission.

¹Derived from optical spectra obtained with KPNO 2.1m telescope during 1997–1999. A dash (–) indicates there is not enough spectral information to classify

²From Brinkmann et al. 2000